Residual tRNA secondary structure in 'denaturing' 8M urea/TBE polyacrylamide gels: effects on electrophoretic mobility and dependency on prior chemical modification of the tRNA

Lisa A.Hegg and David L.Thurlow

Department of Chemistry, Clark University, 950 Main Street, Worcester, MA 01610, USA

Received December 4, 1989; Revised and Accepted February 20, 1990

ABSTRACT

Fifteen individual species of tRNA were treated with the chemical modifiers diethylpyrocarbonate, 50% aqueous hydrazine or hydrazine/3 M NaCl. Following purification of the chemically modified material on polyacrylamide gels containing 8 M urea, variant minor bands, in addition to the expected main band, were observed for 12 of the 15 tRNAs. Characterization of the content of chemically altered bases in material recovered from such bands indicated that tRNAs containing modified nucleotides in base-paired stems were excluded from the main band and present, often in enhanced amounts, in the minor variant bands. The persistence of residual secondary structure on 8 M urea gels run at 45°C and the ability of chemically modified bases to alter electrophoretic mobilities warrant caution in designing and interpreting experiments in which chemically modified RNA is isolated on gels prior to further analysis. tRNA^{Val}(VAC) was unique in that modified bases in non base-paired regions, according to the cloverleaf model of secondary structure, caused exclusion from the main band. Consequently, we propose a secondary structure for partially denatured tRNA^{Val}(VAC), in which these bases are located in double stranded regions of the molecule.

INTRODUCTION

The use of chemical probes including dimethylsulfate, diethylpyrocarbonate (DEP), kethoxal, 50% hydrazine, hydrazine/3 M NaCl, and others has contributed significantly to our understanding of RNA structure, both in solution (1,2) and in RNA-protein complexes (3–11). In most instances the chemically modified RNA was analyzed for its content of modified bases immediately following treatment with the modifying reagent, either by cleaving the RNA chain at chemically altered residues with aniline (12) or by identifying sites where reverse transcriptase pauses during primer extension (11).

However, in some cases the modified RNA was purified on polyacrylamide gels containing 8 M urea prior to characterizing

its content of modified bases (2,3,5–10). While examining the interaction between tRNAs and ATP/CTP:tRNA nucleotidyltransferase, we followed this procedure and observed that some tRNAs, containing certain chemically altered nucleotides, were not recovered from the main band of material on the urea-containing purification gel (13).

Retention of residual secondary structure on gels in 8 M urea is well documented, most frequently in the context of sequence analyses (14). Accordingly, it is not totally surprising that modified bases might interfere with residual structure and thereby alter the RNA's electrophoretic mobility. Because our observation warrants caution in designing and interpreting experiments of this sort, we have undertaken an extensive survey of tRNAs to determine how frequently this phenomenon is encountered. In the course of these studies we have obtained evidence that supports a model of secondary structure for tRNA^{Val}(VAC) under conditions of partial denaturation, which differs significantly from the cloverleaf model.

MATERIALS AND METHODS

Reagent grade chemicals were used to prepare all solutions. Aniline and anhydrous hydrazine were prepared by distillation as described by Perrin *et al.* (15). All aqueous solutions were prepared with deionized distilled water (Barnstead NANOpure II). Individually purified tRNAs were purchased from Boehringer Mannheim Biochemicals (Indianapolis, IN) or Subriden RNA (Rolling Bay, WA). Prior to treatment with chemical modifiers, the tRNAs, together with contaminating species, were labeled with ³²P and purified on a 20% polyacrylamide gel (acrylamide:N,N'-methylene-bisacrylamide=19:1) containing 8 M urea in 100 mM Tris-Borate, pH 8.3, 2.5 mM EDTA (TBE), run at 25 mA constant current until the xylene cyanole tracking dye had migrated 26 cm. The running temperature of such gels was normally ≈45°C.

[α -³²P]ATP (>3000 Ci/mmol) was obtained from ICN Radiochemicals (Irvine, CA). Snake venom phosphodiesterase (46 U/mg) was purchased from Worthington Biochemical (Freehold, NJ). The enzyme ATP/CTP:tRNA nucleotidyltransferase was prepared from 12 g of *Escherichia coli* cells

as follows. Frozen cells were ground with twice their weight of alumina in a chilled mortar and pestle. The slurry was resuspended in 100 mL 2mM Tris-Cl, pH 8.5, 10 mM MgCl₂, 0.1 mM EDTA, 2 mM DTT (Buffer A), centrifuged at 5000 rpm for 10 min in a Sorvall SS-34 rotor; the supernatant was centrifuged again at 16000 rpm for 30 min in the same rotor and at 48000 rpm for 90 min in a Beckman 55.2 Ti rotor. The final supernatant was loaded onto a 3.6 cm×13.5 cm column of Whatman (Hillsboro, OR) DE-52 DEAE cellulose; the column was resuspended in 250 mL Buffer A and protein was eluted with an exponential gradient of KCl (0–0.5 M) in Buffer A. Fractions representing the main peak of activity were concentrated to 0.4 mL using a Centricon-30 (Amicon, Danvers, MA) and an equal volume of glycerol was added. The homologous enzyme from *Saccharomyces cerevisiae* was kindly provided by Dr. Jacek Wower (University of Massachusetts, Amherst, MA) as the DEAE cellulose column eluant in the procedure of Sternbach *et al.* (16).

Removal of the 3'-terminal A, CA or CCA was carried out by digestion with snake venom phosphodiesterase (13). Missing cytosines were restored using 1 μ U nucleotidyltransferase (13) in 20 μ L 50 mM Tris-Glycine, pH 8.5, 5 mM MgCl₂, 8 mM dithiothreitol, 25 μ M CTP for 30 min at 37°C. Following extraction with an equal volume of Tris-EDTA saturated phenol, the RNA was precipitated with four volumes of ethanol, washed with 200 μ L 0.3 M sodium acetate, reprecipitated, rinsed with 500 μ L 95% ethanol, dried and resuspended in H₂O. Labeling with ³²P was accomplished by performing the same reaction in the presence of 50 μ Ci [α -³²P]ATP and in the absence CTP.

Chemical modifications of labeled tRNAs with DEP, 50% hydrazine or hydrazine/3 M NFAcI were performed according to Peattie (12) except the lengths of the incubations were shortened to 90 s, 5 min and 10 min respectively. Cleavage of the RNA at modified bases was accomplished by treatment with 1 M aniline-acetate as described by Peattie (12), except that the length of incubation at 65°C was shortened to 5 min. Resolution of aniline-induced fragments was carried out on 20% gels in TBE and 8 M urea (12). Autoradiograms were made using Kodak X-Omat AR film exposed at -70°C for 12–60 h.

RESULTS

Modified nucleotides in base-paired stems altered electrophoretic mobilities of tRNAs on 8M urea polyacrylamide gels

Fifteen ³²P-labeled tRNAs were purified to homogeneity on 20% polyacrylamide gels in TBE/8 M urea, treated with either DEP, 50% hydrazine or hydrazine/3 M NaCl (12), and then fractionated on another 8 M urea-containing gel. The results obtained with seven of these 15 tRNAs are shown in Figure 1. Most tRNAs treated in this fashion migrated either as a single band (Fig. 1A,B), or sometimes as a main band, accompanied by one or more minor variant bands (Fig. 1C–G). Modification-induced variant bands had greater electrophoretic mobilities than those of the controls, were usually recovered in low yields, less than 10% of main band, and often exhibited variable electrophoretic mobilities depending on the nature of the chemical modification (Fig. 1D,E). Occasionally, some material was lost during treatment with the chemical modifiers, possibly due to inefficient precipitations with ethanol (Fig. 1G, lanes 2 and 4).

Material from the main band of the 20% gels, such as those shown in Figure 1, was recovered and treated with aniline to

Figure 1. Autoradiograms of purification gels showing main bands and some minor electrophoretic variants of tRNAs labeled with ³²P at their 3'-ends. tRNA^{Asn} (A), tRNA^{Gln}(NUG) (B), tRNA^{Asp} (C), tRNA^{Ile} (D), tRNA^{Lys} (E), tRNA^{Gln}(CUG) (F), and tRNA^{Val} (G) were purified on 20% polyacrylamide gels in TBE/8 M urea following no treatment (lanes 1), treatment with DEP (lanes 2), 50% hydrazine (lanes 3) or hydrazine/3 M NaCl (lanes 4). Modification-induced electrophoretic variants are designated by arrows. Note that the upper band in B, present in the unmodified control, is a contaminating species (tRNA^{Gly}), and is not a modification-induced variant. Variant bands of tRNA^{Asp} (V1 and V2) were recovered and analyzed for their content of modified bases.

cleave the RNA chain at modified bases (12). Labeled fragments generated in this manner were resolved on sequencing gels and those that were absent, or present at significantly reduced intensities relative to non-gel purified material, were identified (Fig. 2). We refer to this absence of tRNAs containing a particular modified nucleotide as an 'exclusion,' and note that modification of any of several different nucleotides could cause a tRNA to be excluded from the main band.

A summary of nucleotides responsible for exclusions from the main band of fifteen repurified tRNAs and their general locations within the tRNA's secondary structure are presented in Table 1. Three tRNAs did not exhibit any obvious exclusions following treatment with DEP, 50% hydrazine, or hydrazine/3 M NaCl (Table 1). For example, no aniline-induced fragments of tRNA^{Met} were absent from the gel purified material (Fig. 2A). All other 12 tRNAs examined underwent some exclusion from the main band following treatment with at least one of the chemical modifiers, and some were excluded following treatment with each reagent (e.g. tRNA^{Arg}, Fig. 2B). DEP-modified bases were frequently responsible for exclusions (Table 1, Fig. 2C–F).

Locations of bases responsible for exclusions varied among tRNAs, but were almost always within a base-paired region of the cloverleaf model, usually within the Ψ - or anticodon stem. We cannot rule out the possibility the additional exclusions may have resulted from modified bases near the 5'-end in the acceptor

Figure 2. Exclusions resulting from chemical modification. ^{32}P -labeled tRNA^{Met} (A), tRNA^{Arg} (B), tRNA^{Gln}(NUG) (C), tRNA^{Ser} (D), tRNA^{Gly} (E), and tRNA^{Met} (F) were chemically modified, isolated from the main band on a 20% purification gel in TBE/8 M urea, and treated with aniline. The resulting fragments were separated on sequencing gels run for 2 h (A-D) or 5 h (E,F), which were used to prepare autoradiograms. Control material (lanes 1,3 and 5) was modified, but not gel purified. Modifying reagents were DEP (lanes 1 and 2), 50% hydrazine (lanes 3 and 4) or hydrazine/3 M NaCl (lanes 5 and 6). Nucleotide positions (numbered according to Ref. 21) responsible for exclusions are indicated with arrows.

stem, because fragments of that length were not always fully resolved on our gels. Therefore, the data presented in Table 1 represents a minimum set of nucleotides that, when chemically modified, caused exclusion of the tRNA containing it from the main band of a purification gel.

tRNAs enriched in modified bases were recovered in electrophoretic variant bands

Because the yields of variant bands were generally low, we were able to fully characterize their content of modified bases in only three instances: tRNA^{Asp}, tRNA^{Val} and tRNA^{Gln}(CUG). As discussed above, tRNAs containing certain modified nucleotides were excluded from the main band of the purification gel. However, tRNAs in the modification-induced bands were often

enriched in their content of such modified bases (Figs. 3–5). We refer to this increased recovery as an ‘enhancement,’ and note that modification of any of several different nucleotides in each tRNA could cause this phenomenon.

Where more than one variant band was evident, sites causing enhancements differed from one modification-induced variant band to another. For example, bases responsible for enhancements in the slowest migrating variant band in tRNA^{Asp} (band V1, Fig. 1C) were located primarily in the anticodon stem, positions 27–33 and 39–44. In contrast, nucleotides causing enhancements in the second variant (band V2, Fig. 1C) were in the Ψ -stem, sites 50–53 and 61–64 (Figs. 3 and 6A).

In tRNA^{Gln}(CUG) and tRNA^{Val}(VAC), only one variant band, recovered in significant yield, was evident on the purification gel (Fig. 1F,G). In one case, tRNA^{Val}(VAC) treated with

Table 1. Summary of nucleotides responsible for exclusions and their locations within the cloverleaf model of tRNA.

tRNA	Nucleotide Positions	Position in Cloverleaf Model
Asn _{GUU}	None	
Met _{CAU}	None	
Phe _{GAA} (yeast)	None	
Ser _{UGA}	47G–47J, 46–47, 47A–47C	Variable stem
Arg _{ACG}	65–61, 53–49	Ψ stem
Gln _{UUG}	65–62, 53–49, 43–39, 31–27	Ψ and anticodon stems
Gln _{CUG}	64–61, 53–50, 43–40, 33, 31–27	Ψ and anticodon stems
Asp _{GUC}	64–61, 58, 53–50, 44–39, 33–27	Ψ and anticodon stems
		Ψ and anticodon loops
Lys _{UUU}	66–61, 53–49	Ψ and acceptor stems
Gly _{GCC}	44–38, 33, 31–27	Anticodon stem and loop
Met _t	41–39, 31–29	Anticodon stem
Phe _{GAA}	72–67, 44, 31–27, 7	Anticodon and Acceptor stems
Ile _{NAU}	74–67, 2, 1	Acceptor stem
Trp _{CCA}	72–67, 7–1	Acceptor stem
Val _{VAC}	59–57, 49, 47–43, 33, 22–18	Ψ, D, variable, and anticodon loops

All tRNAs were isolated from *E. coli* unless otherwise specified. Nucleotide positions are numbered according to Sprinzl *et al.* (21).

hydrazine/3 M NaCl, only the modification-dependent band was evident (Fig. 1G, lane 4).

Positions of nucleotides within the secondary structure of tRNA^{Gln}(CUG) that gave rise to enhancements in the variant band were exactly the same as those responsible for exclusions from the main band (Table 1, Fig. 4). Accordingly, they were generally within base-paired regions of the cloverleaf model (Fig. 6B).

Positions in tRNA^{Val}(VAC) responsible for enhancements also corresponded to those causing exclusions (Table 1, Fig. 5). Unlike all 13 other tRNAs examined, bases responsible for exclusions and enhancements were located primarily in single-stranded portions of tRNA^{Val}, according to the cloverleaf model (Table 1, Fig 6C). In fact, they were found within each of the cloverleaf's single-stranded loops, suggesting the possibility that, following treatment with modifying reagents, the tRNA had assumed a secondary structure which differed from the cloverleaf model and which persisted throughout isolation on the purification gel.

DISCUSSION

Alteration of electrophoretic mobility by chemically modified nucleotides

Chemically modified bases, which were responsible for exclusion of tRNAs from the main band of our purification gels (Table 1, Fig. 6A,B), were generally located in double stranded regions of the molecule. This observation suggested that significant secondary structure had persisted during electrophoresis on our 8 M urea gels. We assume that the presence of 8 M urea at our running temperature of 45°C was sufficient to melt out at least the tRNA's tertiary interactions and probably much of the D-stem as well (17,18). In accord with these assumptions, no chemically altered bases involved in tertiary interactions or base pairing in the D-stem were observed to bring about exclusions or enhancements (Table 1), except for tRNA^{Val} (see later in Discussion).

Retention of significant secondary structure on 8 M urea gels has been noted previously in the context of 'band compressions' during sequence analyses (14). Running gels at elevated temperatures (50–60°C) is often sufficient to prevent such

artifacts, but even higher temperatures (80–90°C, with heat resistant or Pyrex plates) may be required (14). Since our gels were run at 45°C, it is likely that tightly base-paired regions had resisted denaturation.

Accordingly, whenever a modified nucleotide interfered with such residual structure, the electrophoretic mobility of the tRNA containing that base could be altered. As a result, that tRNA would not be recovered in the main band of material retaining partial secondary structure, and its aniline-induced fragments would not be observed on sequencing gels (Fig. 2).

Identification of modified bases in material derived from variant bands supports the above interpretation, in that modified bases resulting in exclusion of tRNAs from the main electrophoretic band were simultaneously responsible for enhanced recovery of tRNAs containing them in the lower band. Where more than one variant band was recovered, different regions of secondary structure may have persisted in each. Consistent with this notion, enhancements occurred throughout the anticodon stem in the slowest migrating variant of tRNA^{Asp} (band V1, Fig. 1C) and throughout the Ψ-stem in the faster migrating band (band V2, Fig. 1C, see Figs. 3 and 6A).

In instances where only one additional band resulted from modification (Fig. 1F,G), variant tRNAs may have been devoid of any residual secondary structure. This would follow provided that fully denatured tRNA migrated the same, regardless of the site, or sites, of modification responsible for its inability to retain significant secondary structure. Our data do not explain why the fully denatured form migrated faster than that retaining secondary structure. However, we did use high percentage (20%) polyacrylamide gels for separation, and effects resulting from an increased diameter of base paired regions might bring about a reduced mobility in gels with extremely small pores.

Where mobilities of variant bands varied depending on the nature of the modifying reagent (Fig. 1D,E), differences can be explained by the fact that each modifier used in this study induced vastly different alterations in chemical structure (12).

Material that had been gel purified often displayed an increased reactivity towards treatment with aniline. Frequently, bands corresponding to purines, especially guanosines, were more intense when the sample had been modified with DEP and then isolated on a gel (Fig. 2A and 2E, lanes 2), relative to those in

Figure 3. Exclusions and enhancements resulting from chemical modification of tRNA^{Asp} labeled with ³²P at its 3'-end. Autoradiograms of sequencing gels run for 2 h (A) or 5 h (B) depict separation of aniline-induced fragments. tRNA^{Asp}, modified with DEP (lanes 1–4), 50% hydrazine (lanes 5–7), or hydrazine/3 M NaCl (lanes 8–11), was isolated from a main band and additional variant bands (V1 and V2, Fig. 1C). Following treatment with aniline, fragments were analyzed on sequencing gels as follows: main band, lanes 2,6 and 9; variant V1, lanes 3,7 and 10; variant V2, lanes 4 and 11. Control samples not purified on a gel were analyzed in a similar manner (lanes 1,5 and 8). Nucleotide positions responsible for exclusions from the main band and enhancements in V1 (arrows) or V2 (open circles) are indicated and numbered according to Ref. 21.

material that had not been gel purified following the chemical modification (Fig. 2A and 2E, lanes 1). Alternatively, gel purified material exhibited an increased specificity towards the aniline reaction. Aniline-induced cleavage of samples treated with hydrazine/3 M NaCl and then purified on a gel was much more specific for cytidines than that of non gel purified material (Fig. 2A, compare lane 6 to lane 5). We do not know the basis for the increased reactivity or specificity. However, these phenomena did not interfere with identification of exclusions, because exclusions were characterized by the **absence** of bands in gel purified material; nor did they prevent identification of enhancements, because enhancements were detected by comparisons between samples that had both been purified on gels.

Our observations warrant caution in designing and interpreting experiments involving chemical modification of RNA

The observation that chemically modified tRNAs can migrate differently on 8 M urea gels is important to note in the context of any experiment in which RNA is treated with a modifying reagent and subsequently gel purified. In our lab we have conducted an extensive study of nucleotides required in chemically unaltered form for recognition by the enzyme ATP/CTP:tRNA nucleotidyltransferase (13) in which we utilized such an approach. It was during the course of these damage-selection studies that we first observed exclusions due to gel

Figure 4. Enhancements resulting from chemical modification of tRNA^{Gln}(CUG) labeled with ³²P at its 3'-end. Autoradiograms of sequencing gels run for 2 h (A) or 5 h (B) depict separation of aniline-induced fragments. tRNA^{Gln}(CUG), modified with DEP (lanes 2 and 3), 50% hydrazine (lanes 4 and 5), or hydrazine/3 M NaCl (lanes 6 and 7), was isolated from the main band (lanes 2,4 or 6) or a single additional variant band (lanes 3,5 or 7) (see Fig. 1F) prior to treatment with aniline. Unmodified control material (lanes 1) was gel purified and treated with aniline. Nucleotide positions responsible for enhancements are indicated by arrows and are numbered according to Ref. 21.

purification and, accordingly, have included gel purified controls in our experimental design.

The existence of electrophoretic conformers resulting from treatment with the chemical probe kethoxal has been reported previously in studies examining the conformation of tRNA^{Phe} bound to the ribosome (5). In this instance, the modified tRNA was purified on 12% gels in 8 M urea at pH 7.5, digested with nuclease and then analyzed on 12% gels in 8 M urea at pH 8.3. It was only on the gel at pH 8.3 that the variants were observed. Therefore the two conformers were never individually purified and analyzed for content of modified bases separately. Thus we note that gel purification itself does not necessarily lead to exclusion, unless the conditions of electrophoresis are such that conformers are sufficiently resolved.

Our gels were relatively high in acrylamide content (20%) and were run for extended periods (xylene cyanole tracking dye 26 cm from wells) so as to maximize separation of different species of tRNAs. Consequently, the phenomenon of exclusion was readily

apparent under our conditions. Since one cannot predict *a priori* whether a given set of electrophoresis conditions will lead to resolution of conformers, we suggest that care should be taken in designing experiments to control for the possibility of modification induced electrophoretic conformers.

It is also important to note that previous studies, such as those on tRNA-EFTu (3,6) and tRNA-ribosome (5,6) interactions, and on ribosomal RNA structure in solution (2) and in ribosomes or reconstituted complexes (7–10), utilized purification of chemically modified tRNAs on 8 M urea gels. We cannot say for certain whether any exclusions occurred in these or similar experiments, but suggest that interpretations should be qualified in light of our observations.

Secondary structure for partially denatured tRNA^{Val}(VAC)

Nucleotides responsible for exclusions and enhancements of tRNA^{Val} were located primarily in single-stranded regions of the cloverleaf model. This observation suggested that tRNA^{Val} may

Figure 5. Enhancements resulting from chemical modification of tRNA^{Val} labeled with ³²P at its 3'-end. Autoradiograms of sequencing gels run for 2 h (A) or 5 h (B) depict separation of aniline-induced fragments. tRNA^{Val}, modified with DEP (lanes 2 and 3), 50% hydrazine (lanes 4 and 5), or hydrazine/3 M NaCl (lane 6), was isolated from the main band (lanes 2 and 4) or a single additional variant band (lanes 3, 5 and 6) (see Fig. 1G) prior to treatment with aniline. Unmodified control material (lanes 1) was gel purified and treated with aniline. Nucleotide positions responsible for enhancements are indicated by arrows and are numbered according to Ref. 21.

adopt a secondary structure under conditions of partial denaturation that differs from the cloverleaf model. A structure is proposed (Fig. 7) in which virtually all bases that give rise to exclusions or enhancements are located in double-stranded portions of the molecule.

We note that exclusions or enhancements resulting from modification of base-paired cytidines could not be determined in this instance, because only one band was recovered on the purification gel, and it migrated differently than unmodified tRNA (Fig. 1G). Because the modification was carried out such that each tRNA was modified at only one residue, alteration of any one cytidine must have been sufficient to prevent residual structure in tRNA^{Val}(VAC).

The predicted thermodynamic stability (19) of our proposed

structure (−15.9 kcal/mol) is slightly greater than that of a tRNA folded according to the cloverleaf model without the D-stem (−14.1 kcal/mol). Tertiary base pairs and D-stem base pairs were omitted from the calculations for the cloverleaf model, since previous studies have shown that they were the most easily disrupted by thermal melting in the absence of Mg²⁺ (17,18) and would therefore be unlikely to persist under our conditions of electrophoresis. For the same reasons we did not include the stem formed by base pairing between positions 11–13 and 40–42 in the partially denatured structure (dotted lines, Fig. 7).

Evidence for multiple conformations of tRNA^{Val}(VAC) induced by thermal denaturation has previously been reported based on NMR spectra (20). In this set of experiments the tRNA's native structure underwent a transition with a T_m of 51°C in the

Figure 6. Sites where modified bases caused exclusions and enhancements in tRNA^{Asp} (A), tRNA^{Gln}(CUG) (B), and tRNA^{Val} (C). Nucleotides responsible for exclusions and enhancements are indicated by bold letters on cloverleaf models of secondary structure. For tRNA^{Asp} (A) arrows indicate bases responsible for enhancements (Fig. 3) in band V2 of the purification gel (Fig. 1C), and bases in bold without arrows indicate those that brought about enhancements (Fig. 3) in band V1 (Fig. 1C).

Figure 7. Secondary structural model for partially denatured tRNA^{Val}. Sites responsible for exclusions and enhancements are shown in bold.

absence of Mg²⁺. The enthalpy for this transition was larger than expected for simple disruption of tertiary interactions (20), and based on our data may reflect transition to a structure such as the one we propose (Fig. 7).

Induction of such a conformation by thermal melting, chemical modification, or any other *in vitro* treatment does not signify that this structure is physiologically relevant. However, it does serve to point out the dynamic nature of nucleic acid structure in solution and the care one must take in designing and interpreting structural studies.

REFERENCES

1. Peattie, D.A. and Gilbert, W. (1980) *Proc. Natl. Acad. Sci. USA*, 77, 4679-4682.
2. Van Stolk, B.J. and Noller, H.F. (1984) *J. Mol. Biol.*, 180, 151-177.
3. Bertram, S. and Wagner, R. (1982) *Biochemistry International*, 4, 117-126.
4. Peattie, D.A. and Herr, W. (1981) *Proc. Natl. Acad. Sci. USA*, 78, 2273-2277.
5. Bertram, S., Göringer, U. and Wagner, R. (1983) *Nucleic Acids Res.*, 11, 575-589.
6. Douthwaite, S., Garrett, R.A. and Wagner, R. (1983) *Eur. J. Biochem.*, 131, 261-269.
7. Peattie, D.A., Douthwaite, S., Garrett, R.A. and Noller, H.F. (1981) *Proc. Natl. Acad. Sci. USA*, 78, 7331-7335.
8. Lo, A.C. and Nazar, R.N. (1981) *FEBS Lett.*, 131, 41-44.
9. Wildeman, A.G. and Nazar, R.N. (1982) *Eur. J. Biochem.*, 121, 357-363.
10. Thurlow, D.L., Ehresmann, C. and Ehresmann, B. (1983) *Nucleic Acids Res.*, 11, 6787-6802.
11. Moazed, D., Stern, S. and Noller, H.F. (1986) *J. Mol. Biol.*, 187, 399-416.
12. Peattie, D.A. (1979) *Proc. Natl. Acad. Sci. USA*, 76, 1760-1764.
13. Spacciapoli, P., Doviken, L., Mulero, J.J. and Thurlow, D.L. (1989) *J. Biol. Chem.*, 264, 3799-3805.
14. Deininger, P.L. (1983) *Anal. Biochem.*, 135, 247-263.
15. Perrin, D.D., Armarego, W.L.F., and Perrin, D.R. (1980) *Purification of Laboratory Chemicals*. Pergamon Press, Oxford, England.
16. Sternbach, H., von der Haar, F., Schlimme, E., Gaertner, E. and Cramer, F. (1971) *Eur. J. Biochem.*, 22, 166-172.
17. Crothers, D.M., Cole, P.E., Hilbers, C.W. and Shulman, R.G. (1974) *J. Mol. Biol.*, 87, 63-88.
18. Johnston, P.D. and Redfield, A.G. (1981) *Biochemistry*, 20, 3996-4006.
19. Turner, D.H., Sugimoto, N., Jaeger, J.A., Longfellow, C.E., Freier, S.M. and Kierzek, R. (1987) *Cold Spring Harbor Symp. Quant. Biol.*, 52, 123-133.
20. Kastrup, R.V. and Schmidt, P.G. (1978) *Nucleic Acids Res.*, 5, 257-269.
21. Sprinzl, M., Hartmann, T., Meissner, F., Moll, J. and Vorderwülbecke, T. (1987) *Nucleic Acids Res.*, 15, Supplement, r53-r188.