Published in final edited form as:

Science. 2011 January 14; 331(6014): 220–223. doi:10.1126/science.1197243.

Light-driven changes in energy metabolism directly entrain the cyanobacterial circadian oscillator

Michael J. Rust^{1,2}, Susan S. Golden³, and Erin K. O'Shea¹

¹Howard Hughes Medical Institute, Harvard University Center for Systems Biology, Departments of Molecular and Cellular Biology and Chemistry and Chemical Biology, 52 Oxford Street, Cambridge MA 02138 USA

³Center for Chronobiology, Division of Biological Sciences, University of California San Diego, 9500 Gilman Drive, La Jolla CA 92093 USA

Abstract

Circadian clocks are self-sustained biological oscillators that can be entrained by environmental cues. Though this phenomenon has been studied in many organisms, the molecular mechanisms of entrainment remain unclear. Three cyanobacterial proteins and ATP are sufficient to generate oscillations in phosphorylation *in vitro*. We show that changes in illumination that induce a phase shift in cultured cyanobacteria also cause changes in the ATP/ADP ratio. When these nucleotide changes are simulated in the *in vitro* oscillator, they cause phase shifts similar to those observed *in vivo*. Physiological concentrations of ADP inhibit kinase activity in the oscillator and a mathematical model constrained by data shows that this effect is sufficient to quantitatively explain entrainment of the cyanobacterial circadian clock.

Circadian clocks, based on self-sustained biological oscillators with a ~24 hour period, allow organisms to anticipate regular variations in the environment caused by the Earth's rotation (1). For a circadian system to be most useful to the organism, it must function robustly in the face of fluctuations in the environment but also accept appropriate input signals to stay entrained with the true diurnal cycle. Therefore, a fundamental problem is to elucidate the molecular mechanisms that allow faithful transduction of timing information into the oscillator.

The cyanobacterium *Synechococcus elongatus* PCC 7942 has a core circadian oscillator that can be reconstituted *in vitro* from three purified proteins: KaiA, KaiB and KaiC (2). KaiC is an enzyme that catalyzes the phosphorylation and dephosphorylation of two of its own residues in an ordered pattern (3). KaiA and KaiB work together to modulate the activity of KaiC in a phosphorylation-dependent manner, resulting in stable oscillations in the amount of phosphorylated KaiC (4–6). We sought to identify light-dependent clock input pathways in living cyanobacteria and to study their mechanism in the reconstituted *in vitro* oscillator.

Causal links between metabolic activity and circadian clocks have previously been described. The circadian clock in the rat liver can be entrained by feeding (7), and adenosine monophosphate-activated protein kinase has been implicated in clock function in liver tissue (8). A classical result connecting clocks and metabolism is Aschoff's Rule—the observation

²Current Address: Department of Molecular Genetics and Cell Biology and the Institute for Genomics and Systems Biology, The University of Chicago, 900 East 57th Street, Chicago IL 60637 USA

Light-induced changes in the adenine nucleotide pool directly affect the cyanobacterial circadian oscillator to produce phase shifts.

Supporting Online Material www.sciencemag.org

that varying the light intensity tends to have opposite effects on the circadian clocks of nocturnal versus diurnal organisms (9).

We reasoned that the link to metabolism might be especially important in photoautotrophic cyanobacteria, as these microorganisms are entirely reliant on photosynthesis to extract energy from the environment (10). Darkness elicits profound changes in the physiology of *S. elongatus*, beyond the direct effects on photosynthesis. When cultures are removed from the light, global transcription and translation rates drop and there are large changes in the superhelicity of endogenous DNA (11, 12).

Cyanobacteria growing in the light primarily synthesize ATP by photophosphorylating ADP at the thylakoid membrane (13), and changes in culture illumination have been reported to affect the relative abundance of adenine nucleotides (14–16). We therefore tested whether incubating synchronized cultures in the dark under conditions sufficient to cause a phase shift in the circadian clock would result in changes in the concentrations of ATP and ADP. We monitored changes induced by a dark pulse in the circadian rhythm of an engineered strain of *S. elongatus* with a bioluminescent reporter of clock-driven transcription (17). In agreement with previous studies, an 8-hour dark pulse applied during the subjective day (Fig. 1A) induced a phase shift in the circadian rhythm, and the clock was refractory to a dark pulse applied during the subjective night (Fig. 1B) (18–20). These phase shifts in the transcriptional reporter were mirrored by shifts in the rhythm of KaiC phosphorylation, as measured by immunoblotting (Fig. S1). Irrespective of when the dark pulse was applied, these cultures experienced large changes in their adenine nucleotide pool (Fig. 1C). After 2 to 3 hours in the dark, the ratio of ATP to ADP + ATP had fallen to nearly 50% and remained low for the duration of the dark pulse; when illumination was restored, this ratio returned to ~85% within an hour.

The mechanism of the KaiABC protein oscillator relies on ordered multisite phosphorylation of KaiC (4, 5). Because ADP cannot serve as the phosphoryl donor in the kinase reaction, physiological changes in the relative amounts of ATP and ADP caused by changes in light have to the potential to modulate the activity of enzymes that consume ATP, and might directly affect the function of the circadian clock (21). To test this, we developed an experimental system to mimic *in vitro* the changes in nucleotide ratio we had observed in living cyanobacteria. We initiated oscillator reactions with purified KaiA, KaiB and KaiC in a buffer containing an excess of phosphoenolpyruvate and 4 mM ATP, similar to estimates of adenine nucleotide concentration in cyanobacterial cells in the millimolar range (22). After the oscillation had been established, we simulated the metabolic effects of darkness by adding ADP to bring the ratio of ATP to ADP + ATP to ~50%. To then simulate the effects of a return to growth in light, we added pyruvate kinase to convert the ADP to ATP, a reaction which went to completion in ~1 hour (Fig. S2). Because the ATP/ADP ratio *in vivo* initially falls gradually when cells are incubated in the dark for 8 hours in our liquid culture conditions (Fig. 1C), we used a 5 hour step-like pulse of ADP *in vitro* to approximate the amount of time that the *in vivo* cultures experience the lowest ATP/ADP ratio.

Transient manipulation of the ratio of ATP to ADP in the KaiABC oscillator created large phase shifts in the phosphorylation rhythm (Fig. 2A). Further, the phase response curve obtained by altering the ATP/ADP ratio *in vitro* was similar to that observed in live cells treated with pulses of darkness—the oscillator was most sensitive during the middle of subjective day (when KaiC phosphorylation was increasing) and nearly insensitive during subjective night (when KaiC phosphorylation was decreasing) (Fig. 2B–C) (18, 19, 23). During the pulse of increased ADP, KaiC phosphorylation was decreased relative to that in the control reaction, similar to changes in KaiC phosphorylation *in vivo* when cells are subjected to a dark pulse (Fig. S1) (24).

To determine the increase in the relative amount of ADP needed to produce this effect, we added varying amounts of ADP to several identical KaiABC reactions during a fixed portion of the oscillator cycle. The induced phase shift was a graded function of the amount of ADP added (Fig. 2D–E). Similarly, the magnitude of the phase shift decreased when the length of the pulse of ADP was decreased (Fig. S3). Thus, even small changes in the ATP/ADP ratio, smaller or shorter in duration than those induced by a rapid transition to prolonged darkness, adjusted the time specified by the circadian clock.

To investigate the molecular mechanism by which changes in the relative fraction of ADP are able to alter the oscillator phase, we studied the effect of the ATP/ADP ratio on non-oscillatory partial reactions in which one of the Kai proteins was absent. In the KaiA-KaiC reaction, KaiA chronically stimulates KaiC's autophosphorylation activity resulting in a steady state in which KaiC is highly phosphorylated (5, 25). Increasing the relative fraction of ADP present in the reaction resulted in both lower steady-state phosphorylation and a slower approach to that steady state (Fig. 3A). In the KaiB-KaiC reaction, KaiC has very weak kinase activity and the fraction of phosphorylated KaiC slowly decreases, allowing us to isolate the dephosphorylation reaction (5). We did not detect any effect of varying the ATP/ADP ratio on KaiC's phosphatase activity (Fig. 3B), or on the stability of KaiB-KaiC complexes (Fig. S4).

Quantitative analysis of the kinetics of the partial reactions confirmed the above observations and was consistent with a model in which ADP acts as a competitive inhibitor of KaiC's kinase activity. In fitting the phosphorylation kinetics to an initial velocity approximation, we estimated an effective relative affinity for ATP and ADP that is close to unity ($K_{rel} = 0.76 \pm 0.15$) (Fig. 3C). This value is compatible with two simple mechanistic hypotheses: either the true binding affinities for ATP and ADP are comparable in the CII lobe [the seat of KaiC autophosphorylation (26)] or the rate of ATP hydrolysis in CII is much faster than the rate of nucleotide dissociation, so that any difference in affinity is negligible. In either case, ADP acts in proportion to its relative abundance to decrease the KaiC phosphorylation rate.

To test whether the competitive inhibition mechanism we had identified was sufficient to explain the large phase shifts in the full oscillating reaction, we turned to a mathematical model of the circadian clock based on experimental measurements of the rates of phosphorylation and dephosphorylation at two sites on KaiC: S431 and T432 (5). In this model, stable oscillations arise because KaiC phosphorylated only on S431 interacts strongly with KaiB, trapping KaiA stoichiometrically and globally promoting dephosphorylation. In this sense, KaiC phosphorylated on S431 is autocatalytic, generating a nonlinearity that drives the oscillator.

We modified this model so that, in accord with our experimental observations, the kinase rates were modulated by the ratio of ATP to ADP + ATP present in the reaction, as well as the level of KaiA activity (Fig. 4A) (5). Thus, at each point in the oscillation, KaiC integrates clock information (through KaiA's kinase-stimulating activity) with metabolic information (through the adenosine pool). We simulated our *in vitro* experiments within this model by transiently lowering the ratio of ATP to ADP + ATP from 100% to 50% at various times in the circadian cycle. In simulation, pulses of competitive inhibition by ADP were sufficient to generate large phase shifts in the oscillator when these pulses occurred during the phase of the cycle when KaiC phosphorylation was rising (Fig. 4B). In order to obtain a strong resetting effect, KaiC must have sufficiently high sensitivity to ADP, at the upper end of the range of effective affinities we estimated experimentally (Fig. S5). We used this model to predict a full phase response curve based on this competitive inhibition mechanism and compared it to our experimental data and to measurements of the phase shift induced by

darkness *in vivo* (Fig. 4C) (19). Thus, varying relative nucleotide concentrations in the reconstituted oscillator approximates the response of the circadian clock in living cyanobacteria to changes in illumination, and an increase in the amount of ADP appears to alter the phase of the oscillator through inhibition of KaiC's kinase activity.

In the model, the ATP/ADP ratio describes a family of limit cycles that differ in the amplitude of the phosphorylation rhythm. Indeed, KaiABC reactions in buffers with tonically lowered ATP/ADP continue to oscillate with a circadian period, but KaiC cycles through distinct patterns of phosphorylated states (Fig. S6). If the ATP/ADP ratio is lowered abruptly, phosphorylation is inhibited, and the system must adjust to a new limit cycle. If this transition produces amounts of S431-phosphorylated KaiC sufficient to block KaiA activity, a large phase shift can result.

We have described a simple entrainment mechanism for a circadian clock in which enzymatic activity is directly tied to the availability of biochemical energy in the cell. In this view, no signaling pathway that specifically targets the oscillator is required to couple the clock to the environment, and the core oscillator proteins are sensitive to the metabolic state of the cell (27). Although factors outside the Kai proteins have been implicated in light-driven input to the cyanobacterial circadian system, including LpdA and the histidine kinase CikA (18, 20), strains lacking these proteins can be effectively entrained with repeated light-dark cycles, supporting the hypothesis that there exist basic synchronization mechanisms intrinsic to the KaiABC core oscillator itself. Because unexpected darkness will unavoidably lead to changes in the production and consumption of ATP in an obligate phototroph, KaiC's sensitivity to ADP represents a robust intrinsic mechanism for maintaining synchrony with the environment.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

29. We thank G. Dong for advice on culture conditions and K. Cook, V. Denic, Q. Justman, J. Markson, A. Rizvi and V. Vijayan for valuable discussions and comments on the manuscript. E. K. O. is an investigator of the Howard Hughes Medical Institute. NIH grant GM62419 supported work in the lab of S.S.G. M. J. R. was supported by a Helen Hay Whitney Foundation postdoctoral fellowship and by a Burroughs-Wellcome Foundation Career Award at the Scientific Interface.

References and Notes

1. Roenneberg T, Foster RG. *Photochem Photobiol.* Nov.1997 66:549. [PubMed: 9383985]
2. Nakajima M. *Science.* Apr 15.2005 308:414. [PubMed: 15831759]
3. Xu Y. *Proc Natl Acad Sci U S A.* Sep 21.2004 101:13933. [PubMed: 15347809]
4. Nishiwaki T. *EMBO J.* Sep 5.2007 26:4029. [PubMed: 17717528]
5. Rust MJ, Markson JS, Lane WS, Fisher DS, O'Shea EK. *Science.* Nov 2.2007 318:809. [PubMed: 17916691]
6. Qin X. *Proc Natl Acad Sci U S A.* Aug 2.2010
7. Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M. *Science.* Jan 19.2001 291:490. [PubMed: 11161204]
8. Lamia KA. *Science.* Oct 16.2009 326:437. [PubMed: 19833968]
9. Aschoff J. *Cold Spring Harb Symp Quant Biol.* 1960; 25:11. [PubMed: 13684695]
10. Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. *J General Microbiol.* 1979; 111:1.
11. Ito H. *Proc Natl Acad Sci U S A.* Aug 18.2009 106:14168. [PubMed: 19666549]

12. Woelfle MA, Xu Y, Qin X, Johnson CH. *Proc Natl Acad Sci U S A*. Nov 20.2007 104:18819. [PubMed: 18000054]
13. Scherer S, Almon H, Boger P. *Photosynthesis research*. 1988; 15:95. 1988.
14. Bornefeld T, Simonis W. *Planta*. 1974; 115:309. 1974.
15. Kallas T, Castenholz RW. *J Bacteriol*. Jan.1982 149:229. [PubMed: 6798019]
16. Lubberding HJ, Schroten W. *FEMS Microbiology Letters*. 1984; 22:93. 1984.
17. Andersson CR. *Methods Enzymol*. 2000; 305:527. [PubMed: 10812624]
18. Schmitz O, Katayama M, Williams SB, Kondo T, Golden SS. *Science*. Aug 4.2000 289:765. [PubMed: 10926536]
19. Kiyohara YB, Katayama M, Kondo T. *J Bacteriol*. Apr.2005 187:2559. [PubMed: 15805501]
20. Katayama M, Kondo T, Xiong J, Golden SS. *J Bacteriol*. Feb.2003 185:1415. [PubMed: 12562813]
21. Atkinson D. *Biochemistry*. November.1968 7:4030. 1968. [PubMed: 4972613]
22. Ihlenfeldt MJ, Gibson J. *Arch Microbiol*. 1975; 102:13. [PubMed: 235244]
23. Ivleva NB, Bramlett MR, Lindahl PA, Golden SS. *EMBO J*. Mar 23.2005 24:1202. [PubMed: 15775978]
24. Ivleva NB, Gao T, LiWang AC, Golden SS. *Proc Natl Acad Sci U S A*. Nov 14.2006 103:17468. [PubMed: 17088557]
25. Iwasaki H, Nishiwaki T, Kitayama Y, Nakajima M, Kondo T. *Proc Natl Acad Sci U S A*. Nov 26.2002 99:15788. [PubMed: 12391300]
26. Pattanayek R. *Mol Cell*. Aug 13.2004 15:375. [PubMed: 15304218]
27. Wood TL. *Proc Natl Acad Sci U S A*. Mar 30.2010 107:5804. [PubMed: 20231482]
28. Materials and methods are available as supporting material on *Science* Online.
29. We thank G. Dong for advice on culture conditions and K. Cook, V. Denic, Q. Justman, J. Markson, A. Rizvi and V. Vijayan for valuable discussions and comments on the manuscript. E. K. O. is an investigator of the Howard Hughes Medical Institute. NIH grant GM62419 supported work in the lab of S.S.G. M. J. R. was supported by a Helen Hay Whitney Foundation postdoctoral fellowship and by a Burroughs-Wellcome Foundation Career Award at the Scientific Interface.

Fig. 1.

Sustained drop in the ATP/ADP ratio and a phase shift in the circadian clock in response to a pulse of darkness. **(A)** Bioluminescence rhythms of a reporter under circadian control (*PkaIBC^ΔluxAB*) in synchronized cultures either grown in constant light (*black symbols*) or subjected to an 8-hour pulse of darkness (*shaded region*) at *t* = 29 hours (*blue symbols*). Bars show the range of duplicate measurements from the same experiment. Time is measured from the release of cultures into constant light. **(B)** Same as (A) except that the dark pulse was applied at *t* = 40 hours. **(C)** Relative levels of ATP and ADP extracted from cultures before, during (*shaded region*) and after a dark pulse in the conditions shown in (A) (*blue symbols*) and in (B) (*red symbols*). Bars indicate the range from duplicate cultures.

Fig. 2.

Phase shifts in the *in vitro* KaiABC oscillator caused by changing the ATP/ADP ratio. **(A)** Phase shift induced by adding an amount of ADP equal to the amount of ATP in the reaction buffer ~6 hours before peak phosphorylation (*blue symbols*) relative to a control (*black symbols*). The shaded region indicates the time after addition of ADP, but before the addition of pyruvate kinase to convert ADP to ATP. **(B)** Same as (A) except that ADP was added ~6 hours after peak phosphorylation (*red symbols*). **(C)** Phase shifts induced by a pulse of ADP in the *in vitro* oscillator reaction (*lower panel*) shown adjacent to the rhythm of KaiC phosphorylation in the control (*upper panel*) to indicate where in the cycle the pulses were applied. Differences in phase were measured by fitting the data to a sinusoidal function. **(D)** Effect of addition of mixtures of nucleotides to lower the ATP/ADP ratio (*colored symbols*) to various levels relative to a control which received only ATP (*black symbols*). Total adenine nucleotide concentration was the same across all reactions. The shaded region indicates the time before addition of pyruvate kinase. **(E)** Quantification of phase shifts from the experiment shown in (D). The phase shift when only ATP is added is zero by definition. Differences in phase were measured by fitting the data to a sinusoidal function.

Fig. 3.

Inhibition of KaiC phosphorylation by ADP. **(A)** KaiC phosphorylation in a KaiA-KaiC reaction with different ratios of ADP to ATP + ADP. Solid lines are fits to the first 4 hours of data used to isolate the first phosphorylation step (see Supplemental Text). **(B)** KaiC dephosphorylation in a KaiB-KaiC reaction with different ratios of ADP to ATP + ADP. Solid lines are fit to a first-order reversible reaction. **(C)** Kinase (k_{phos}) and phosphatase (k_{dephos}) rate constants are obtained from the data in (A). The dashed line is a fit to the

competitive inhibition model $k \frac{[ATP]}{[ATP] + K_{rel} [ADP]}$, where K_{rel} is an effective relative affinity for ADP vs. ATP, fit to 0.76 ± 0.15 . Error bars represent the standard error in the fit. **(D)** Phosphatase rate constants obtained from the data in (B). Error bars represent the standard error in the fit.

Fig. 4.

A mathematical model of the KaiABC oscillator predicts entrainment by varying ATP/ADP. **(A)** A schematic of a mathematical model for nucleotide-driven entrainment of the KaiABC oscillator. Oscillations occur because the S431-phosphorylated form of KaiC promotes its own production from KaiC phosphorylated at both S431 and T432 through a double-negative feedback loop involving sequestration of KaiA. Free KaiA promotes phosphorylation of KaiC and inhibits dephosphorylation. Changing illumination conditions *in vivo* drive changes in the adenosine pool, which modulate phosphorylation rates (k_{phos}). Though the figure shows only the effect on S431-P KaiC, changing KaiA activity and the ATP/ADP ratio affect all phosphorylation steps in the model. K_{rel} is an effective relative affinity for ADP versus ATP. $[\text{KaiA}]_{\text{free}}$ is the free concentration of [KaiA] and $K_{1/2}$ is the concentration of [KaiA] needed to produce a half-maximal effect on KaiC. k_{phos}^A is the maximum phosphorylation rate with saturating KaiA and 100% ATP. See the Supporting Material and (5) for complete details. **(B)** Simulation of the effect of lowering the ratio of ATP to ATP + ADP from 100% to 50% for 5 hours during the subjective day (*shaded region*) using equal affinities for ATP and ADP. **(C)** Phase response curve predicted by the model using equal affinities for ATP and ADP (*blue dashed line*) as in (B) compared with *in vitro* data (*black circles*) and *in vivo* data (*green squares*) from (19) (4-hour dark pulse, solid media) and from this study (*hollow triangles*) (8-hour dark pulse, liquid media). All data sets were aligned so that peak phosphorylation occurs at 36 hours (*grid line*), and the modeling results were scaled on both axes to make the period equal 24 hours.