Published in final edited form as:

Tissue Antigens. 2012 April ; 79(4): 249–262. doi:10.1111/j.1399-0039.2012.01849.x.

Analysis of MHC class I folding: novel insights into intermediate forms

Laura C. Simone^{1,*}, Amit Tuli^{1,2,**}, Peter D. Simone¹, Xiaojian Wang¹, and Joyce C. Solheim^{1,2,3}

¹Eppley Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, Omaha, NE 68198

²Department of Biochemistry and Molecular Biology, University of Nebraska Medical Center, Omaha, NE 68198

³Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE 68198

Abstract

Folding around a peptide ligand is integral to the antigen presentation function of major histocompatibility complex (MHC) class I molecules. Several lines of evidence indicate that the broadly cross-reactive 34-1-2 antibody is sensitive to folding of the MHC class I peptide-binding groove. Here, we show that peptide-loading complex proteins associated with the murine MHC class I molecule K^d are found primarily in association with the 34-1-2⁺ form. This led us to hypothesize that the 34-1-2 antibody may recognize intermediately, as well as fully, folded MHC class I molecules. In order to further characterize the form(s) of MHC class I molecules recognized by 34-1-2, we took advantage of its cross-reactivity with L^d. Recognition of the open and folded forms of L^d by the 64-3-7 and 30-5-7 antibodies, respectively, has been extensively characterized, providing us with parameters against which to compare 34-1-2 reactivity. We found that the 34-1-2⁺ L^d molecules displayed characteristics indicative of incomplete folding, including increased tapasin association, endoplasmic reticulum retention, and instability at the cell surface. Moreover, we demonstrate that an L^d-specific peptide induced folding of the 34-1-2⁺ L^d intermediate. Altogether, these results yield novel insights into the nature of MHC class I molecules recognized by the 34-1-2 antibody.

Keywords

antigen processing and presentation; conformation; epitope; folding intermediate; MHC class I; monoclonal antibody; peptide; tapasin

Correspondence: Joyce C. Solheim, Eppley Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, 986805 Nebraska Medical Center, Omaha, NE 68198-6805, United States. Phone: +1 402 559 4539. Fax: +1 402 559 4651. jsolheim@unmc.edu.

*Current address: Department of Biochemistry and Molecular Biology, University of Nebraska Medical Center, Omaha, NE 68198, United States.

**Current address: Division of Rheumatology, Immunology, and Allergy, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, United States.

Conflicts of interest notification: The authors confirm that there are no conflicts of interests.

INTRODUCTION

Cytotoxic T-cell responses against infected and aberrant cells are elicited through recognition of major histocompatibility complex (MHC) class I-peptide complexes at the cell surface. The fully assembled MHC class I molecule is a heterotrimer consisting of a polymorphic MHC class I heavy chain, the conserved light chain β_2m , and an 8–10 amino acid peptide. The MHC class I heavy chain structure is comprised of three extracellular domains. The $\alpha 1$ and $\alpha 2$ domains form the peptide-binding groove, and the $\alpha 3$ domain associates with β_2m and connects with the transmembrane region. Folding of MHC class I molecules occurs in a sequential manner. Upon translocation into the endoplasmic reticulum (ER) lumen, the MHC class I $\alpha 3$ domain quickly acquires its native disulfide bond and associates with β_2m (1,2). At this point, the resulting MHC class I heavy chain/ β_2m heterodimer is devoid of peptide and the $\alpha 1$ and $\alpha 2$ domains remain incompletely folded. Peptide loading is facilitated by the recruitment of empty MHC class I heavy chain/ β_2m heterodimers into a complex with the transporter associated with antigen processing (TAP), tapasin, ERp57, and calreticulin (3). Binding of an appropriate peptide induces folding of the $\alpha 1$ and $\alpha 2$ domains and release of the MHC class I heavy chain/ β_2m /peptide heterotrimer from the loading complex.

Given that the folding and conformation of MHC class I molecules are integral to antigen presentation, the identification and characterization of MHC class I conformational states are important issues. In terms of assessment of folding, one of the most extensively studied MHC class I molecules is the murine L^d molecule. The $\alpha 1$ domain-specific monoclonal antibody (mAb) 64-3-7 recognizes incompletely folded L^d molecules (4,5). Completely denatured L^d heavy chains, as well as L^d/β_2m heterodimers, are identified by the 64-3-7 mAb (6). However, the addition of appropriate peptide ligands induces a dose-dependent loss of 64-3-7⁺ L^d molecules (6–8), indicating that conformational changes occurring upon peptide binding mask the 64-3-7 epitope. Moreover, peptides have not been detected in association with 64-3-7⁺ L^d , suggesting that the peptide-binding groove of this conformer is indeed empty (4,7). Yet, as previously noted, it remains possible that the 64-3-7 mAb may recognize L^d molecules with poorly bound peptides which are unable to induce complete folding of the $\alpha 1/\alpha 2$ domains (8).

In contrast to the 64-3-7 mAb, the 30-5-7 mAb recognizes fully conformed L^d/β_2m /peptide heterotrimers (4,9). Furthermore, 64-3-7⁺ and 30-5-7⁺ L^d molecules exhibit a precursor-product relationship (6,7,9), such that peptide binding induces the folding of 64-3-7⁺ L^d molecules into 30-5-7⁺ L^d molecules. Importantly, the ability to serologically detect open, peptide-free MHC class I conformers has been extended to other mouse and human MHC class I molecules (including K^b , K^d , and HLA-B27), through introduction of the 64-3-7 epitope into the heavy chain by site-directed mutagenesis (8,10,11).

Within the peptide-loading complex, tapasin enables the selective loading of high-affinity peptides onto MHC class I molecules (12–15). Biophysical and *in vitro* studies examining tapasin-mediated peptide editing suggest that MHC class I molecules transition through an intermediate form during peptide loading (16–18). However, the molecular nature of such a transition state has yet to be fully defined. In the following studies, we took a serologic approach to defining MHC class I folding status. Although serology cannot discern fine molecular detail, mAbs are valuable tools in understanding MHC class I folding. For instance, the characterization of L^d folding with the 64-3-7 and 30-5-7 mAbs revealed important insights into the conformational changes that occur within the MHC class I peptide-binding groove upon β_2m /peptide association (4–7,9). In addition to the 64-3-7 and 30-5-7 mAbs described above, we utilized the 34-1-2 mAb. The 34-1-2 mAb recognizes the $\alpha 1$ domain of K^d molecules and has wide cross-reactivity with many MHC class Ia and Ib

molecules in the d, b, s, r, q, and p haplotypes, including weak cross-reactivity with L^d (19–23). Both mutation of L^d at positions within the peptide-binding groove and incubation of L^d with specific peptides greatly increases the level of 34-1-2⁺ L^d, indicating that the 34-1-2 mAb detects folded, peptide-bound MHC class I molecules (23). Furthermore, the 34-1-2 mAb is sensitive to nuances in the interaction between the MHC class I heavy chain and β_2m (22,23).

We have previously found that tapasin is readily detected in association with the 34-1-2⁺ form of K^d molecules (24,25). Therefore, we asked if, in addition to recognizing folded, peptide-bound MHC class I molecules, the 34-1-2 mAb also identifies a disparate population of intermediately folded MHC class I. Specifically, we asked whether the 34-1-2⁺ MHC class I molecules display characteristics consistent with sub-optimal peptide loading, including ER retention and cell surface instability. We also tested whether a high-affinity peptide could induce the folding of the 34-1-2⁺ intermediate. Our findings provide valuable information necessary for the correct interpretation of immunoprecipitation and flow cytometric experiments utilizing the 34-1-2 mAb. Furthermore, by distinguishing a functionally relevant tapasin substrate, these studies open new doors to the investigation of the molecular mechanisms underlying peptide optimization.

MATERIAL AND METHODS

Antibodies

The $\alpha 1$ domain-specific mAb 64-3-7 was raised against L^d (26), but also strongly recognizes L^d and has been mainly used in the analysis of L^d. The 64-3-7 mAb recognizes open, peptide-free L^d molecules that are associated with tapasin and other members of the peptideloading complex (4–7,27). The 64-3-7 mAb additionally detects the open forms of K^d and K^b molecules to which the 64-3-7 epitope has been transferred by site-directed mutagenesis (8,10). The 30-5-7 mAb recognizes folded, peptide-bound L^d molecules (4,9,28). The 34-1-2 mAb binds to K^d on the $\alpha 1$ domain (19,22). In addition to recognizing the MHC class Ia d haplotype molecules (K^d, D^d, and weakly L^d), 34-1-2 also cross-reacts with the $\alpha 1/\alpha 2$ regions of the Qa molecules encoded by the BALB/c Q5 and Q6 genes, and with MHC class Ia molecules expressed by b, s, r, q, and p haplotype mice (19–23,29). Weak cross-recognition of L^d by 34-1-2 is enhanced when L^d is bound to human β_2m or particular peptide ligands, and by mutation of L^d at residues within the peptide-binding region, indicating that the 34-1-2 epitope is sensitive to the conformation of the peptide-binding groove (22,23). All the anti-H2 antibodies (64-3-7, 30-5-7, and 34-1-2) used in this study were gracious gifts from Dr. T. Hansen (Washington University, St. Louis, MO). The following antibodies were used to detect proteins on western blots: mouse anti-human tapasin mAb (BD Biosciences, San Jose, CA), hamster anti-mouse tapasin mAb (gift from Dr. T. Hansen (Washington University, St. Louis, MO)), rabbit anti-human TAP1 serum (Genetex, Irvine, CA), and rabbit anti-human ERp57 (Abcam, Cambridge, MA). Additionally, the 64-3-7 mAb was used to detect denatured L^d and epitope-tagged K^d molecules on western blots.

Cell lines

HeLa cells and DAP-3 L-cells were grown at 37°C in 5% CO₂ in RPMI 1640 medium supplemented with 15% fetal bovine serum, 2 mM glutamine, 1 mM sodium pyruvate, 0.1 mM MEM non-essential amino acids solution, 4 mM HEPES, penicillin (100 U/ml), and streptomycin (100 μ g/ml). The fetal bovine serum was purchased from Atlanta Biologicals (Lawrenceville, GA). All other media reagents were purchased from Invitrogen (Carlsbad, CA). Where indicated, HeLa cells were cultured for 48 h in the presence of 100 U recombinant human interferon-gamma (IFN- γ)/ml (R&D Systems, Minneapolis, MN).

Similarly, in some experiments, DAP-3 L-cells were cultured for 48 h in the presence of 100 U recombinant mouse IFN- γ /ml (BD Biosciences, San Jose, CA). The HeLa cell line was a gift from Dr. Wendy Maury (University of Iowa, Iowa City, IA). Epitope-tagged K^d and L^d were cloned into the mammalian expression vector RSV.5-neo and stably transfected into HeLa and DAP-3 L-cells using the Amaxa Nucleofector system (Lonza, Basel, Switzerland). Stable transfectants were selected with 0.4 mg/ml Geneticin (Invitrogen, Carlsbad, CA). The tapasin knockout mouse fibroblast cell lines used for the data shown in Supplementary Figure 1A have been previously described (30).

Immunoprecipitations and western blots

In preparation for immunoprecipitations, cells were washed in phosphate-buffered saline (PBS) containing 20 mM iodoacetamide (Sigma-Aldrich, St. Louis, MO) three times and then lysed in 3-[(3-cholamidopropyl) dimethylammonio]-1-propanesulfonate (CHAPS) lysis buffer. The lysis buffer was composed of 1% CHAPS (Roche Applied Science, Indianapolis, IN) in Tris-buffered saline (TBS) (pH 7.4) with freshly added 0.2 mM phenylmethanesulfonyl fluoride (PMSF), 20 mM iodoacetamide, and excess antibody. After 1 h incubation on ice, the lysates were centrifuged to pellet the cell nuclei and the lysate supernatants were added to Protein A-Sepharose (PAS) (GE Healthcare, Piscataway, NJ) and mixed intermittently for 45 min. The beads were then washed four times in 0.1% CHAPS/20 mM iodoacetamide in TBS (pH 7.4) and boiled for 5 min in elution buffer (0.125 M Tris (pH 6.8)/2% sodium dodecyl sulfate (SDS)/12% glycerol/0.02% bromophenol blue) to elute the proteins from the beads. In some experiments, the immunoprecipitated proteins were reduced by adding glycoprotein denaturing buffer (0.5% SDS/40 mM dithiothreitol) (New England Biolabs, Ipswich, MA) and boiling the samples for 10 min. Where indicated, the immunoprecipitated proteins were treated with Endo H (New England Biolabs, Ipswich, MA) at 37°C for 2 h according to the manufacturer's instructions.

For some experiments, immunoprecipitation was used to deplete forms of MHC class I molecules recognized by specific antibodies. For these experiments, first cells were lysed in 1% CHAPS/0.2 mM PMSF/20 mM iodoacetamide in TBS (pH 7.4) with or without 500 μ M P29 peptide (YPNVNIHNF) and incubated at 4°C overnight. After the overnight incubation, the lysates were centrifuged, and the supernatant divided into 3 equal aliquots. The aliquots were sequentially incubated with either PAS alone (for the mock depletion), or PAS pre-absorbed to either the 64-3-7 or the 30-5-7 mAb. After the final round of immunodepletion, the lysates were transferred to PAS pre-absorbed to the 34-1-2 mAb. Following incubation with the cell lysate, the beads were washed and eluted as described above.

For western blotting, after elution the immunoprecipitates were electrophoresed on precast Tris-glycine SDS gels (Invitrogen, Carlsbad, CA) and the proteins were transferred to Immobilon-P membranes (Millipore, Billerica, MA). Following overnight blocking in 10% dry milk dissolved in 0.05% Tween 20 in PBS, the blotting membranes were incubated with primary antibody diluted in milk for 2 h and then washed three times with 0.05% Tween 20/PBS. Next, the membranes were incubated for 1 h with the appropriate secondary antibody diluted in 0.05% Tween 20/PBS. The secondary antibodies (goat anti-mouse IgG [light chain specific], mouse anti-rabbit IgG [light chain specific], and goat anti-Armenian hamster IgG [heavy and light chain specific]) were conjugated to horseradish peroxidase or biotin. All secondary antibodies were purchased from Jackson ImmunoResearch (West Grove, PA). Membranes incubated with biotinconjugated secondary antibody were washed three times with 0.05% Tween 20/PBS and then incubated with streptavidin-conjugated horseradish peroxidase (Invitrogen, Carlsbad, CA) diluted in 0.05% Tween 20/PBS for 1 h. Next, the membranes were washed three times with 0.3% Tween 20/PBS, and treated with enhanced chemiluminescence western blot developing reagents (Pierce Chemical Co., Rockford, IL). The blots were exposed to Kodak BioMax film (Eastman Kodak, Rochester, NY) to

visualize the bands. For some experiments, protein bands were quantified using the Molecular Imager ChemiDoc XRS system with Quantity One 1-D Analysis Software (Bio-Rad, Hercules, CA).

Flow cytometry assays

In flow cytometry assays, cells were suspended at $5 \times 10^6/\text{ml}$ in PBS with 0.2% bovine serum albumin and 0.1% sodium azide. Cell suspension aliquots in volumes of 0.1 ml were distributed to the wells of a 96-well plate. The cells were incubated with excess mAb or with bovine serum albumin/azide/PBS alone (as a control) at 4°C for 30 min, washed three times, and incubated with a phycoerythrin-conjugated, Fc-specific F(ab')_2 portion of goat anti-mouse IgG (Jackson ImmunoResearch, West Grove, PA) at 4°C for 30 min. The cells were washed three times, resuspended in bovine serum albumin/azide/PBS, and analyzed on a FACSCalibur flow cytometer (BD Biosciences, San Jose, CA). Statistical analyses were done with the Cell Quest software (BD Biosciences, San Jose, CA).

Where indicated, cells were incubated with 2 $\mu\text{g}/\text{ml}$ of brefeldin A (Bfa) (Sigma-Aldrich, St. Louis, MO) for various times prior to collection and analysis of the cells. To calculate the cell surface half-life of MHC class I, the following model was employed:

where F is folded, peptide-bound MHC class I, O is open, peptide-free MHC class I, s signifies that the protein is on the cell surface, and i signifies that the protein has been internalized. Thus, both folded, peptide-bound and open, peptide-free MHC class I can be internalized into the cell, and the folded MHC class I can melt into open MHC class I on the cell surface. Assuming that each process is a first-order elementary reaction, it can be shown that folded and open MHC class I obey the following decay equations:

$$\begin{aligned}
 [\text{F}_s] &= [\text{F}_s]_0 e^{-(k_1+k_2)t} \\
 [\text{O}_s] &= [\text{O}_s]_0 e^{-k_3t} + \frac{k_2[\text{F}_s]_0}{k_3-k_1-k_2} (e^{-(k_1+k_2)t} - e^{-k_3t}).
 \end{aligned}$$

Since the initial concentration of protein on the surface was set to 100%, the data for folded MHC class I were fit to the equation $[\text{F}_s] = 100e^{-at}$, using Mathematica (Wolfram Research). Fitting the data for 34-1-2⁺ L^d to the open MHC class I equation $[\text{O}_s] = 100e^{-bt} + c(e^{-at} - e^{-bt})$ gave the best fit. The half-life was calculated for folded MHC class I (34-1-2⁺ K^d and 30-5-7⁺ L^d) using the standard equation for exponential decay ($t_{1/2} = \ln 2 / a$). For 34-1-2⁺ L^d, according to the model for open MHC class I, the half-life is not constant and was calculated by setting $[\text{O}_s]$ equal to 50% in the best fit equation and solving for $t_{1/2}$ numerically.

Structure analysis

The crystal structure of L^d (Protein Data Bank ID 1LD9) (31) and K^d (Protein Data Bank ID 2FWO) (32) were superimposed using LSQKAB (33) in CCP4 (34). Hydrogen bonds were

verified by measuring interatomic distances with Coot (35). Structural figures were created using PyMOL (36).

RESULTS

Comparison of L^d and K^d association with peptide-loading complex proteins using MHC class I heavy chain conformation-specific mAbs

Previous studies from our lab have reported that tapasin co-immunoprecipitates strongly with the 34-1-2⁺ form of K^d (24,25). In this study, we expanded upon this initial observation by comparing the association of tapasin with serologically detectable forms of L^d or K^d. Immunoprecipitation of L^d molecules from HeLa-L^d or L-L^d cell lysates with the 64-3-7 mAb or the 30-5-7 mAb (Figures 1A and B, top panel) revealed association of tapasin with the open (64-3-7⁺), but not the folded (30-5-7⁺), L^d molecules (Figures 1A and B, bottom panel). In both HeLa-K^b and L-K^b cells, tapasin was also found to associate with the open (64-3-7⁺) form of K^b, but not folded (Y3⁺) K^b molecules (data not shown). Moreover, HLA molecules endogenously expressed in HeLa cells exhibited binding to tapasin exclusively in their peptide-free (HC10⁺) form (data not shown). These results are in accord with the view that peptide-free MHC class I molecules are integrated into the peptide-loading complex through association with tapasin (24,37–41). In contrast, tapasin was more readily detected in association with the 34-1-2⁺ form of K^d rather than with open (64-3-7⁺) K^d molecules in both HeLa-K^d and L-K^d cells (Figures 1C and D). In agreement with a previous study (8), we have also observed tapasin in association with 64-3-7⁺ K^d molecules in some experiments (Supplementary Figure 1A). However, as shown in Figures 1C and D, the level of tapasin bound to the 34-1-2⁺ form of K^d is substantially greater than the level bound to 64-3-7⁺ K^d molecules. Thus, in both human and mouse systems, tapasin preferentially binds to K^d molecules that possess the 34-1-2 epitope over 64-3-7⁺ K^d molecules.

We hypothesized that the association of other peptide-loading complex proteins with L^d or K^d would mirror tapasin binding. Indeed, we found that both TAP and ERp57 associate with the 64-3-7⁺ form of L^d (Figures 2A and B) whereas TAP and ERp57 preferentially bind to 34-1-2⁺ (as compared to 64-3-7⁺) K^d molecules (Figures 2C and D). These data indicate that other assembly complex proteins, in addition to tapasin, share association with 34-1-2⁺ K^d.

Serologic analysis of MHC class I surface expression indicates that 34-1-2⁺ L^d molecules are more unstable than are 30-5-7⁺ L^d molecules

If the 34-1-2 mAb exclusively recognizes completely folded MHC class I molecules, then the interpretation of the data shown in Figures 1 and 2 would be that most K^d molecules within the peptide-loading complex have folded around a peptide ligand. This conclusion opposes the model whereby conformational changes occurring within the $\alpha 1/\alpha 2$ domains upon peptide binding induce dissociation of the MHC class I molecule from tapasin and other peptide-loading complex proteins. An alternative explanation is that some K^d molecules have bound to peptides that have induced partial folding of the $\alpha 1/\alpha 2$ domains (as detected by the 34-1-2 mAb); however, this partial folding is insufficient to cause release from tapasin. This would suggest that, in addition to recognizing fully folded MHC class I heterotrimers, the 34-1-2 mAb also detects an intermediate in the folding process. Therefore, we sought to further characterize the 34-1-2⁺ form of MHC class I molecules.

To do so, we took advantage of the cross-reactivity of the 34-1-2 mAb with L^d, since we could also distinguish incompletely and fully folded L^d molecules with the 64-3-7 and 30-5-7 mAbs, respectively. As the level of 34-1-2⁺ L^d molecules is very low, we chose to analyze HeLa cells since human $\beta 2m$ enhances the detection of 34-1-2⁺ L^d (22,23). Additionally, the HeLa-L^d cells were treated with IFN- γ , which was found to boost the level

of 34-1-2⁺ L^d (Supplementary Figure 1B), likely through transcriptional induction of peptide-loading complex proteins. The 34-1-2 epitope was also found on L^d molecules expressed in mouse L-cell fibroblasts (Supplementary Figure 1C), indicating that 34-1-2⁺ L^d molecules are not simply artifacts induced by L^d association with human β_2m . In further support of the theory that the 34-1-2 epitope naturally occurs on L^d molecules, previous work has demonstrated that various peptide ligands selectively augment the level of 34-1-2⁺ L^d molecules at the surface of mouse L-cell fibroblasts (23).

First, to gain a better understanding of the nature of 34-1-2⁺ MHC class I molecules, we evaluated the level of 64-3-7⁺, 30-5-7⁺, and 34-1-2⁺ L^d expressed on HeLa-L^d cells incubated at either 37°C or 25°C (Figure 3A). Incubation of cells at 25°C permits the surface expression of unstable MHC class I molecules (42). For comparison, we also assessed the level of 64-3-7⁺ and 34-1-2⁺ K^d expressed on HeLa-K^d cells (Figure 3A). Relative to incubation at 37°C, the 25°C incubation decreased the surface expression of 64-3-7⁺ L^d molecules. Surface expressed 64-3-7⁺ L^d molecules arise from unfolding of 30-5-7⁺ L^d molecules at the plasma membrane (7). Thus, incubation at a lower temperature appears to promote the stabilization of cell surface L^d molecules, preventing them from melting to the open form. In contrast, the level of 64-3-7⁺ K^d was not influenced by the reduced temperature incubation. A likely explanation is that the folded K^d molecules at the cell surface are highly stable, and do not readily melt at 37°C. The levels of 30-5-7⁺ L^d and 34-1-2⁺ L^d were both increased by the 25°C incubation, suggesting that colder temperatures stabilize intracellular peptide-MHC class I complexes and promote their trafficking to the cell surface. Importantly, while the level of 30-5-7⁺ L^d was increased 2-fold at 25°C, the level of 34-1-2⁺ L^d was increased 8-fold. The greater extent of surface induction of the 34-1-2⁺ form of L^d as compared to 30-5-7⁺ L^d suggests that the 34-1-2⁺ form is intrinsically more unstable and thus more amenable to stabilization at reduced temperatures.

We also compared the surface stability of the serologic forms of L^d and K^d by assessing the level of each form that remained at the plasma membrane on cells treated with Bfa for varying lengths of time. Because the pool of open, 64-3-7⁺ MHC class I molecules at the cell surface can be sustained by the melting of folded K^d and L^d molecules, we limited our analysis to the turnover of 30-5-7⁺ and/or 34-1-2⁺ L^d and K^d molecules. Using a different method to detect MHC class I internalization, Mahmutefendić, et al found that the 64-3-7⁺ forms of L^d are endocytosed at a faster rate than are the folded, 30-5-7⁺ L^d molecules (43). In our studies, we found that the cell surface 34-1-2⁺ K^d molecules were highly stable, having a half-life of over 16 h (Figure 3B). As compared to K^d, the folded, 30-5-7⁺ L^d decayed more quickly from the cell surface, with a half-life of 7.2 h (Figure 3B). In striking contrast to the 34-1-2⁺ form of K^d, the 34-1-2⁺ L^d molecules were exceptionally unstable. Virtually all of the 34-1-2⁺ L^d molecules were lost from the cells within 3 hours (Figure 3B). The extreme cell surface instability of 34-1-2⁺ L^d molecules is consistent with the idea that this pool of molecules is imperfectly folded.

These data also reveal important differences between the K^d and L^d molecules. First, in contrast to the substantial induction of 30-5-7⁺ L^d surface expression, incubation at 25°C only moderately increased the level of cell surface 34-1-2⁺ K^d molecules (Figure 3A). This is in agreement with a prior observation that culturing cells at 23-26°C induces cell surface L^d molecules to a greater extent as compared to other MHC class I molecules (9). Thus, at steady state, cell surface K^d molecules appear to be maximally stable and cannot be further stabilized by reduced temperature. In addition, 34-1-2⁺ K^d molecules persisted at the cell surface for a substantially longer time than did either 30-5-7⁺ or 34-1-2⁺ L^d molecules (Figure 3B). Together, these findings likely reflect the peptide-binding characteristics of the L^d and K^d molecules. Indeed, whereas the peptide-binding groove of L^d is shallow and hydrophobic (31,44,45), the K^d peptide-binding groove possesses deep pockets that allow

for tight peptide binding (32,46–49). Therefore, the large induction of 34-1-2⁺ L^d surface expression at 25°C as compared to the minor increase in cell surface 34-1-2⁺ K^d at 25°C (Figure 3A), as well as the relative cell surface stabilities of L^d and K^d (Figure 3B), may be explained by the peptide-binding properties of these two MHC class I molecules.

The majority of 34-1-2⁺ L^d molecules display immature glycosylation, and tapasin preferentially binds to the 34-1-2⁺ over the 30-5-7⁺ form of L^d

Next, we characterized the 34-1-2⁺ L^d molecules by immunoprecipitation and western blotting. L^d molecules were isolated from HeLa-L^d cells using the 64-3-7, 34-1-2, or 30-5-7 mAb. Because the level of 34-1-2⁺ L^d is very low as compared to the amount of 30-5-7⁺ and 64-3-7⁺ L^d, the number of cells that were used for the immunoprecipitation with 34-1-2 was increased four fold. The immunoprecipitated proteins were then mock treated or treated with the enzyme Endo H. Maturation of glycoproteins in the medial Golgi yields complex oligosaccharides which are resistant to Endo H cleavage (50). Thus, Endo H digestion can be used to differentiate immature glycosylation (Endo H-sensitive) from mature glycosylation (Endo H-resistant) on MHC class I molecules (51,52). As is apparent in Figure 4A, the 64-3-7⁺ L^d molecules are predominantly Endo H-sensitive, whereas the 30-5-7⁺ L^d molecules are approximately equal parts Endo H-sensitive and Endo H-resistant, consistent with a previous report (4).

In order to investigate the relative level of tapasin association with the three serologic forms of L^d, equivalent amounts of L^d molecules immunoprecipitated with the 64-3-7, 34-1-2, or 30-5-7 mAb were probed on western blots for the L^d heavy chain (Figure 4B, upper panel) or for tapasin (Figure 4B, lower panel). Consistent with the premise that open, peptide-free MHC class I molecules are incorporated into the peptide-loading complex, the 64-3-7⁺ L^d molecules showed the highest level of tapasin association, and very little tapasin was found in association with 30-5-7⁺ L^d molecules. Markedly, the 34-1-2⁺ L^d molecules were bound to tapasin at a level that was only 30% less than that of the 64-3-7⁺ L^d molecules. This suggests that the 34-1-2⁺ form of L^d is not readily released from the peptide-loading complex. Tapasin was also detected in association with 34-1-2⁺ L^d molecules expressed in mouse L-cell fibroblasts (Supplementary Figure 1C), indicating that the binding of tapasin to 34-1-2⁺ L^d molecules does not require the presence of xenogeneic β₂m.

Maturation of the L^d heavy chain carbohydrate moiety can also be inferred from the western blot shown in Figure 4B. The L^d heavy chain bands with a faster migration on the gel correspond to L^d molecules with immature glycosylation, whereas carbohydrate modification results in slower migration of the MHC class I molecule (compare migration of 30-5-7⁺ L^d, mock-treated or Endo H-treated, in Figure 4A with 30-5-7⁺ L^d in Figure 4B). Quantification of the L^d heavy chain bands in Figure 4B revealed that the majority (65%) of the 34-1-2⁺ L^d molecules display immature glycosylation (Figure 4C). Thus, most of the 34-1-2⁺ L^d molecules have not trafficked past the medial Golgi. In contrast, slightly over half of the 30-5-7⁺ L^d molecules possess mature glycosylation (Figure 4C). The vast majority (95%) of the 64-3-7⁺ L^d molecules display immature glycosylation (Figure 4C). Together, the findings that most of the 34-1-2⁺ L^d molecules are retained within the ER and that this pool of L^d molecules is found in association with tapasin to a greater extent than are folded, 30-5-7⁺ L^d molecules strongly suggest that the 34-1-2 epitope can be present on incompletely folded MHC class I molecules.

The 34-1-2⁺ L^d intermediate conformer overlaps with the pools of 64-3-7⁺ L^d and 30-5-7⁺ L^d molecules, and peptide induces folding of the 64-3-7⁺34-1-2⁺30-5-7⁺ L^d intermediate

Finally in our analysis of the 34-1-2⁺ L^d molecules, we asked whether the 34-1-2⁺ L^d form is separate from the 64-3-7⁺ L^d and 30-5-7⁺ L^d molecules. To address this, we pre-cleared

HeLa-L^d cell lysates with fourteen rounds of incubation with either the 64-3-7 mAb or the 30-5-7 mAb (Figures 5A and 5B, lanes 1–6). As a control, HeLa-L^d cell lysates were mock pre-cleared. Following removal (or mock removal) of the 64-3-7⁺ or 30-5-7⁺ L^d molecules, the HeLa-L^d cell lysates were subjected to immunoprecipitation using the 34-1-2 mAb (Figure 5C, lanes 1-3). Removal of the 64-3-7⁺ L^d molecules resulted in a reduction, but not a complete loss, of the 34-1-2⁺ L^d molecules (Figure 5C, compare lanes 1 and 2), indicating that a portion of the 34-1-2⁺ L^d molecules are not also 64-3-7⁺. On the other hand, depletion of the 30-5-7⁺ L^d molecules eliminated all of the detectable 34-1-2⁺ L^d molecules (Figure 5C, compare lanes 1 and 3). Thus, all of the 34-1-2⁺ L^d molecules also possess the 30-5-7 epitope. Together, the findings that the 64-3-7 and 30-5-7 immunodepletions both impacted the level of 34-1-2⁺ L^d, and that the 30-5-7⁺ and 34-1-2⁺ pools of L^d completely overlap, imply that the 30-5-7⁺ and 64-3-7⁺ pools of L^d must partially overlap (Figure 5D). Furthermore, this overlapping pool is that which is recognized by the 34-1-2 mAb, and is therefore 64-3-7⁺, 34-1-2⁺, and 30-5-7⁺ (Figure 5D). In support of the overlap between the 64-3-7⁺ and 30-5-7⁺ pools of L^d, we have observed tapasin in association with 30-5-7⁺ L^d in IFN- γ -treated cells on very long film exposures (Supplementary Figure 1D). Additionally, because the 64-3-7 depletion did not completely eliminate the 34-1-2⁺ L^d pool, there must also exist a separate pool of 34-1-2⁺30-5-7⁺ L^d molecules (Figure 5D).

We reasoned that if the 34-1-2⁺ L^d molecules represent an intermediately folded form, then addition of an L^d-specific peptide should induce folding of the intermediate. To test this, HeLa-L^d cell lysates were first incubated with the P29 peptide, which is a known endogenous L^d ligand (53). Following overnight incubation with peptide, the lysates were mock depleted, or immunodepleted with either the 64-3-7 mAb or the 30-5-7 mAb (Figures 5A and 5B, lanes 8–13). As has been previously shown (6,9), addition of the P29 peptide induced L^d folding, as seen in the drop in the level of open, 64-3-7⁺ L^d molecules (Figure 5A, compare lanes 1-5 with lanes 8-13), and a rise in the level of folded, 30-5-7⁺ L^d molecules (Figure 5B, compare lanes 1–5 with lanes 8–13). Finally, the lysates were immunoprecipitated with the 34-1-2 mAb (Figure 5C, lanes 5–7). Similar to the situation without peptide, removal of the 30-5-7⁺ L^d molecules from peptidetreated lysates resulted in the loss of all detectable 34-1-2⁺ L^d molecules (Figure 5C, compare lanes 5 and 7). However, in the presence of P29, removal of the 64-3-7⁺ L^d molecules did not impact the level of 34-1-2⁺ L^d (Figure 5C, compare lanes 5 and 6). This demonstrates that the P29 peptide induces folding of the 64-3-7⁺34-1-2⁺30-5-7⁺ L^d intermediate (Figure 5D).

DISCUSSION

Multiple lines of evidence presented throughout this report indicate that, besides recognizing MHC class I molecules that are optimally folded with peptide, the 34-1-2 mAb is additionally capable of identifying MHC class I molecules with an incompletely folded peptide-binding groove. Our data do not exclude 34-1-2 mAb recognition of fully folded, peptide-bound MHC class I molecules. Indeed, the high level of 34-1-2⁺ MHC class I molecules on the surface of cells which endogenously express K^d (19), as well as the ability of the 34-1-2 mAb to block the cytolytic and proliferative response of MHC class I-restricted T-cells (54), attest to the ability of the 34-1-2 mAb to detect folded MHC class I molecules. However, our current studies demonstrate that the 34-1-2 epitope is also present on a subset of 64-3-7⁺ L^d molecules, and that this pool of MHC class I molecules is peptide-receptive (Figure 5). In contrast to L^d, immunodepletion of 64-3-7⁺ K^d molecules did not impact the amount of 34-1-2⁺ K^d that was immunoprecipitated from HeLa-K^d cells (Supplementary Figure 2), indicating that the pools of 64-3-7⁺ and 34-1-2⁺ K^d molecules do not overlap to an appreciable extent. L^d molecules bind peptides relatively weakly and are prone to peptide dissociation (45). Thus, in contrast to K^d, L^d may be able to adopt an intermediately folded (34-1-2⁺) conformation that is more unstable than the K^d form, and/or

one on which the 64-3-7 epitope is also exposed. In order to further characterize 34-1-2⁺ K^d molecules, it will be useful to identify mAbs that exclusively recognize fully folded K^d. In this regard, our preliminary studies show that tapasin does not co-immunoprecipitate with K^d molecules recognized by the MA-215 or Do4 mAbs (Supplementary Fig 3). MA-215 binds to an undefined epitope on the $\alpha 1/\alpha 2$ domains (55). The domain specificity of Do4 remains unknown; however, the finding that Do4 blocks cytotoxic T-cell responses suggests that this mAb detects folded MHC class I (56). Ongoing studies in our lab are aimed to determine the extent to which MA-215 and Do4 may discriminate between suboptimally folded and fully folded MHC class I.

The 34-1-2 epitope has been broadly mapped to a region spanning residues 63-85 of the MHC class I $\alpha 1$ -helix (Figure 6A) (22,23,57). Within the $\alpha 1$ -helix, K^d and L^d differ by nine amino acids (Supplementary Fig 4). Thus, the 34-1-2 epitope is likely, in part, composed of one or more of these non-conserved residues. Indeed, mutation of Ld at position 70 of the $\alpha 1$ helix increases 34-1-2 mAb recognition (23,57), indicating that amino acid 70 influences the 34-1-2 epitope. The side chain of amino acid 70 is also part of the B pocket, which accommodates the second amino acid (P2) of antigenic peptides (58). It has been suggested that upon association with β_2m , newly synthesized MHC class I molecules quickly bind to peptides, which are likely to be an imperfect fit, but serve to stabilize the MHC class I molecule and prevent degradation (59,60). We propose that, by causing structural shifts within the $\alpha 1$ domain, these initial peptide cargoes induce the intermediate conformation that is detected by the 34-1-2 mAb. We will refer to this group of sub-optimally loaded MHC class I conformers as 34-1-2-INT (for intermediately folded, 34-1-2⁺ molecules) to distinguish them from fully folded, 34-1-2⁺ MHC class I molecules. Alternatively, the MHC class I peptide-binding groove may be able to adopt the 34-1-2-INT conformation in the absence of a peptide ligand. Indeed, molecular dynamics simulations of peptide-free MHC class I molecules indicate that regions of the groove which contact the peptide N-terminus (including residues 56-70 of the $\alpha 1$ helix) are able to adopt a “pre-formed” conformation that is similar to the structure of the peptide-bound state (61).

Regardless of peptide occupancy, the 34-1-2-INT molecules likely represent a n appropriate substrate for tapasin-mediated peptide optimization. We previously found that the presence of tapasin enhances the level of 34-1-2⁺ K^d molecules (24). In light of our current findings, this could result from tapasin-mediated folding of K^d and/or from tapasin stabilization of the 34-1-2-INT. As shown in Figure 6A, the 34-1-2 antibody binding site is located across the peptide-binding groove from the tapasin binding site, which has been proposed to include the short $\alpha 2$ -helix (residues 138–149) along with a conserved loop comprising residues 128–138 (27,39,60,62–64). Biophysical studies suggest that tapasin achieves peptide editing by destabilizing binding of the peptide C-terminus to promote dissociation of low-affinity peptides (16). During the peptide selection process, binding of the P2 anchor residue into the B pocket may shift the $\alpha 1$ -helix and expose the 34-1-2 epitope; at the same time, tapasin may remain bound to the $\alpha 2$ domain of the 34-1-2-INT molecule, until sufficiently stabilizing interactions with the peptide C-terminus induce complete folding. The P2 anchor residue for K^d-binding peptides is tyrosine, which fits into the deep B pocket of the K^d peptide-binding groove (32,46–49). In the crystal structure of a K^d-peptide complex, the hydroxyl group of P2 tyrosine forms a hydrogen bond with an aspartic acid residue at position 70 (Figure 6B) (32). Thus, tight binding of the P2 peptide anchor residue may permit K^d molecules to adopt the 34-1-2-INT conformation readily while the peptide C-terminus remains incompletely bound. On the other hand, in L^d molecules residue 70 is glutamine, which does not hydrogen bond with the P2 anchor residue proline (31). Thus, L^d molecules may acquire the 34-1-2 epitope through an alternative mode. As such, in the crystal structures of both L^d and K^d, residue 70 forms a hydrogen bond with peptide position 5 (P5) (Figure 6B). Thus, $\alpha 1$ domain interactions with P5 may permit L^d molecules to adopt

the 34-1-2-INT conformation. Additionally, residue 73 (which is tryptophan in both K^d and L^d molecules) has also been implicated in promoting 34-1-2 reactivity (65). Therefore, shifts in other α 1 domain residues, besides amino acid 70, during peptide binding may reveal the epitope of the 34-1-2 mAb. Overall, this model suggests that the 34-1-2-INT MHC class I molecules are undergoing tapasin-mediated peptide editing.

In order to attain a fully folded conformation, 34-1-2-INT MHC class I molecules may require peptide exchange. Alternatively, the sub-optimal peptide itself may be able to induce the conversion of 34-1-2-INT MHC class I molecules to the fully folded conformer. Support for the latter hypothesis comes from a study by Praveen et al that assessed the kinetics of peptide binding (17). This study demonstrated that upon the initial interaction between the MHC class I molecule and peptide, an intermediate complex is formed which slowly converts to a stable, folded state. These authors also concluded that, in the absence of peptide competition, lower-affinity peptides may be able to induce the conversion of the intermediate complex into fully folded MHC class I molecules. Furthermore, additional layers of quality control may facilitate the optimization of the cargo bound to 34-1-2-INT MHC class I molecules. Along these lines, Cresswell and colleagues have recently demonstrated that UDP-glucose:glycoprotein glucosyltransferase 1 (UGT1) recognizes and re-glycosylates sub-optimally folded MHC class I molecules within the ER, thereby allowing these MHC class I molecules to undergo another round of peptide-editing (66). Our present studies indicate that 34-1-2-INT MHC class I molecules may be appropriate substrates for UGT1. Thus, our findings could provide an appropriate model for studying how UGT1 detects sub-optimal folding. Altogether, 34-1-2-INT molecules may be able to transition into stably folded, peptide-bound molecules, either by a mechanism involving peptide exchange or by peptide-induced conformational changes.

On the other hand, 34-1-2-INT MHC class I molecules might also arise from the partial melting of peptide-folded MHC class I molecules. If so, then the immunodepletion of 30-5-7⁺ L^d molecules may partially impact the level of 34-1-2⁺ L^d (as observed in Figure 5C) through the removal of the 34-1-2-INT precursor rather than through the removal of a theoretical 34-1-2⁺/30-5-7⁺ double-positive population. On the other hand, a substantial proportion (40%) of the 34-1-2⁺ L^d molecules have acquired Endo H resistance (Figures 4B and 4C). Two non-mutually exclusive possibilities could account for this pool of Endo-H resistant, 34-1-2⁺ L^d molecules. First, this pool of molecules may represent MHC class I molecules which left the ER in a fully folded conformation, but then melted to the 34-1-2-INT conformer. Secondly, the Endo H-resistant, 34-1-2⁺ L^d molecules may represent molecules that were permitted to leave the ER with the 34-1-2 epitope. At this point, it is unclear to what extent 34-1-2⁺, Endo H-resistant L^d molecules egress to the cell surface or are recycled back to the ER through ER/Golgi quality control mechanisms (67,68). Furthermore, it has recently been demonstrated that proprotein convertase 7 facilitates the stabilization of sub-optimally loaded MHC class I molecules in a post-ER compartment (69). Thus, through the action of proprotein convertase 7 or an analogous mechanism, the 34-1-2⁺, Endo H-resistant L^d molecules may acquire peptides capable of inducing the fully folded conformation before reaching the plasma membrane. Indeed, 34-1-2⁺ L^d molecules are stably expressed at the cell surface when they are appropriately loaded with peptide (23).

In future studies, the characterization of the nature and affinity of peptide ligands bound by 34-1-2-INT MHC class I molecules should shed more light on the transitional states of MHC class I. Indeed, the ability of an MHC class I molecule to adopt the 34-1-2-INT conformation may depend on particular peptide sequences (23). Many of the peptide sequences that bind to 34-1-2-INT, but do not generate the folded conformer, are likely to be displaced during tapasin-mediated peptide selection. Thus, the identification of these peptides should permit the classification of endogenous antigens that tapasin specifically

selects against. The ability of the 34-1-2 mAb to recognize many MHC class I allotypes (19–21) will make it broadly useful in such analyses. Moreover, precise knowledge of the structural states of the MHC class I molecules that associate with tapasin has been long sought, and will provide insight regarding the tapasin-MHC class I interface. In this regard, the purification of 34-1-2-INT molecules may make possible the structural determination of the MHC class I intermediate conformation. Overall, the ability to identify and capture MHC class I molecules in the process of peptide editing will aid in advancing our understanding of the molecular mechanisms underlying antigenic peptide selection.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Abbreviations used in this paper

β₂m	beta 2-microglobulin
Bfa	brefeldin A
CHAPS	3-[(3-cholamidopropyl) dimethylammonio]-1-propanesulfonate
Endo H	endoglycosidase H
ER	endoplasmic reticulum
IFN-γ	interferon-gamma
mAb	monoclonal antibody
MHC	major histocompatibility complex
PAS	Protein A-Sepharose
PBS	phosphate-buffered saline
PMSF	phenylmethanesulfonyl fluoride
SDS	sodium dodecyl sulfate
TAP	transporter associated with antigen processing
TBS	Tris-buffered saline
UGT1	UDP-glucose:glycoprotein glucosyltransferase 1

Acknowledgments

This work was supported by NIH Grants R01 GM057428 and R01 GM057428-09S1 (J.C.S.), a Nebraska Department of Health and Human Services LB506 grant (J.C.S.), and by the Nebraska Center for Cellular Signaling (funded by National Institute of Research Resources Centers of Biomedical Research Excellence Grant P20 RR018759) (J.C.S.). Support for this work was also provided by UNMC Graduate Studies Fellowships (L.C.S. and A.T.), an NIH/NCI Training Grant T32 CA009476 Fellowship (L.C.S.), and a Department of Education Graduate Assistance in Areas of National Need Fellowship (P.D.S.). We gratefully acknowledge the personnel of the UNMC Cell Analysis Core Facility (funded in part from National Cancer Institute Cancer Center Support Grant P30 CA036727). We thank Dr. Ted Hansen and Dr. Wendy Maury for generously providing cell lines, antibodies, and cDNA.

REFERENCES

1. Ribaldo RK, Margulies DH. Independent and synergistic effects of disulfide bond formation, beta 2-microglobulin, and peptides on class I MHC folding and assembly in an in vitro translation system. *J Immunol.* 1992; 149:2935–2944. [PubMed: 1401922]

2. Neefjes JJ, Hammerling GJ, Momburg F. Folding and assembly of major histocompatibility complex class I heterodimers in the endoplasmic reticulum of intact cells precedes the binding of peptide. *J Exp Med.* 1993; 178:1971–1980. [PubMed: 8245776]
3. Chapman DC, Williams DB. ER quality control in the biogenesis of MHC class I molecules *Semin. Cell Dev Biol.* 2010; 21:512–519.
4. Lie WR, Myers NB, Connolly JM, Gorka J, Lee DR, Hansen TH. The specific binding of peptide ligand to L^d class I major histocompatibility complex molecules determines their antigenic structure. *J Exp Med.* 1991; 173:449–459. [PubMed: 1703208]
5. Solheim JC, Carreno BM, Smith JD, et al. Binding of peptides lacking consensus anchor residue alters H-2L^d serologic recognition. *J Immunol.* 1993; 151:5387–5397. [PubMed: 7693810]
6. Smith JD, Solheim JC, Carreno BM, Hansen TH. Characterization of class I MHC folding intermediates and their disparate interactions with peptide and beta 2-microglobulin. *Mol Immunol.* 1995; 32:531–540. [PubMed: 7783756]
7. Smith JD, Myers NB, Gorka J, Hansen TH. Model for the in vivo assembly of nascent L^d class I molecules and for the expression of unfolded L^d molecules at the cell surface. *J Exp Med.* 1993; 178:2035–2046. [PubMed: 8245780]
8. Myers NB, Harris MR, Connolly JM, Lybarger L, Yu YY, Hansen TH. K^b, K^d, and L^d molecules share common tapasin dependencies as determined using a novel epitope tag. *J Immunol.* 2000; 165:5656–5663. [PubMed: 11067922]
9. Smith JD, Lie WR, Gorka J, Kindle CS, Myers NB, Hansen TH. Disparate interaction of peptide ligand with nascent versus mature class I major histocompatibility complex molecules: comparisons of peptide binding to alternative forms of L^d in cell lysates and the cell surface. *J Exp Med.* 1992; 175:191–202. [PubMed: 1730917]
10. Yu YY, Myers NB, Hilbert CM, Harris MR, Balendiran GK, Hansen TH. Definition and transfer of a serological epitope specific for peptide-empty forms of MHC class I. *Int Immunol.* 1999; 11:1897–1906. [PubMed: 10590255]
11. Harris MR, Lybarger L, Myers NB, et al. Interactions of HLA-B27 with the peptide loading complex as revealed by heavy chain mutations. *Int Immunol.* 2001; 13:1275–1282. [PubMed: 11581172]
12. Barnden MJ, Purcell AW, Gorman JJ, McCluskey J. Tapasin-mediated retention and optimization of peptide ligands during the assembly of class I molecules. *J Immunol.* 2000; 165:322–330. [PubMed: 10861068]
13. Garbi N, Tan P, Diehl AD, et al. Impaired immune responses and altered peptide repertoire in tapasin-deficient mice. *Nat Immunol.* 2000; 1:234–238. [PubMed: 10973281]
14. Barber LD, Howarth M, Bowness P, Elliott T. The quantity of naturally processed peptides stably bound by HLA-A*0201 is significantly reduced in the absence of tapasin. *Tissue Antigens.* 2001; 58:363–368. [PubMed: 11929586]
15. Williams AP, Peh CA, Purcell AW, McCluskey J, Elliott T. Optimization of the MHC class I peptide cargo is dependent on tapasin. *Immunity.* 2002; 16:509–520. [PubMed: 11970875]
16. Chen M, Bouvier M. Analysis of interactions in a tapasin/class I complex provides a mechanism for peptide selection. *EMBO J.* 2007; 26:1681–1690. [PubMed: 17332746]
17. Praveen PV, Yaneva R, Kalbacher H, Springer S. Tapasin edits peptides on MHC class I molecules by accelerating peptide exchange. *Eur J Immunol.* 2010; 40:214–224. [PubMed: 20017190]
18. Garstka MA, Fritzsche S, Lenart I, et al. Tapasin dependence of major histocompatibility complex class I molecules correlates with their conformational flexibility. *FASEB J.* 2011
19. Ozato K, Mayer NM, Sachs DH. Monoclonal antibodies to mouse major histocompatibility complex antigens. *Transplantation.* 1982; 34:113–120. [PubMed: 7135466]
20. Straus DS, Stroynowski I, Schiffer SG, Hood L. Expression of hybrid class I genes of the major histocompatibility complex in mouse L cells. *Proc Natl Acad Sci U S A.* 1985; 82:6245–6249. [PubMed: 3862131]
21. Stroynowski I, Forman J, Goodenow RS, et al. Expression and T cell recognition of hybrid antigens with amino-terminal domains encoded by Qa-2 region of major histocompatibility complex and carboxyl termini of transplantation antigens. *J Exp Med.* 1985; 161:935–952. [PubMed: 2580938]

22. Nieto MC, Song ES, McKinney D, McMillan M, Goodenow RS. The association of H-2L^d with human beta-2 microglobulin induces localized conformational changes in the alpha-1 and-2 superdomain. *Immunogenetics*. 1989; 30:361–369. [PubMed: 2478461]
23. Solheim JC, Carreno BM, Myers NB, Lee DR, Hansen TH. Peptide-induced rescue of serologic epitopes on class I MHC molecules. *J Immunol*. 1995; 154:1188–1197. [PubMed: 7529793]
24. Simone LC, Wang X, Tuli A, McIlhaney MM, Solheim JC. Influence of the tapasin C terminus on the assembly of MHC class I allotypes. *Immunogenetics*. 2009; 61:43–54. [PubMed: 18958466]
25. Wang X, Simone LC, Tuli A, Solheim JC. Comparative analysis of the impact of a free cysteine in tapasin on the maturation and surface expression of murine MHC class I allotypes. *Int J Immunogenet*. 2009; 36:183–187. [PubMed: 19490214]
26. Shiroishi T, Evans GA, Appella E, Ozato K. In vitro mutagenesis of a mouse MHC class I gene for the examination of structure-function relationships. *J Immunol*. 1985; 134:623–629. [PubMed: 3964822]
27. Yu YY, Turnquist HR, Myers NB, Balendiran GK, Hansen TH, Solheim JC. An extensive region of an MHC class I alpha 2 domain loop influences interaction with the assembly complex. *J Immunol*. 1999; 163:4427–4433. [PubMed: 10510384]
28. Ozato K, Hansen TH, Sachs DH. Monoclonal antibodies to mouse MHC antigens. II. Antibodies to the H-2L^d antigen, the products of a third polymorphic locus of the mouse major histocompatibility complex. *J Immunol*. 1980; 125:2473–2477. [PubMed: 7191868]
29. Evans GA, Margulies DH, Shykind B, Seidman JG, Ozato K. Exon shuffling: mapping polymorphic determinants on hybrid mouse transplantation antigens. *Nature*. 1982; 300:755–757. [PubMed: 6184620]
30. Simone LC, Wang X, Tuli A, Solheim JC. Effect of a tapasin mutant on the assembly of the mouse MHC class I molecule H2-K^d. *Immunol Cell Biol*. 2010; 88:57–62. [PubMed: 19687800]
31. Balendiran GK, Solheim JC, Young AC, Hansen TH, Nathenson SG, Sacchettini JC. The three-dimensional structure of an H-2L^d-peptide complex explains the unique interaction of L^d with beta-2 microglobulin and peptide. *Proc Natl Acad Sci U S A*. 1997; 94:6880–6885. [PubMed: 9192660]
32. Mitaksov V, Fremont DH. Structural definition of the H-2K^d peptide-binding motif. *J Biol Chem*. 2006; 281:10618–10625. [PubMed: 16473882]
33. Kabsch W. A solution for the best rotation to relate two sets of vectors. *Acta Crystallographica Section A*. 1976; 32:922–923.
34. Collaborative Computational Project, Number 4. The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr*. 1994; 50:760–763. [PubMed: 15299374]
35. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr*. 2010; 66:486–501. [PubMed: 20383002]
36. Schrödinger LLC. The PyMOL Molecular Graphics System, Version 1.3. 2010
37. Ortmann B, Androlewicz MJ, Cresswell P. MHC class I/beta 2-microglobulin complexes associate with TAP transporters before peptide binding. *Nature*. 1994; 368:864–867. [PubMed: 8159247]
38. Suh WK, Cohen-Doyle MF, Fruh K, Wang K, Peterson PA, Williams DB. Interaction of MHC class I molecules with the transporter associated with antigen processing. *Science*. 1994; 264:1322–1326. [PubMed: 8191286]
39. Elliott T. How does TAP associate with MHC class I molecules? *Immunol Today*. 1997; 18:375–379. [PubMed: 9267079]
40. Harris MR, Yu YY, Kindle CS, Hansen TH, Solheim JC. Calreticulin and calnexin interact with different protein and glycan determinants during the assembly of MHC class I. *J Immunol*. 1998; 160:5404–5409. [PubMed: 9605141]
41. Rizvi SM, Raghavan M. Direct peptide-regulatable interactions between MHC class I molecules and tapasin. *Proc Natl Acad Sci U S A*. 2006; 103:18220–18225. [PubMed: 17116884]
42. Ljunggren HG, Stam NJ, Ohlen C, et al. Empty MHC class I molecules come out in the cold. *Nature*. 1990; 346:476–480. [PubMed: 2198471]
43. Mahmutefendic H, Blagojevic G, Tomas MI, Kucic N, Lucin P. Segregation of open Major Histocompatibility Class I conformers at the plasma membrane and during endosomal trafficking

- reveals conformation-based sorting in the endosomal system. *Int J Biochem Cell Biol.* 2011; 43:504–515. [PubMed: 21176792]
44. Speir JA, Garcia KC, Brunmark A, et al. Structural basis of 2C TCR allorecognition of H-2Ld peptide complexes. *Immunity.* 1998; 8:553–562. [PubMed: 9620676]
 45. Hansen T, Balendiran G, Solheim J, Ostrov D, Nathenson S. Structural features of MHC class I molecules that might facilitate alternative pathways of presentation. *Immunol Today.* 2000; 21:83–88. [PubMed: 10652466]
 46. Maryanski JL, Romero P, Van Pel A, et al. The identification of tyrosine as a common key residue in unrelated H-2K^d restricted antigenic peptides. *Int Immunol.* 1991; 3:1035–1042. [PubMed: 1721832]
 47. Romero P, Corradin G, Luescher IF, Maryanski JL. H-2K^d-restricted antigenic peptides share a simple binding motif. *J Exp Med.* 1991; 174:603–612. [PubMed: 1714934]
 48. Quesnel A, Casrouge A, Kourilsky P, Abastado JP, Trudelle Y. Use of synthetic peptide libraries for the H-2K^d binding motif identification. *Pept Res.* 1995; 8:44–51. [PubMed: 7756754]
 49. Suri A, Walters JJ, Levisetti MG, Gross ML, Unanue ER. Identification of naturally processed peptides bound to the class I MHC molecule H-2K^d of normal and TAP-deficient cells. *Eur J Immunol.* 2006; 36:544–557. [PubMed: 16479539]
 50. Kornfeld R, Kornfeld S. Assembly of asparagine-linked oligosaccharides. *Annu Rev Biochem.* 1985; 54:631–664. [PubMed: 3896128]
 51. Krangel MS, Orr HT, Strominger JL. Assembly and maturation of HLA-A and HLA-B antigens in vivo. *Cell.* 1979; 18:979–991. [PubMed: 93026]
 52. Owen MJ, Kissonerghis AM, Lodish HF. Biosynthesis of HLA-A and HLA-B antigens in vivo. *J Biol Chem.* 1980; 255:9678–9684. [PubMed: 7000762]
 53. Corr M, Boyd LF, Frankel SR, Kozlowski S, Padlan EA, Margulies DH. Endogenous peptides of a soluble major histocompatibility complex class I molecule, H-2L^ds: sequence motif, quantitative binding, and molecular modeling of the complex. *J Exp Med.* 1992; 176:1681–1692. [PubMed: 1281216]
 54. Yui K, Hashimoto Y, Wadsworth S, Greene MI. Characterization of Lyt-2-, L3T4- class I-specific cytolytic clones in C3H-gld/gld mice. Implications for functions of accessory molecules and programmed development. *J Exp Med.* 1987; 166:1026–1040. [PubMed: 3498785]
 55. Feuerbach D, Etteldorf S, Ebenau-Jehle C, Abastado JP, Madden D, Burgert HG. Identification of amino acids within the MHC molecule important for the interaction with the adenovirus protein E3/19K. *J Immunol.* 1994; 153:1626–1636. [PubMed: 8046235]
 56. Hauptfeld V, Nahm M, Hauptfeld M, Shreffler DC. Monoclonal antibodies to mouse MHC antigens. I. Serologic characterization of ten anti-H-2 and anti-Ia reagents. *Immunogenetics.* 1984; 19:169–173. [PubMed: 6583159]
 57. Koeller D, Lieberman R, Miyazaki J, et al. Introduction of H-2D^d determinants into the H-2L^d antigen by site-directed mutagenesis. *J Exp Med.* 1987; 166:744–760. [PubMed: 2442290]
 58. Saper MA, Bjorkman PJ, Wiley DC. Refined structure of the human histocompatibility antigen HLA-A2 at 2.6 Å resolution. *J Mol Biol.* 1991; 219:277–319. [PubMed: 2038058]
 59. Lewis JW, Elliott T. Evidence for successive peptide binding and quality control stages during MHC class I assembly. *Curr Biol.* 1998; 8:717–720. [PubMed: 9637925]
 60. Van Hateren A, James E, Bailey A, Phillips A, Dalchau N, Elliott T. The cell biology of major histocompatibility complex class I assembly: towards a molecular understanding. *Tissue Antigens.* 2010; 76:259–275. [PubMed: 21050182]
 61. Zacharias M, Springer S. Conformational flexibility of the MHC class I alpha1-alpha2 domain in peptide bound and free states: a molecular dynamics simulation study. *Biophys J.* 2004; 87:2203–2214. [PubMed: 15454423]
 62. Lewis JW, Neisig A, Neeffes J, Elliott T. Point mutations in the alpha 2 domain of HLA-A2.1 define a functionally relevant interaction with TAP. *Curr Biol.* 1996; 6:873–883. [PubMed: 8805302]
 63. Peace-Brewer AL, Tussey LG, Matsui M, Li G, Quinn DG, Frelinger JA. A point mutation in HLA-A*0201 results in failure to bind the TAP complex and to present virus-derived peptides to CTL. *Immunity.* 1996; 4:505–514. [PubMed: 8630735]

64. Dong G, Wearsch PA, Peaper DR, Cresswell P, Reinisch KM. Insights into MHC class I peptide loading from the structure of the tapasin-ERp57 thiol oxidoreductase heterodimer. *Immunity*. 2009; 30:21–32. [PubMed: 19119025]
65. Smith MJ, Basora T, Kieran JE, Nieto MC. Alteration within a discrete region of the H-2L^d alpha 1 helix upon association with human beta 2 microglobulin. *Immunol Cell Biol*. 1993; 71(Pt 2): 145–149. [PubMed: 8486398]
66. Zhang W, Wearsch PA, Zhu Y, Leonhardt RM, Cresswell P. A role for UDP-glucose glycoprotein glucosyltransferase in expression and quality control of MHC class I molecules. *Proc Natl Acad Sci U S A*. 2011
67. Paulsson KM, Jevon M, Wang JW, Li S, Wang P. The double lysine motif of tapasin is a retrieval signal for retention of unstable MHC class I molecules in the endoplasmic reticulum. *J Immunol*. 2006; 176:7482–7488. [PubMed: 16751394]
68. Garstka M, Borchert B, Al-Balushi M, et al. Peptide-receptive major histocompatibility complex class I molecules cycle between endoplasmic reticulum and cis-Golgi in wild-type lymphocytes. *J Biol Chem*. 2007; 282:30680–30690. [PubMed: 17656363]
69. Leonhardt RM, Fiegl D, Rufer E, Karger A, Bettin B, Knittler MR. Post-endoplasmic reticulum rescue of unstable MHC class I requires proprotein convertase PC7. *J Immunol*. 2010; 184:2985–2998. [PubMed: 20164418]
70. McKinney DM, Chen P, McMillan M. Single D^d amino acid substitutions in the H-2L^d molecule identify antibody epitopes. *Immunogenetics*. 1994; 39:440–443. [PubMed: 7514573]

Figure 1. Tapasin preferentially interacts with the 34-1-2⁺ conformation of K^d over the open, peptide-free conformation

Either L^d or epitope-tagged K^d were stably expressed in HeLa cells (HeLa-L^d and HeLa-K^d, respectively) and in L-cell fibroblasts (L-L^d and L-K^d, respectively). Cell lysates were immunoprecipitated with mAb 64-3-7 (for open, peptide-free L^d and K^d), 34-1-2 (for K^d), or 30-5-7 (for folded, peptide-bound L^d). Immunoprecipitation of untransfected cell lysates served as negative controls. The precipitated proteins were electrophoresed on 4→20% acrylamide Tris-glycine SDS gels, transferred to membranes, and probed with 64-3-7 mAb to visualize L^d and K^d heavy chains (A–D, top panel). To visualize MHC class I-associated tapasin, equivalent amounts of immunoprecipitated heavy chain proteins were electrophoresed on 10% acrylamide Tris-glycine SDS gels, transferred to membranes, and probed with mouse anti-human tapasin mAb (A and C, bottom panel) or Armenian hamster antimouse tapasin mAb (B and D, middle panel and bottom panel). In the experiment shown in (C), equivalent loading of immunoprecipitated K^d molecules was verified by the heavy chain blot shown in the middle panel. For immunoprecipitation experiments in L-cells, the immunoprecipitated proteins were reduced with dithiothreitol, which accounts for the

reduced antibody heavy chain band seen on the western blots. In **(C)** a non-specific band was seen in all of the samples when probed for tapasin on western blot. In **(B)** and **(D)**, on shorter exposures of the western blots to film, a weak tapasin band was apparent in association with 64-3-7⁺ L^d and 34-1-2⁺ K^d (middle panel), respectively, but with prolonged exposure a strong tapasin band could be detected (bottom panel). The specific band corresponding to tapasin in **(B)**, **(C)** and **(D)** is highlighted with a bold arrowhead.

Figure 2. TAP and ERp57 co-precipitate with the 34-1-2⁺ conformation of K^d

Components of the peptide-loading complex, TAP and ERp57, bind to the open, peptide-free form of L^d and to the 34-1-2⁺ form of K^d. Cell lysates of HeLa-K^d and HeLa-L^d cells were immunoprecipitated with mAb 64-3-7 (for open, peptide-free L^d and K^d), 34-1-2 (for K^d), or 30-5-7 (for folded, peptide-bound L^d). Untransfected HeLa cell lysates were used as negative controls. The precipitated proteins were electrophoresed on 4→20% acrylamide Tris-glycine SDS gels, transferred to membranes, and probed with 64-3-7 mAb to visualize immunoprecipitated L^d and K^d heavy chains (data not shown). To visualize ERp57 and TAP molecules co-immunoprecipitated with MHC heavy chains, equivalent amounts of immunoprecipitated MHC heavy chain molecules were electrophoresed on 10% acrylamide Tris-glycine SDS gels, transferred to membranes, and probed with rabbit anti-serum to human TAP1 (**A** and **C**) and with rabbit anti-serum to human ERp57 (**B** and **D**). A non-specific band (marked with an asterisk) was present when 30-5-7 and 34-1-2 immunoprecipitates were probed on a western blot for ERp57.

A

B

Figure 3. The 34-1-2⁺ L^d molecules are stabilized at the cell surface by 25°C incubation and display an exceptionally fast rate of cell surface decay at 37°C

(A) HeLa-L^d and HeLa-K^d cells were incubated either at 37°C or at 25°C overnight prior to flow cytometric analysis with the 64-3-7, 30-5-7, and 34-1-2 mAbs. Each sample was prepared in triplicate, and the data are represented as the average mean fluorescence units $\pm$ the standard deviation. (B) HeLa-L^d and HeLa-K^d cells (incubated at 37°C) were treated with 2 μ g/ml Bfa for the indicated times prior to flow cytometric analysis with the 64-3-7, 30-5-7, and 34-1-2 mAbs. The level of each serologic conformer present on the cells at 0 hr Bfa treatment was set to 100% and used to calculate the level of each serologic conformer that remained at each subsequent time point. The cell surface half life (t_{1/2}) of the 34-1-2⁺ K^d and 30-5-7⁺ L^d molecules was determined by fitting the data to the exponential decay equation ($[A] = [A]_0 e^{-at}$), such that t_{1/2} is equal to $\ln(2)/a$. The decay of the 34-1-2⁺ L^d

molecules was better described by the equation $[B] = [B]_0 e^{-bt} + c(e^{-at} - e^{-bt})$, which allows for the generation of 34-1-2⁺ conformers from the melting of folded, 30-5-7⁺ L^d molecules. The $t_{1/2}$ of 34-1-2⁺ L^d molecules was calculated by setting $[B] = (1/2)[B]_0$ and solving for t numerically.

Figure 4. The 34-1-2⁺ L^d molecules are predominantly immature and display a strong association with tapasin

(A) HeLa-L^d cells were treated with IFN- γ and the cell lysates were immunoprecipitated with the 64-3-7, 34-1-2, or 30-5-7 mAb. For the immunoprecipitation with the 64-3-7 or 30-5-7 mAb, 1×10^7 cells were used, whereas 4×10^7 cells were used for the 34-1-2 immunoprecipitation. Immunoprecipitated proteins were mock treated or treated with Endo H and the samples were electrophoresed on a 4 $\rightarrow$ 20% acrylamide Tris-glycine SDS gel, transferred to a membrane, and probed with 64-3-7 mAb to visualize the immunoprecipitated L^d heavy chain. Equivalent numbers of untransfected HeLa cells treated with IFN- γ were used as negative controls. The top panel shows a long exposure of the

western blot to film in order to reveal the low level of 34-1-2⁺ L^d. A non-specific band which migrates just above the MHC class I heavy chain band in the upper panel is marked with an asterisk. **(B)** Equivalent amounts of the L^d heavy chain immunoprecipitated by the indicated mAb from IFN- γ -treated HeLa-L^d cells were electrophoresed on a 4 $\rightarrow$ 20% acrylamide Tris-glycine SDS gel (for the L^d heavy chain) or on a 10% acrylamide Tris-glycine SDS gel (for tapasin), transferred to membranes, and probed with 64-3-7 mAb (for the L^d heavy chain blot) or with mouse anti-human tapasin mAb. HeLa cells not treated with IFN- γ were used as negative controls. **(C)** Densitometry of the L^d heavy chain bands shown in **(B)**. L^d molecules with mature glycosylation migrate slower than do immature L^d glycoproteins. Thus, to calculate the percentage of L^d molecules with mature and immature glycosylation that were precipitated by each mAb, the level of the slow and fast migrating bands was divided by the total amount of L^d heavy chain. The background band (apparent in HeLa cells not expressing L^d), which migrates at the same position as the mature L^d molecules, was subtracted for this analysis.

Figure 5. The 34-1-2⁺ L^d molecules overlap with the pools of 64-3-7⁺ L^d and 30-5-7⁺ L^d, and addition of an L^d-specific peptide induces folding of the 64-3-7⁺34-1-2⁺ L^d molecules
 Lysates of equivalent numbers of IFN- γ -treated HeLa-L^d cells were incubated at 4°C overnight in the absence or presence of 500 μ M P29 peptide (YPNVNIHNF). Following the overnight incubation, the lysates were centrifuged to pellet the cell nuclei, and the supernatants were divided into three aliquots. The aliquots were either mock depleted, or subjected to 14 consecutive rounds of immunoprecipitation with either the 64-3-7 mAb (**A**) or the 30-5-7 mAb (**B**). Following the mock depletion, 64-3-7 depletion, or 30-5-7 depletion, each sample was immunoprecipitated with the 34-1-2 mAb (**C**). An equivalent number of IFN- γ -treated HeLa cells (not expressing K^d or L^d) were immunoprecipitated with either 64-3-7 (**A**, lane 7), 30-5-7 (**B**, lane 7), or 34-1-2 (**C**, lane 8) and served as negative controls. Immunoprecipitated proteins were electrophoresed on 4 $\rightarrow$ 20% acrylamide Tris-glycine SDS gels, transferred to membranes, and probed with the 64-3-7 mAb to visualize the L^d heavy chain. (**D**) Model of the overlap between the 64-3-7⁺, 34-1-2⁺, and 30-5-7⁺ L^d serologic forms, and the influence of the P29 peptide on these forms. (Areas are not intended as representative of the actual amount of each L^d serologic form.)

Figure 6. Location of antibody binding sites, the proposed tapasin binding site, and peptide-heavy chain hydrogen bonds within the MHC class I peptide-binding groove
(A) Ribbon diagram of the $\alpha 1/\alpha 2$ domains of the L^d molecule (Protein Data Bank ID 1LD9) (31). Residues 128–149 of the MHC class I $\alpha 2$ domain (shown in red) have been proposed to associate with a conserved surface on tapasin (39,64). The 34-1-2 epitope (highlighted in yellow) is located within residues 63–85 (22,23,57). Also shown are the positions of residue 70 (triangle) and residue 73 (circle) which are important for 34-1-2 mAb recognition (57,65). Residues 107, 169, and 173 are part of the 30-5-7 epitope (shown in magenta) (23,70). The 64-3-7 mAb epitope maps to residues 46-52 (shown in blue) (8). **(B)** Superposition of the crystal structures of the K^d -TYQRTRALV peptide (magenta) (Protein

Data Bank ID 2FWO) (32) and the L^d-YPNVNIHNF peptide (green) (Protein Data Bank ID 1LD9) (31) complexes. Aspartic acid 70 in K^d forms two hydrogen bonds with the peptide: one occurring between Oδ2 and the hydroxyl group of P2 tyrosine, and the other between Oδ1 and the hydroxyl of P5 threonine. Glutamine 70 in L^d forms one hydrogen bond between Nε2 and P5 Oδ1. The side chain of tryptophan 73 in both K^d and L^d is also shown.