Obstructive Sleep Apnea Syndrome and Perioperative Complications: A Systematic Review of the Literature

Tajender S. Vasu, M.D., M.S.¹; Ritu Grewal, M.D.²; Karl Doghramji, M.D.²

¹Division of Pulmonary, Critical Care, and Sleep Medicine, Stony Brook University Medical Center, Stony Brook, NY;

²Division of Sleep Medicine, Thomas Jefferson University Hospital, Philadelphia, PA

Obstructive sleep apnea syndrome (OSAS) is a common sleep related breathing disorder. Its prevalence is estimated to be between 2% and 25% in the general population. However, the prevalence of sleep apnea is much higher in patients undergoing elective surgery. Sedation and anesthesia have been shown to increase the upper airway collapsibility and therefore increasing the risk of having postoperative complications in these patients. Furthermore, the majority of patients with sleep apnea are undiagnosed and therefore are at risk during the perioperative period. It is important to identify these patients so that appropriate actions can be taken in a timely fashion. In this review article, we will discuss the epidemiology of sleep apnea in the

surgical population. We will also discuss why these patients are at a higher risk of having postoperative complications, with the special emphasis on the role of anesthesia, opioids, sedation, and the phenomenon of REM sleep rebound. We will also review how to identify these patients preoperatively and the steps that can be taken for their perioperative management.

Keywords: Obstructive sleep apnea syndrome, perioperative complications, postoperative complications, STOP Questionnaire, Berlin Questionnaire

Citation: Vasu TS; Grewal R; Doghramji K. Obstructive sleep apnea syndrome and perioperative complications: a systematic review of the literature. *J Clin Sleep Med* 2012;8(2):199-207.

Obstructive sleep apnea is characterized by intermittent and recurrent episodes of partial or complete obstruction of the upper airway during sleep. These episodes disrupt sleep architecture, causing fragmented sleep and daytime sleepiness. OSAS has been shown to be associated with various health-related consequences, including increased rate of motor vehicle accidents, hypertension, diabetes mellitus, congestive heart failure, stroke, and all-cause mortality.¹⁻⁹

Recently, numerous studies have demonstrated that surgical patients with sleep apnea are at increased risk of having perioperative complications, including hypoxemia, pneumonia, difficult intubation, myocardial infarction, pulmonary embolism, atelectasis, cardiac arrhythmias, and unanticipated admission to the ICU. The majority of patients with OSA are undiagnosed upon admission and are at risk during the perioperative period, presumably due to their underlying sleep apnea. Therefore, it is very important to identify these patients preoperatively so that one can initiate appropriate perioperative measures.

Epidemiology

OSAS is an extremely common sleep related breathing disorder, and its prevalence has been increasing throughout the world because of obesity and increasing age of the general population. Its prevalence is between 2% and 25% in the general population, depending upon how sleep apnea is defined. In an epidemiological study, Young et al. noted that the prevalence of sleep apnea, defined as apnea-hypopnea index (AHI) $\geq 5/h$ was 9% for women and 24% for men.¹⁰ However, the prevalence of OSAS (defined as AHI $\geq 5/h$ and daytime sleepiness) was 2% in women and 4% in men.¹⁰ The National Sleep Foundation (NSF)

Sleep in America 2005 Poll found that 1 in 4 Americans are at high risk of having sleep apnea based on the Berlin Questionnaire.¹¹ The prevalence of sleep apnea is much higher in surgical patients and depends on the type of surgery. In the bariatric surgery population, the prevalence of sleep apnea has been found to be $> 70\%$.^{12,13} It is the standard of care for these patients to get a formal sleep evaluation prior to undergoing the bariatric surgery. However, patients who are coming for general surgery also have a higher prevalence of sleep apnea.¹⁴ Chung et al. used the Berlin Questionnaire preoperatively and found that 24% of surgical patients were at high risk for sleep apnea.¹⁵ We used the STOP-BANG Questionnaire in our elective surgical population and found that 41% of patients were at high risk for sleep apnea based on the questionnaire.¹⁶ In a cross-sectional study, 39 patients underwent nocturnal polysomnography (NPSG) prior to undergoing epilepsy surgery. It was found that 1 in 3 patients undergoing epilepsy surgery had sleep apnea.¹⁷ In another study, the prevalence of sleep apnea was found to be 64% in a small population of patients undergoing surgery for intracranial tumor.¹⁸ The majority of these patients are unaware of their sleep apnea prior to undergoing elective surgery. In an observational study, Finkel et al. noted that $> 80\%$ of surgical patients were unaware that they had sleep apnea prior to undergoing surgery.¹⁴

Pathophysiology

Surgical patients receive sedation, anesthesia, and opioids during the perioperative period. These medicines have been shown to increase pharyngeal collapse, decrease ventilatory response, and impair the arousal response, leading to worsening of sleep apnea in the perioperative period.

Impact of Sedation, Anesthesia, and Opioids

Patients with sleep apnea have recurrent episodes of partial or complete obstruction of the upper airway during sleep. These episodes usually occur when the negative pressure of inspiratory muscles exceeds the upper airway dilator muscle activity (critical airway pressure).^{19,20} General anesthetics have been shown to decrease the upper airway dilator muscle activity in a dose-dependent manner and thereby increase upper airway collapsibility.²¹⁻²³ In 12 healthy subjects undergoing minor surgery, increasing depth of propofol anesthesia was associated with a progressive increase in critical airway pressure and upper airway collapsibility.^{23,24} This increased upper airway collapsibility was found to be secondary to progressive decrease in the genioglossus muscle activity. Upper airway collapsibility may cause worsening of the sleep apnea and increase the risk of hypoxemia, cardiac arrhythmias, and postoperative complications.

Anesthetic medicines also impair the arousal response, a protective defense mechanism against sleep apnea that helps in overcoming the airway obstruction. Anesthetics, opioids, hypnotics, and benzodiazepines may also cause respiratory depression and thereby decrease the minute ventilation. Studies have shown that halothane reduces the ventilatory response to hypoxemia and hypercapnia in humans.^{25,26} This depression is most likely secondary to a selective effect of halothane on the peripheral chemoreflex loop. Similarly, a subanesthetic dose of isoflurane has been shown to reduce the hypoxic ventilatory response via peripheral chemoreceptors.^{27,28}

Patients undergoing surgery frequently receive opioids for the pain control. Opioids have been shown to impair ventilatory function by affecting both peripheral and central carbon dioxide chemoreflex loops.^{29,30} Studies have shown that small doses of narcotics administered epidurally may also depress the respiratory function, even in healthy adults.³¹⁻³³ The ventilatory depression of opiates appears to be affected by sex and ethnicity, as well.^{34,35} Morphine has been shown to reduce hypoxic and hypercapnic ventilatory response in women, but not in men.³⁴ On the other hand, it raises the apneic threshold in men with no effect in women.³⁴ The combination of opiates and benzodiazepines has been shown to cause more significant episodes of hypoxemia and apnea.³⁶ This is most likely secondary to significant reduction of hypoxic ventilatory response to both opiates and benzodiazepines.³⁷⁻³⁹

REM Sleep Rebound

Surgical patients have been shown to have highly fragmented sleep on postoperative nights 1 or 2, with a significant reduction in REM sleep, slow wave sleep, and increased stage 2 NREM sleep.⁴⁰⁻⁴⁶ These sleep disturbances usually occur secondary to surgical stress, pain, and the use of anesthetic and pain medications.^{40,47,48} The stress of surgical trauma leads to increased level of cortisol, and cortisol has been shown to cause significant reduction in REM sleep.^{49,50} Surgical trauma has also been shown to induce significant inflammatory response, characterized by the increased level of pro-inflammatory markers like tumor necrosis factor alpha (TNF- α), interleukin 1 (IL-1), and IL-6.⁵¹⁻⁵³ These inflammatory markers, especially IL-1 and TNF- α , have been shown to suppress REM sleep.⁵⁴⁻⁵⁶

There are studies that have shown that REM sleep is usually absent on postoperative nights 1 and 2. This is usually followed by a profound increase in the amount and density of REM sleep (REM sleep rebound) during recovery nights 3 to 5.^{44,57-59} The episodes of sleep disordered breathing and hypoxemia are usually worse during REM sleep due to hypotonia and unstable breathing. REM sleep is also associated with increased sympathetic discharge leading to tachycardia, hemodynamic instability, and myocardial ischemia.⁶⁰⁻⁶⁴

It is well known that most complications after the surgery occur in the first postoperative week, especially between postoperative days 2 and 5, corresponding to periods of REM rebound. Episodes of hypoxemia after surgery have been reported to occur mostly between postoperative nights 2 to 5.^{58,59,65} These episodes may increase the risk of wound infection, cerebral dysfunction, and cardiac arrhythmias.⁶⁶ In an observational study at Mayo Clinic, it was found that the incidence of acute myocardial infarction peaked on day 3 after surgery.⁶⁷ Similarly, episodes of delirium, nightmares, and psychomotor dysfunction have been reported to occur between postoperative nights 3 to 5.⁶⁸⁻⁷⁰

Evidence on Sleep Apnea as a Risk Factor for Perioperative Complications

We systematically searched the literature on PubMed, Embase, and Scopus databases to identify relevant studies on association between obstructive sleep apnea and perioperative outcome. We included studies conducted in adults who underwent elective surgery. We excluded bariatric surgery and sleep apnea surgery population. Two authors (TSV and RG) independently searched for the relevant articles published in the English literature from 1966 to December 2011. We used the combination of terms including surgery, perioperative outcome, perioperative complications, perioperative pulmonary outcome, perioperative risk, postoperative complications, and obstructive sleep apnea. Bibliographies of all selected articles and review articles were also reviewed to find any other relevant article. These 2 authors (TSV and RG) also independently assigned the Oxford level of evidence and the grade of recommendation to each article.⁷¹ There was 100% agreement between the 2 authors. We identified 11 articles on obstructive sleep apnea and perioperative outcome. These articles along with their Oxford level of evidence are reported in the **Table 1** and **Table 2**.

Surgical patients are at higher risk of having complications for a variety of reasons, including ASA (American Society of Anesthesiologists) class,^{72,73} age,⁷³⁻⁷⁶ type of paralytics,^{77,78} current smoking,⁷⁹⁻⁸¹ low albumin,⁸¹⁻⁸³ duration of surgery,⁸⁴⁻⁸⁶ type of anesthesia,^{73,78,87,88} and other comorbidities—especially chronic obstructive pulmonary disease, coronary artery disease, and renal failure.^{79,80,89} The risk of postoperative complications also depends on the type of surgery. The rate of complication is higher in patients undergoing abdominal surgery⁹⁰⁻⁹² and is also increased with aortic aneurysm repair,⁹³⁻⁹⁶ vascular,^{74,78,79,93} thoracic,^{83,90,93,96} and neck surgery.^{83,96,97}

Gupta et al. have shown an increased risk of postoperative complications (39% vs 18%), higher rate of transfer to ICU (24% vs 9%), and increased length of hospital stay in patients with obstructive sleep apnea compared with control subjects matched for age, sex, and body mass index (BMI).⁹⁸ In this study, these investigators also reported that OSA patients who

Table 1—Studies reporting association between obstructive sleep apnea and perioperative complications

Author	Type of Study	Number of Patients	Diagnosis of OSAS	Type of Surgeries	Complications	Results
Gupta et al. ⁹⁸	Case control study	101 patients with OSA and 101 matched controls	Polysomnography (PSG)	Orthopedic (hip or knee replacement)	Reintubation, hypoxemia, acute hypercapnia, myocardial infarction, arrhythmia, delirium, and ICU transfer	Patients with OSA had higher rate of postoperative complications (39% vs 18%). These patients also had increased hospital length of stay.
Auckley et al. ¹⁰⁵	Historical cohort study	81 patients with completed Berlin Questionnaire	Berlin Questionnaire	Elective surgery (type of surgeries is not included in the abstract)	Hypoxemia, hypercapnia, reintubation, atelectasis, pneumonia, arrhythmia, thromboembolism	Patients with high-risk of sleep apnea based on the Berlin Questionnaire had a higher rate of postoperative complications (20% vs 4.5%).
Sabers et al. ¹⁰⁶	Case control study	234 patients with OSA and 234 matched controls	Polysomnography	Non-otorhinolaryngologic outpatient surgical procedures	Unplanned hospital admission, bronchospasm, upper airway obstruction, hypotension, atrial fibrillation, pulmonary edema	No significant difference in the rate of unplanned hospital admissions (23.9% vs 18.8%) or other adverse events (2.1% vs 1.3%)
Kaw et al. ¹⁰⁰	Case control study	37 patients with OSA and 185 matched controls	Polysomnography	Cardiac	Encephalopathy, postoperative infections, and ICU length of stay	Patients with sleep apnea had higher rate of encephalopathy, postoperative infections (mediastinitis), and increased ICU length of stay.
Hwang et al. ¹⁰²	Historical cohort study	172 patients underwent home nocturnal oximetry	Home nocturnal oximetry	Abdominal, ENT, Thoracic, Vascular, Gyn, Neurosurgical, Urologic, Cardiothoracic, and Orthopedic	Arrhythmia, hypoxemia, atelectasis, GI bleed, pneumonia, pulmonary embolism,	Patients with ODI4% $\geq$ 5/h had a higher rate of postoperative complications than those with ODI4% < 5/h (15.3% vs 2.7%).
Gali et al. ¹⁰⁴	Prospective cohort study	693 patients with completed Flemons Criteria and SACS score	Flemons Criteria and SACS score	Orthopedic, Gyn, ENT, Urologic, Thoracic, Plastics, Neurosurgery, General abdominal	Arrhythmia, MI, ICU admission, pneumonia, need for the ventilator support	Postoperative respiratory events were associated with high SACS and PACU events
Liao et al. ⁹⁹	Retrospective matched cohort study	240 patients with OSA and 240 matched controls	International Classification of Disease (ICD-9) codes	Cardiac, ENT, Orthopedic, Spine, Urologic, General, Gyn, and Plastic	Hypoxemia, pulmonary edema, bronchospasm, arrhythmia, confusion	Patients with OSA had a higher incidence of postoperative complications (48% vs 36%)
Vasu et al. ¹⁶	Historical cohort study	135 patients with completed STOP BANG Questionnaire	STOP BANG Questionnaire	Orthopedic, Abdominal, Head and Neck, ENT, Gyn, Vascular, Cardiothoracic	Hypoxemia, pneumonia, pulmonary embolism, atelectasis, hypotension, atrial fibrillation	Patients with high-risk of sleep apnea based on STOP BANG Questionnaire had a higher rate of postoperative complications (19.6% vs 1.3%) and the hospital length of stay.

PSG, polysomnography; ICU, Intensive Care Unit; SACS, Sleep Apnea Clinical Score; PACU, Postanesthesia Care Unit.

Table 1 continues on the following page

Table 1 (continued)—Studies reporting association between obstructive sleep apnea and perioperative complications

Author	Type of Study	Number of Patients	Diagnosis of OSAS	Type of Surgeries	Complications	Results
Stierer et al. ¹⁰⁷	Prospective cohort study	A cohort of 2139 patients who underwent ambulatory surgical procedure	Probability of OSA based on demographic and questionnaire including Maislin index score	Orthopedic, ENT, Gyn, Plastic, Neurologic, Urologic, and general outpatient surgical procedures	Unplanned hospital admission, hypoxemia, cardiac arrhythmia, re-intubation, re-admission within 24 h of discharge, and need for lung ventilation	Increased propensity for OSA was not associated with unplanned hospital admission. However, it was associated with difficult intubation, increased oxygen requirement, and intraoperative tachycardia with increased need for labetalol or metoprolol.
Memtsoudis et al. ¹⁰¹	Case control study	58358 orthopedic patients with OSA and 45547 general surgery patients with OSA were matched for controls in 1:3 manner	International Classification of Disease (ICD-9) codes	Orthopedic and general surgery	Aspiration pneumonia, pulmonary embolism, need for intubation and mechanical ventilation, ARDS	Patients with sleep apnea undergoing orthopedic and general surgeries were at a higher risk of aspiration pneumonia, ARDS, and the need for intubation and mechanical ventilation.
Kaw et al. ¹⁰³	Cohort study	471 patients who underwent non-cardiac surgery within 3 years of PSG	Patients with an apnea-hypopnea index (AHI) $\geq 5/h$ were defined as OSA, and those with AHI < 5 as controls	Non-cardiac surgery	Atrial fibrillation, respiratory failure, hypoxemia, delirium, transfer to ICU, congestive heart failure, myocardial infarction, hospital length of stay	Patients with OSA had a higher rate of postoperative hypoxemia (12.4% vs 2.1%), transfer to ICU (6.7% vs 1.6%), any complication (14.2% vs 2.6%), and hospital length of stay.

PSG, polysomnography; ICU, Intensive Care Unit; SACS, Sleep Apnea Clinical Score; PACU, Postanesthesia Care Unit.

Table 2—The Oxford level of evidence and grade of recommendation for individual studies

Author	Type of Study	Journal	Year of Publication	Oxford Level of Evidence	Grade of Recommendation
Gupta et al. ⁹⁸	Case control study	<i>Mayo Clin Proc</i>	2001	3b	B
Auckley et al. ¹⁰⁵	Historical cohort study	<i>Abstract in Sleep</i>	2003	2b	B
Sabers et al. ¹⁰⁶	Case control study	<i>Anesth Analg</i>	2003	3b	B
Kaw et al. ¹⁰⁰	Case control study	<i>J Cardiovasc Surg</i>	2006	3b	B
Hwang et al. ¹⁰²	Historical cohort study	<i>Chest</i>	2008	2b	B
Gali et al. ¹⁰⁴	Prospective cohort study	<i>Anesthesiology</i>	2009	2b	B
Liao et al. ⁹⁹	Retrospective matched cohort study	<i>Can J Anesth</i>	2009	3b	B
Vasu et al. ¹⁶	Historical cohort study	<i>Arch Otolaryngol Head Neck Surg</i>	2010	2b	B
Stierer et al. ¹⁰⁷	Prospective cohort study	<i>J Clin Sleep Med</i>	2010	2b	B
Memtsoudis et al. ¹⁰¹	Case control study	<i>Anesth Analg</i>	2011	3b	B
Kaw et al. ¹⁰³	Cohort study	<i>Chest</i>	2011	2b	B

received continuous positive airway pressure (CPAP) therapy prior to surgery had a reduced rate of serious complications and a one-day reduction in the length of hospital stay. In another case-control study, Liao et al. found that patients with OSA had higher rate of postoperative complications (44% vs 28%).⁹⁹ Interestingly, they also noted that OSA patients who were not compliant with their CPAP had the highest rate of postoperative complications. Kaw et al. also demonstrated that patients with

OSA had higher incidence of encephalopathy, postoperative infections (mediastinitis), and increased ICU length of stay.¹⁰⁰ Similarly, Memtsoudis et al. in their case-control study found that orthopedic and general surgical patients with sleep apnea were at a higher risk of having perioperative pulmonary complications.¹⁰¹

Hwang et al. demonstrated that the rate of postoperative complications was increased in proportion to episodes of over-

night desaturation during home nocturnal oximetry.¹⁰² In this study, 172 patients underwent home nocturnal oximetry during preoperative evaluation for elective surgery. The number of episodes per hour of oxygen desaturation (or oxygen desaturation index) $\geq 4\%$ (known as the ODI 4%) was calculated for every patient from the home nocturnal oximetry. Patients with an ODI 4% $\geq 5/h$ had a significantly higher rate of postoperative complications than those with ODI 4% $< 5/h$ (15.4% vs 2.7%). Interestingly, the rate of postoperative complications increased with increasing ODI severity. Patients with an ODI 4% of 5-15 had a 13.8% incidence of complications, compared to 17.5% of those with an ODI 4% > 15 .

In a recent cohort study, 471 patients who underwent non-cardiac surgery within 3 years of polysomnography were evaluated for postoperative complications and hospital length of stay. It was found that patients with sleep apnea had higher rate of postoperative complications and hypoxemia.¹⁰³ We used the STOP-BANG Questionnaire preoperatively to identify patients at high risk for OSAS.¹⁶ We found that patients at high risk of OSAS had a higher rate of postoperative pulmonary and cardiac complications than patients at low risk (19.6% vs 1.3%). We also noted that patients at high risk of OSAS had significantly higher length of stay in the hospital compared to patients at low risk. Gali et al. used Flemons criteria and SACS (sleep apnea clinical score) to identify patients at high or low risk for obstructive sleep apnea.¹⁰⁴ They also noted that patients with high SACS and PACU (postanesthesia care unit) events had a higher rate of postoperative respiratory events. Similarly, Auckley et al. used the Berlin Questionnaire to identify the high-risk patients.¹⁰⁵ They demonstrated that patients at high risk of sleep apnea had more postoperative complications (20% vs 4.5%); however, this difference was not statistically significant.

There are two studies that have been conducted in patients undergoing ambulatory surgery to assess the impact of obstructive sleep apnea.^{106,107} These studies found that the presence of OSA did not increase the rate of unplanned hospital admissions in patients who underwent outpatient surgical procedures. Stierer et al. used a prediction model in a cohort of ambulatory surgical population to assess the probability of sleep apnea.¹⁰⁷ They demonstrated that patients with $\geq 70\%$ propensity for OSA had increased rate of difficult intubation, increased oxygen requirement, and intraoperative tachycardia.

How to Identify Patients with Sleep Apnea

Nocturnal polysomnography (NPSG) is considered to be the gold standard to identify patients with obstructive sleep apnea.¹⁰⁸ However, in the perioperative setting, it is difficult to implement due to a variety of factors, including its prolongation of the process of surgery and contribution to the overall cost. In many hospital settings, it may also not be readily available. There are other methods that have been shown to identify patients who are at risk for obstructive sleep apnea including questionnaires, nocturnal pulse oximetry, and home sleep testing.

Questionnaires

There are many questionnaires that are available to identify surgical patients who are at high risk of having obstructive sleep apnea. Three of these questionnaires have been validated

in the surgical population: the Berlin Questionnaire, ASA checklist, and the STOP-BANG questionnaire. The Berlin questionnaire is the most widely used questionnaire to identify patients at high risk for OSA. It contains 11 questions that are organized in 3 symptom categories and has been validated in the primary care patients.¹⁰⁹ It has a sensitivity of 86% and positive predictive value of 89% for identifying patients with respiratory disturbance index (RDI) $> 5/h$ in the primary care clinic. Recently, Chung et al. validated the Berlin questionnaire in surgical population and found that it had a sensitivity of 74.3% to 79.5% and a negative predictive value of 76% to 89.3% in identifying patients with moderate-to-severe OSA.¹¹⁰ However, this questionnaire has complex scoring system and is time consuming.¹¹¹

The American Society of Anesthesiologists (ASA) checklist also appears to be very useful and promising and has been proposed by ASA Task Force to identify patients with OSA.¹¹² It contains 12 items for adults and 14 items for children. The ASA checklist has been shown to have a sensitivity of 78.6% to 87.2% and a negative predictive value of 72.7% to 90.9% in identifying surgical patients with moderate-to-severe OSA.¹¹⁰

Recently, the STOP-BANG (Snoring, Tiredness during daytime, Observed apnea, high blood Pressure, Body mass index, Age, Neck circumference, Gender) questionnaire was validated as a screening modality for OSAS in the preoperative setting.¹¹³ It is a concise, self-administered, and easy-to-use questionnaire that consists of 8 yes/no questions. Patients are considered to be at high risk of OSAS if they answer yes to ≥ 3 items. The sensitivity of the STOP-BANG questionnaire at an AHI > 5 , > 15 , and > 30 cutoff values was 83.6%, 92.9%, and 100%, respectively; corresponding negative predictive values (NPV) were 60.8%, 90.2%, and 100%. This questionnaire has a moderately high level of sensitivity, specificity, and NPV to detect patients with moderate (AHI > 15) to severe (AHI > 30) sleep apnea in the surgical population. Therefore, if the patient is ranked as a low risk for OSA by the STOP-BANG scoring model, practitioners can exclude the possibility that the patient would have moderate-to-severe sleep apnea with a good degree of accuracy.

Abrishami et al. conducted a systematic review to identify and evaluate the different screening questionnaires for sleep apnea.¹¹⁴ They noted that the Berlin and STOP-BANG questionnaires had high sensitivities and specificities in predicting moderate or severe sleep apnea. However, they found that the STOP and STOP-BANG questionnaires had the highest methodological validity, and these questionnaires are very easy to use.

Nocturnal Pulse Oximetry

Nocturnal pulse oximetry has also been used for the screening of OSA. Malbois et al. compared the sensitivity of nocturnal oximetry to ambulatory monitoring in order to identify patients with sleep apnea prior to undergoing bariatric surgery.¹¹⁵ They demonstrated that nocturnal oximetry with a 3% desaturation index as a screening tool for OSA could rule out significant OSA (AHI > 10) and also detect patients with severe OSA. This cheap and widely available technique could accelerate preoperative work-up of these patients.

Table 3—Perioperative management of patients at high risk of obstructive sleep apnea syndrome**Preoperative Evaluation**

1. History
2. Physical Examination
3. Screening Questionnaires like Berlin, ASA, or STOP BANG to identify high-risk patients
4. Consider a formal sleep evaluation in very high-risk group

Intraoperative Management

1. Minimize the surgical stress
2. Reduce the duration of surgery
3. Consider regional or local anesthesia instead of general anesthesia
4. Anticipate difficult intubation
5. Consider awake extubation preferably in semi-upright position

Postoperative Management

1. Minimize the use of opioids and sedation after the surgery
2. Consider using acetaminophen, NSAIDs, or regional analgesia for the pain control
3. Continuously monitor oxygenation in the postoperative period
4. Patients with a known diagnosis of sleep apnea should use their CPAP after the surgery
5. High-risk patients for sleep apnea should use Auto CPAP during the postoperative period
6. Follow-up at the sleep center for the management of sleep apnea upon discharge from the hospital

Home Sleep Testing

Ambulatory monitoring is another modality that can be employed in patients at high risk for sleep apnea. However, this is recommended only in patients with a high pre-test probability of sleep apnea and is not recommended in patients with coexisting cardiopulmonary complications.¹¹⁶ It also has its limitations in terms of ease of use by the patients. There is one study that has shown the effectiveness of ambulatory monitoring in confirming the diagnosis of OSA in 82 % of adult surgical patients, identified as high risk prior to undergoing surgery, in a large academic medical center.¹⁴ It would be helpful to evaluate this modality of diagnosing OSAS in specific surgical patient population.

Perioperative Management

The American Society of Anesthesiologists published practice guidelines in 2006 on the perioperative management of patients with obstructive sleep apnea.¹¹² It also proposed OSA scoring system to assess the perioperative risk. Based on these guidelines, perioperative care can be subdivided in 3 parts: preoperative evaluation, intraoperative management, and postoperative management (Table 3).

Preoperative Evaluation

Patients should undergo thorough history and physical examination preoperatively with the special emphasis on the evaluation of sleep apnea. One should obtain history pertinent to sleep apnea including snoring, excessive daytime sleepiness, witnessed apneas, frequent awakenings at night, and morning headaches. A focused physical examination should be

conducted to evaluate neck circumference, body mass index, modified Mallampati score, tongue volume, tonsillar size, and nasopharyngeal characteristics. It is important to administer screening questionnaires like the Berlin, ASA, or STOP-BANG to identify patients at high risk for OSA. These questionnaires are simple and easy to administer preoperatively and have been validated in the surgical population. There should be an action plan for the management of high-risk patients during the perioperative period. In few circumstances, anesthesiologists and surgeons may decide to obtain formal sleep evaluation for the management of sleep apnea prior to performing surgery.

Intraoperative Management

Intraoperative management usually focuses on surgical measures and the type of anesthesia. One should minimize the surgical stress and the duration of surgery as these factors have been shown to increase the perioperative complications. Whenever possible, consider using regional or local anesthesia instead of general anesthesia. These patients should be extubated when they are fully awake, preferably in the semi-upright position.

Postoperative Management

Patients at increased perioperative risk from OSA should be very closely monitored in the post anesthesia care unit (PACU) for hypoxemia or other complications. They should have continuous monitoring of oxygenation with the help of pulse oximetry. Whenever possible, these patients should be placed in the non-supine position after the surgery to decrease the severity of apnea. These patients are very susceptible to opioids and benzodiazepines and one should minimize the use of these medicines in the perioperative period. Consider using NSAIDs, acetaminophen, tramadol, and regional analgesia for pain control. Dexmedetomidine can be very useful for sedation because of its opioid sparing effect and the lack of respiratory depression. Patients with the known diagnosis of sleep apnea should use their CPAP after surgery. There is no randomized controlled trial that has demonstrated that CPAP is beneficial in the postoperative setting. However, one may consider using auto-CPAP in high-risk patients after surgery, although it might be difficult for the CPAP-naïve patient to get used to it in the perioperative period. Once again, the use of auto-CPAP has not been formally studied in this population, and there may be a need to conduct a randomized controlled trial to assess the efficacy of auto-CPAP. These patients should also get formal sleep evaluation after discharge from the hospital.

Evidence for Perioperative use of CPAP

CPAP acts as a pneumatic splint and helps in opening the collapsed upper airway at night. The application of CPAP also improves functional residual capacity (FRC) and oxygenation with reduction in work of breathing. CPAP has been shown to improve excessive daytime sleepiness in patients with OSAS. There is some evidence that the perioperative use of CPAP may help in reducing postoperative complications. In a case control study, Gupta et al. noted that OSA patients who were compliant with their CPAP had reduced rate of complications and also decreased hospital length of stay.⁹⁸ Similarly, Liao et al. noted that OSA patients who were not compliant with their CPAP were at the greatest risk of having postoperative complications.⁹⁹

Squadron et al. demonstrated that the use of CPAP leads to reduction in the incidence of endotracheal intubation and other severe complications in patients who develop hypoxemia after elective major abdominal surgery.¹¹⁷ In a randomized controlled trial, Kingen-Milles et al. found that the prophylactic use of nasal CPAP was associated with a reduction in pulmonary complications and hospital length of stay in patients undergoing thoracoabdominal aortic aneurysm repair.¹¹⁸ In another study, Zarbock et al. also noted significant reduction in the rate of pulmonary complications with the prophylactic use of nasal CPAP in patients undergoing elective cardiac surgery.¹¹⁹ A recent meta-analysis of nine randomized controlled trials in the abdominal surgical population reported reduction in the rate of atelectasis, postoperative pulmonary complications, and pneumonia with the perioperative use of CPAP.¹²⁰

Conclusions

Obstructive sleep apnea syndrome (OSAS) is a common type of sleep disordered breathing, with a high prevalence in the surgical population. The majority of patients with sleep apnea are undiagnosed and are therefore unaware of their OSAS at the time of the surgery. These patients are at increased risk for perioperative complications. Sedation, anesthesia, opioids, and REM sleep rebound have been shown to cause worsening of sleep apnea in the perioperative period that may lead to increase in the rate of perioperative complications. It is important to identify these patients preoperatively so that appropriate actions can be taken during their perioperative care. Screening questionnaires such as the Berlin, STOP-BANG, or ASA checklist are easy to administer preoperatively and have been shown to identify high-risk patients. There should be a standard protocol for the perioperative management of high-risk patients in order to reduce the rate of complications. These high-risk patients should also have a formal sleep evaluation for the long-term management of their sleep apnea after the discharge from the hospital.

REFERENCES

- Howard ME, Desai AV, Grunstein RR, et al. Sleepiness, sleep-disordered breathing, and accident risk factors in commercial vehicle drivers. *Am J Respir Crit Care Med* 2004;170:1014-21.
- Tregear S, Reston J, Schoelles K, Phillips B. Obstructive sleep apnea and risk of motor vehicle crash: systematic review and meta-analysis. *J Clin Sleep Med* 2009;5:573-81.
- Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000;342:1378-84.
- Tasali E, Mokhlesi B, Van Cauter E. Obstructive sleep apnea and type 2 diabetes: interacting epidemics. *Chest* 2008;133:496-506.
- Sharma B, Owens R, Malhotra A. Sleep in congestive heart failure. *Med Clin North Am* 2010;94:447-64.
- Yaggi HK, Concato J, Kernan WN, et al. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med* 2005;353:2034-41.
- Marin JM, Carrizo SJ, Vicente E, et al. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet* 2005;365:1046-53.
- Punjabi NM, Caffo BS, Goodwin JL, et al. Sleep-disordered breathing and mortality: a prospective cohort study. *PLoS Med* 2009;6:e1000132.
- Marshall NS, Wong KK, Liu P, et al. Sleep apnea as an independent risk factor for all-cause mortality: the Busselton Health Study. *Sleep* 2008;31:1079-85.
- Young T, Palta M, Dempsey J, et al. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 1993;328:1230-5.
- Hiestand DM, Britz P, Goldman M, et al. Prevalence of symptoms and risk of sleep apnea in the US population: Results from the national sleep foundation sleep in America 2005 poll. *Chest* 2006;130:780-6.
- Frey WC, Pilcher J. Obstructive sleep-related breathing disorders in patients evaluated for bariatric surgery. *Obes Surg* 2003;13:676-83.
- O'Keeffe T, Patterson EJ. Evidence supporting routine polysomnography before bariatric surgery. *Obes Surg* 2004;14:23-6.
- Finkel KJ, Searleman AC, Tymkew H, et al. Prevalence of undiagnosed obstructive sleep apnea among adult surgical patients in an academic medical center. *Sleep Med* 2009;10:753-8.
- Chung F, Ward B, Ho J, Yuan H, et al. Preoperative identification of sleep apnea risk in elective surgical patients, using the Berlin questionnaire. *J Clin Anesth* 2007;19:130-4.
- Vasu TS, Doghramji K, Cavallazzi R, et al. Obstructive sleep apnea syndrome and postoperative complications: clinical use of the STOP-BANG questionnaire. *Arch Otolaryngol Head Neck Surg* 2010;136:1020-4.
- Malow BA, Levy K, Maturen K, Bowes R. Obstructive sleep apnea is common in medically refractory epilepsy patients. *Neurology* 2000;55:1002-7.
- Pollak L, Shpirer I, Rabey JM, et al. Polysomnography in patients with intracranial tumors before and after operation. *Acta Neurol Scand* 2004;109:56-60.
- Gold AR, Schwartz AR. The pharyngeal critical pressure. The whys and hows of using nasal continuous positive airway pressure diagnostically. *Chest* 1996;110:1077-88.
- Gold AR, Marcus CL, Dipalo F, et al. Upper airway collapsibility during sleep in upper airway resistance syndrome. *Chest* 2002;121:1531-40.
- Drummond GB. Comparison of sedation with midazolam and ketamine: effects on airway muscle activity. *Br J Anaesth* 1996;76:663-7.
- Drummond GB. Influence of thiopentone on upper airway muscles. *Br J Anaesth* 1989;63:12-21.
- Eastwood PR, Platt PR, Shepherd K, et al. Collapsibility of the upper airway at different concentrations of propofol anesthesia. *Anesthesiology* 2005;103:470-7.
- Hillman DR, Walsh JH, Maddison KJ, et al. Evolution of changes in upper airway collapsibility during slow induction of anesthesia with propofol. *Anesthesiology* 2009;111:63-71.
- Knill RL, Clement JL. Site of selective action of halothane on the peripheral chemoreflex pathway in humans. *Anesthesiology* 1984;61:121-6.
- Dahan A, van den Elsen MJ, Berkenbosch A, et al. Effects of subanesthetic halothane on the ventilatory responses to hypercapnia and acute hypoxia in healthy volunteers. *Anesthesiology* 1994;80:727-38.
- van den Elsen M, Dahan A, DeGoede J, et al. Influences of subanesthetic isoflurane on ventilatory control in humans. *Anesthesiology* 1995;83:478-90.
- Knill RL, Kierszewicz HT, Dodgson BG, et al. Chemical regulation of ventilation during isoflurane sedation and anaesthesia in humans. *Can Anaesth Soc J* 1983;30:607-14.
- Borison HL. Central nervous respiratory depressants--narcotic analgesics. *Pharmacol Ther B* 1977;3:227-37.
- Berkenbosch A, Teppema LJ, Olivier CN, Dahan A. Influences of morphine on the ventilatory response to isocapnic hypoxia. *Anesthesiology* 1997;86:1342-9.
- Christensen V. Respiratory depression after extradural morphine. *Br J Anaesth* 1980;52:841.
- Reiz S, Westberg M. Side-effects of epidural morphine. *Lancet* 1980;2:203-4.
- Knill RL, Lam AM, Thompson WR. Epidural morphine and ventilatory depression. *Anesthesiology* 1982;56:486-8.
- Dahan A, Sarton E, Teppema L, et al. Sex-related differences in the influence of morphine on ventilatory control in humans. *Anesthesiology* 1998;88:903-13.
- Sarton E, Teppema L, Dahan A. Sex differences in morphine-induced ventilatory depression reside within the peripheral chemoreflex loop. *Anesthesiology* 1999;90:1329-38.
- Bailey PL, Pace NL, Ashburn MA, et al. Frequent hypoxemia and apnea after sedation with midazolam and fentanyl. *Anesthesiology* 1990;73:826-30.
- Weil JV, McCullough RE, Kline JS, et al. Diminished ventilatory response to hypoxia and hypercapnia after morphine in normal man. *N Engl J Med* 1975;292:1103-6.
- Bailey PL, Lu JK, Pace NL, et al. Effects of intrathecal morphine on the ventilatory response to hypoxia. *N Engl J Med* 2000;343:1228-34.
- Alexander CM, Gross JB. Sedative doses of midazolam depress hypoxic ventilatory responses in humans. *Anesth Analg* 1988;67:377-82.
- Ellis BW, Dudley HA. Some aspects of sleep research in surgical stress. *J Psychosom Res* 1976;20:303-8.
- Kavey NB, Ahshuler KZ. Sleep in herniorrhaphy patients. *Am J Surg* 1979;138:683-7.
- Lehmkuhl P, Prass D, Pichlmayr I. General anesthesia and postnarcotic sleep disorders. *Neuropsychobiology* 1987;18:37-42.

43. Aurell J, Elmqvist D. Sleep in the surgical intensive care unit: continuous polygraphic recording of sleep in nine patients receiving postoperative care. *Br Med J (Clin Res Ed)* 1985;290:1029-32.
44. Knill RL, Moote CA, Skinner MI, et al. Anesthesia with abdominal surgery leads to intense REM sleep during the first postoperative week. *Anesthesiology* 1990;73:52-61.
45. Rosenberg J. Sleep disturbances after non-cardiac surgery. *Sleep Med Rev* 2001;5:129-37.
46. Rosenberg-Adamsen S, Skarbye M, Wildschiodtz G, et al. Sleep after laparoscopic cholecystectomy. *Br J Anaesth* 1996;77:572-5.
47. Helton MC, Gordon SH, Nunnery SL. The correlation between sleep deprivation and the intensive care unit syndrome. *Heart Lung* 1980;9:464-8.
48. de Andres I, Corpas I. Morphine effects in brainstem-transected cats: II. Behavior and sleep of the decerebrate cat. *Behav Brain Res* 1991;44:21-6.
49. Fehm HL, Benkowitz R, Kern W, et al. Influences of corticosteroids, dexamethasone and hydrocortisone on sleep in humans. *Neuropsychobiology* 1986;16:198-204.
50. Moser NJ, Phillips BA, Guthrie G, et al. Effects of dexamethasone on sleep. *Pharmacol Toxicol* 1996;79:100-2.
51. Lin E, Calvano SE, Lowry SF. Inflammatory cytokines and cell response in surgery. *Surgery* 2000;127:117-26.
52. Menger MD, Vollmar B. Surgical trauma: hyperinflammation versus immunosuppression? *Langenbecks Arch Surg* 2004;389:475-84.
53. Ni Chioileain N, Redmond HP. Cell response to surgery. *Arch Surg* 2006;141:1132-40.
54. Opp MR, Kapas L, Toth LA. Cytokine involvement in the regulation of sleep. *Proc Soc Exp Biol Med* 1992;201:16-27.
55. Kapas L, Hong L, Cady AB, et al. Somnogenic, pyrogenic, and anorectic activities of tumor necrosis factor-alpha and TNF-alpha fragments. *Am J Physiol* 1992;263(3 Pt 2):R708-15.
56. Krueger JM, Fang J, Taishi P, et al. Sleep. A physiologic role for IL-1 beta and TNF-alpha. *Ann N Y Acad Sci* 1998;856:148-59.
57. Knill RL, Moote CA, Skinner MI, et al. Anesthesia with abdominal surgery leads to intense REM sleep during the first postoperative week. *Anesthesiology* 1990;73:52-61.
58. Rosenberg JF, Ullstad TF, Rasmussen J, et al. Time course of postoperative hypoxaemia. *Eur J Surg* 1994;160:137-43.
59. Rosenberg JF, Wildschiodtz G, Pedersen MH, et al. Late postoperative nocturnal episodic hypoxaemia and associated sleep pattern. *Br J Anaesth* 1994;72:145-50.
60. Knill RL, Skinner MI, Novick T, et al. The night of intense REM sleep after anesthesia and surgery increases urinary catecholamines. *Can J Anaesth* 1990;37(4 Pt 2):S12.
61. Reeder MK, Muir AD, Foex P, et al. Postoperative myocardial ischaemia: temporal association with nocturnal hypoxaemia. *Br J Anaesth* 1991;67:626-31.
62. Somers VK, Dyken ME, Mark AL, et al. Sympathetic-nerve activity during sleep in normal subjects. *N Engl J Med* 1993;328:303-7.
63. Mancia G. Autonomic modulation of the cardiovascular system during sleep. *N Engl J Med* 1993;328:347-9.
64. Gogenur I, Rosenberg-Adamsen S, Lie C, et al. Relationship between nocturnal hypoxaemia, tachycardia and myocardial ischaemia after major abdominal surgery. *Br J Anaesth* 2004;93:333-8.
65. Galatius-Jensen S, Hansen J, Rasmussen V, et al. Nocturnal hypoxaemia after myocardial infarction: association with nocturnal myocardial ischaemia and arrhythmias. *Br Heart J* 1994;72:23-30.
66. Kehlet H, Rosenberg J. Late post-operative hypoxaemia and organ dysfunction. *Eur J Anaesthesiol Suppl* 1995;10:31-4.
67. Tarhan S, Moffitt EA, Taylor WF, et al. Myocardial infarction after general anesthesia. *JAMA* 1972;220:1451-4.
68. Edwards H, Rose EA, Schorow M, et al. Postoperative deterioration in psychomotor function. *JAMA* 1981;245:1342-3.
69. Brimacombe J, Macfie AG. Peri-operative nightmares in surgical patients. *Anaesthesia* 1993;48:527-9.
70. Galanakis P, Bickel H, Gradinger R, et al. Acute confusional state in the elderly following hip surgery: incidence, risk factors and complications. *Int J Geriatr Psychiatry* 2001;16:349-55.
71. Oxford Center for Evidence-Based Medicine. Levels of Evidence Working Group. The Oxford 2011 Levels of Evidence. Oxford Centre for Evidence-Based Medicine. <http://www.cebm.net/index.aspx?o=5653>.
72. Pedersen T. Complications and death following anaesthesia. A prospective study with special reference to the influence of patient-, anaesthesia-, and surgery-related risk factors. *Dan Med Bull* 1994;41:319-31.
73. Schwilk B, Muche R, Treiber H, et al. A cross-validated multifactorial index of perioperative risks in adults undergoing anaesthesia for non-cardiac surgery. Analysis of perioperative events in 26907 anaesthetic procedures. *J Clin Monit Comput* 1998;14:283-94.
74. Ballotta E, Da Giau G, Militello C, et al. Major elective surgery for vascular disease in patients aged 80 or more: perioperative (30-day) outcomes. *Ann Vasc Surg* 2007;21:772-9.
75. Basilico FC, Sweeney G, Losina E, et al. Risk factors for cardiovascular complications following total joint replacement surgery. *Arthritis Rheum* 2008;58:1915-20.
76. McAlister FA, Bertsch K, Man J, et al. Incidence of and risk factors for pulmonary complications after nonthoracic surgery. *Am J Respir Crit Care Med* 2005;171:514-7.
77. Leung JM, Dzankic S. Relative importance of preoperative health status versus intraoperative factors in predicting postoperative adverse outcomes in geriatric surgical patients. *J Am Geriatr Soc* 2001;49:1080-5.
78. Tiet L, Hatton F, Desmonts JM, et al. Prediction of outcome of anaesthesia in patients over 40 years: a multifactorial risk index. *Stat Med* 1988;7:947-54.
79. Scholes RL, Browning L, Sztendur EM, et al. Duration of anaesthesia, type of surgery, respiratory co-morbidity, predicted VO2max and smoking predict postoperative pulmonary complications after upper abdominal surgery: an observational study. *Aust J Physiother* 2009;55:191-8.
80. Pereira ED, Fernandes AL, da Silva Ancao M, et al. Prospective assessment of the risk of postoperative pulmonary complications in patients submitted to upper abdominal surgery. *Sao Paulo Med J* 1999;117:151-60.
81. Khuri SF, Daley J, Henderson W, et al. The National Veterans Administration Surgical Risk Study: risk adjustment for the comparative assessment of the quality of surgical care. *J Am Coll Surg* 1995;180:519-31.
82. Gibbs J, Cull W, Henderson W, et al. Preoperative serum albumin level as a predictor of operative mortality and morbidity: results from the National VA Surgical Risk Study. *Arch Surg* 1999;134:36-42.
83. Arozullah AM, Daley J, Henderson WG, et al. Multifactorial risk index for predicting postoperative respiratory failure in men after major noncardiac surgery. The National Veterans Administration Surgical Quality Improvement Program. *Ann Surg* 2000;232:242-53.
84. Mitchell CK, Smoger SH, Pfeifer MP, et al. Multivariate analysis of factors associated with postoperative pulmonary complications following general elective surgery. *Arch Surg* 1998;133:194-8.
85. Mircea N, Constantinescu C, Jianu E, et al. Risk of pulmonary complications in surgical patients. *Resuscitation* 1982;10:33-41.
86. Calligaro KD, Azurin DJ, Dougherty MJ, et al. Pulmonary risk factors of elective abdominal aortic surgery. *J Vasc Surg* 1993;18:914-20; discussion 920-1.
87. Parker MJ, Handoll HH, Griffiths R. Anaesthesia for hip fracture surgery in adults. *Cochrane Database Syst Rev* 2004;18:CD000521.
88. Hallan S, Molaug PO, Arnulf V, et al. Causes and risk factors of intraoperative anesthesia complications. A prospective study of 14,735 anesthetics. *Tidsskr Nor Laegeforen* 1990;110:38-41.
89. Pedersen T, Eliassen K, Henriksen E. A prospective study of mortality associated with anaesthesia and surgery: risk indicators of mortality in hospital. *Acta Anaesthesiol Scand* 1990;34:176-82.
90. Kroenke K, Lawrence VA, Theroux JF, et al. Postoperative complications after thoracic and major abdominal surgery in patients with and without obstructive lung disease. *Chest* 1993;104:1445-51.
91. Kanat F, Golcuk A, Teke T, et al. Risk factors for postoperative pulmonary complications in upper abdominal surgery. *ANZ J Surg* 2007;77:135-41.
92. Wong DH, Weber EC, Schell MJ, et al. Factors associated with postoperative pulmonary complications in patients with severe chronic obstructive pulmonary disease. *Anesth Analg* 1995;80:276-84.
93. Vodinh J, Bonnet F, Touboul C, et al. Risk factors of postoperative pulmonary complications after vascular surgery. *Surgery* 1989;105:360-5.
94. Svensson LG, Hess KR, Coselli JS, et al. A prospective study of respiratory failure after high-risk surgery on the thoracoabdominal aorta. *J Vasc Surg* 1991;14:271-82.
95. Money SR, Rice K, Crockett D, et al. Risk of respiratory failure after repair of thoracoabdominal aortic aneurysms. *Am J Surg* 1994;168:152-5.
96. Arozullah AM, Khuri SF, Henderson WG, Daley J; Participants in the National Veterans Affairs Surgical Quality Improvement Program. Development and validation of a multifactorial risk index for predicting postoperative pneumonia after major noncardiac surgery. *Ann Intern Med* 2001;135:847-57.
97. McCulloch TM, Jensen NF, Girod DA, et al. Risk factors for pulmonary complications in the postoperative head and neck surgery patient. *Head Neck* 1997;19:372-7.

98. Gupta RM, Parvizi J, Hanssen AD, et al. Postoperative complications in patients with obstructive sleep apnea syndrome undergoing hip or knee replacement: a case-control study. *Mayo Clin Proc* 2001;76:897-905.
99. Liao P, Yegneswaran B, Vairavanathan S, et al. Postoperative complications in patients with obstructive sleep apnea: a retrospective matched cohort study. *Can J Anaesth* 2009;56:819-28.
100. Kaw R, Golish J, Ghamande S, et al. Incremental risk of obstructive sleep apnea on cardiac surgical outcomes. *J Cardiovasc Surg (Torino)* 2006;47:683-9.
101. Memsoudis S, Liu SS, Ma Y, et al. Perioperative pulmonary outcomes in patients with sleep apnea after noncardiac surgery. *Anesth Analg* 2011;112:113-21.
102. Hwang D, Shakir N, Limann B, et al. Association of sleep-disordered breathing with postoperative complications. *Chest* 2008;133:1128-34.
103. Kaw R, Pasupuleti V, Walker E, Ramaswamy A, Foldvary-Schafer N. Postoperative complications in patients with obstructive sleep apnea. *Chest* 2011 Aug 25 (Epub ahead of print).
104. Gali B, Whalen FX, Schroeder DR, et al. Identification of patients at risk for postoperative respiratory complications using a preoperative obstructive sleep apnea screening tool and postanesthesia care assessment. *Anesthesiology* 2009;110:869-77.
105. Auckley DH, Steinel J, Southwell C, et al. Does screening for sleep apnea with the berlin questionnaire predict elective surgery postoperative complications? *Sleep (Abstract Supplement)* 2003;26:A238-9.
106. Sabers C, Plevak DJ, Schroeder DR, et al. The diagnosis of obstructive sleep apnea as a risk factor for unanticipated admissions in outpatient surgery. *Anesth Analg* 2003;96:1328-35.
107. Stierer TL, Wright C, George A, et al. Risk assessment of obstructive sleep apnea in a population of patients undergoing ambulatory surgery. *J Clin Sleep Med* 2010;6:467-72.
108. Kushida CA, Littner MR, Morgenthaler T, et al. Practice parameters for the indications for polysomnography and related procedures: an update for 2005. *Sleep* 2005;28:499-521.
109. Netzer NC, Stoohs RA, Netzer CM, et al. Using the Berlin Questionnaire to identify patients at risk for the sleep apnea syndrome. *Ann Intern Med* 1999;131:485-91.
110. Chung F, Yegneswaran B, Liao P, et al. Validation of the Berlin questionnaire and American Society of Anesthesiologists checklist as screening tools for obstructive sleep apnea in surgical patients. *Anesthesiology* 2008;108:822-30.
111. Senthilvel E, Auckley D, Dasarathy J. Evaluation of sleep disorders in the primary care setting: history taking compared to questionnaires. *J Clin Sleep Med* 2011;7:4148.
112. Gross JB, Bachenberg KL, Benumof JL, et al. Practice guidelines for the perioperative management of patients with obstructive sleep apnea: a report by the American Society of Anesthesiologists Task Force on Perioperative Management of patients with obstructive sleep apnea. *Anesthesiology* 2006;104:1081-93; quiz 1117-8.
113. Chung F, Yegneswaran B, Liao P, et al. STOP questionnaire: a tool to screen patients for obstructive sleep apnea. *Anesthesiology* 2008;108:812-21.
114. Abrishami A, Khajehdehi A, Chung F. A systematic review of screening questionnaires for obstructive sleep apnea. *Can J Anaesth* 2010; 57:423-38.
115. Malbois M, Giusti V, Suter M, et al. Oximetry alone versus portable polygraphy for sleep apnea screening before bariatric surgery. *Obes Surg* 2010; 20:326-31.
116. Collop NA, Anderson WM, Boehlecke B, et al. Clinical guidelines for the use of unattended portable monitors in the diagnosis of obstructive sleep apnea in adult patients. Portable Monitoring Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2007;3:737-47.
117. Squadrone V, Coxa M, Cerutti E, et al. Continuous positive airway pressure for treatment of postoperative hypoxemia: a randomized controlled trial. *JAMA* 2005;293:589-95.
118. Kindgen-Milles D, Muller E, Buhl R, et al. Nasal-continuous positive airway pressure reduces pulmonary morbidity and length of hospital stay following thoracoabdominal aortic surgery. *Chest* 2005;128:821-8.
119. Zarbock A, Mueller E, Netzer S, et al. Prophylactic nasal continuous positive airway pressure following cardiac surgery protects from postoperative pulmonary complications: a prospective, randomized, controlled trial in 500 patients. *Chest* 2009;135:1252-9.
120. Ferreyra GP, Baussano I, Squadrone V, et al. Continuous positive airway pressure for treatment of respiratory complications after abdominal surgery: a systematic review and meta-analysis. *Ann Surg* 2008;247:617-26.

SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication October, 2011

Submitted in final revised form December, 2011

Accepted for publication January, 2012

Address correspondence to: Tajender S Vasu, M.D., M.S., Division of Pulmonary, Critical Care, and Sleep Medicine, Assistant Professor of Medicine, Stony Brook University Medical Center, HSC T 17-040, Stony Brook, NY 11794; Tel: (631) 444-1707; Fax: (631) 444-7502; E mail: Tajender.Vasu@sbumed.org

DISCLOSURE STATEMENT

This was not an industry supported study. The authors have indicated no financial conflicts of interest.