Published in final edited form as:

J Adolesc Health. 2012 April ; 50(4): 358–364. doi:10.1016/j.jadohealth.2011.08.009.

A Family Intervention to Reduce Sexual Risk Behavior, Substance Use, and Delinquency Among Newly Homeless Youth

Norweeta G. Milburn, PhD, Francisco Javier Iribarren, MSW-PsyD, Eric Rice, PhD, Marguerita Lightfoot, PhD, Rosa Solorio, MD, Mary Jane Rotheram-Borus, PhD, Katherine Desmond, MS, Alex Lee, BA, Kwame Alexander, BA, Katherine Maresca, BA, Karen Eastmen, PhD, Elizabeth Mayfield Arnold, PhD, and Naihua Duan, PhD

Department of Psychiatry and Biobehavioral Sciences, Semel Institute for Neuroscience and Human Behavior, Center for Community Health, University of California, Los Angeles (Drs. Milburn, Rotheram-Borus, and Eastmen; Ms. Desmond and Ms. Maresca; Mr. Lee and Mr. Alexander); Chicano Studies Research Center, University of California, Los Angeles (Dr. Iribarren); School of Social Work, University of Southern California, Los Angeles (Dr. Rice); School of Public Health, University of Washington (Dr. Solorio); Center for AIDS Prevention Studies, University of California, San Francisco (Dr. Lightfoot); Wake Forest University School of Medicine (Dr. Arnold); and Columbia University (Dr. Duan)

Abstract

Purpose—We evaluate the efficacy of a short family intervention in reducing sexual risk behavior, drug use, and delinquent behaviors among homeless youth.

Methods—A randomized controlled trial of 151 families with a homeless adolescent aged 12 to 17 years. Adolescents were recruited from diverse sites in southern California from March 2006 through June 2009 and assessed at recruitment (baseline), 3, 6, and 12 months later. Families were randomly assigned to an intervention condition with five weekly home-based intervention sessions or a control condition (standard care). Main outcome measures reflect self-reported sexual, substance use and delinquent behaviors over the last 90 days.

Results—Sexual risk (e.g., mean number of partners) ($p < .001$), alcohol use ($p = .003$), hard drug use ($p < .001$), and delinquent behaviors ($p = .001$) decreased significantly more over 12 months in the intervention condition compared to the control condition. Marijuana use, however, significantly increased in the intervention condition compared to the control condition ($p < .001$).

Conclusion—An intervention to re-engage families of homeless youth has significant benefits in reducing risk over 12 months.

© 2011 Society for Adolescent Medicine. Published by Elsevier Inc. All rights reserved.

Corresponding Author: Norweeta G. Milburn, PhD, Center for Community Health, Semel Institute for Neuroscience and Human Behavior, 10920 Wilshire Blvd., Suite 350, Los Angeles, CA 90024. Telephone: 310-794-3773; Facsimile: 310-794-8297; nmilburn@mednet.ucla.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Implications and Contribution: Homeless youth engage in risky acts that increase substantially with chronic homelessness. Typically, street-based interventions with chronic homeless youth have been the focus of interventions. In this article, we present findings that demonstrate successful efficacy of an intervention on sexual and drug-taking HIV risk behaviors among newly homeless youth.

Keywords

delinquency; HIV; homelessness; intervention

INTRODUCTION

There are an estimated 1.6 million runaway and homeless youth in the United States each year (1). Although recent reports indicate that the prevalence of HIV among the general population of young people in the United States is only 0.2% (2), the rate of infection among homeless youth far exceeds this number in urban centers, with estimates ranging from 2% to 11% (3–4). Research has consistently shown that the longer youth are away from home and living on the streets, the more heavily involved they become in risky behavior, especially HIV-related sexual and drug-taking risk (5–12). Intervening when homeless youth first leave home may provide the best opportunity for reducing their risk behaviors.

Recent work has demonstrated that most runaways return home within 3 months of first becoming homeless (13). Contact with parents, especially mothers, increases the chances that newly homeless youth will return home and remain at home over time (14). To date, however, the only effective HIV prevention programs for homeless youth have targeted individual youth in the context of street life (15). Interventions involving the family that occur before the youth becomes chronically homeless may prove an efficacious alternative (16). Behavioral family interventions can be effective in preventing negative developmental outcomes for children and adolescents such as delinquent behaviors (17). Family interventions have been successfully employed with other high-risk youth, including those who are suicidal (18), delinquent (19), or engaged in HIV-related risk behaviors such as substance abuse (17, 20–23).

To address the need for family-based risk reduction and HIV prevention for homeless youth, we evaluated the efficacy of a brief family intervention in reducing risk-taking and delinquent behaviors among newly homeless youth in Southern California. We hypothesized that risk behaviors are prevalent among homeless youth, but that a family intervention would result in less risk-taking and fewer delinquent behaviors over time among those in the intervention compared to the control condition.

METHODS

Study Design and Participants

A randomized controlled trial was conducted from March 2006 through June 2009. Newly homeless youth were recruited from community-based organizations (e.g., presentations at shelters or schools) and from direct recruitment (e.g., flyers, advertisements) in Los Angeles and San Bernardino counties. Eligibility criteria included having been away from home for at least two nights in the past six months, not being away for more than six months, and having the potential to return home. Additional eligibility criteria included no current abuse or neglect, no active psychosis, or no current substance intoxication; these additional criteria could not be determined prior to obtaining informed consent and were screened for in the baseline assessment.

Flyers and recruitment scripts indicated eligibility criteria for the study. Adolescents expressed interest in the study by calling phone numbers provided on recruitment materials or by approaching recruiters. Attempts were then made to obtain written informed consent from both the adolescent and parent to participate.* The Institutional Review Board of the University of California, Los Angeles, approved the protocol for this study.

After both the adolescent and parent assented/consented, they completed baseline assessments. Eligible families were then randomly assigned to one of two treatment conditions, intervention or control. Families in the control condition received the “standard care” that they were receiving from the agencies that referred them. If they were not actively receiving any type of services, the families were given appropriate referrals based on their needs. Families in the intervention condition received the STRIVE (Support To Reunite, Involve and Value Each Other) intervention.

Separate teams were involved in the recruitment/assessment and intervention aspects of the study. Once a family provided consent and both baseline assessments, the recruitment/assessment team referred the family to the intervention team, which used a computerized coin toss to assign families to study condition, and provided the intervention to families randomized to receive it. The recruitment/assessment team continued to follow all families through June 2009, blinded as to study arm.

Following outreach efforts, 442 adolescents initially indicated interest in the study. Among these, 151 provided full consent from parent and child, completed both the parent and adolescent baseline assessments, and were randomized. Sixty-eight were assigned to receive the intervention; 83 were in the control condition. Parents included mothers/female guardians (76%), fathers/male guardians (10%), grandparents (6%) and other adults (7%).

Intervention Methods

The STRIVE intervention consisted of five sessions administered to the youth and parent(s) together by a trained facilitator at a site selected by the family (usually their home, where youth could be found despite runaway episodes). Five sessions were determined as the minimum number of sessions required for behavioral change and the maximum number of sessions anticipated to be replicable (24, 25). The session content was based upon cognitive-behavioral theories, designed to improve families’ problem-solving and conflict resolution skills (17, 26–28). Since the intervention was specifically designed for newly homeless youth, running away from home was framed as an ineffective attempt to resolve family conflicts. The conceptual frame underlying the intervention highlights the importance of establishing a positive family climate, improving family functioning by routinely resolving conflicts in a mutually beneficial manner, learning how to recognize and effectively manage feelings, increasing positive affirmations, learning and practicing problem solving skills, and providing role clarity (29–32). As can be seen in Table 1, the sessions are based on a set of very highly interactive, semi-structured tasks involving repeated feedback and practice. New skills are learned based on the preceding sessions, and the skills learned in one session are constantly reinforced in the subsequent sessions. For example, nurturing a positive family climate is reinforced in all five sessions. The intervention includes the following tools: tokens to strengthen desired behaviors, a feeling thermometer to teach emotional regulation, a “think-feel-do” problem solving model to operationalize and tackle problems, role playing for safely practicing new skills, and reframing to conceptualize problems and solutions in a non-blaming manner.

An intervention manual was created to ensure fidelity. Intervention sessions usually lasted between 1½ and 2 hours, and were conducted once weekly. All sessions were audio-recorded and 20% were monitored for quality assurance. A 13-item rating instrument was developed to assess fidelity to session content and goals; overall 98% of tasks were implemented with fidelity. Attendance was high: 76% of families completed all 5 sessions;

*Parent refers to parent or guardian including a birth, adoptive, step or foster parent. In two parent families, both parents could participate in the intervention sessions if the family wanted both parents to participate but data were collected from only one parent that the family selected as the parent participant.

6% attended 3–4 sessions; 16% attended 1–2 sessions; and only one family did not attend any sessions. In addition, satisfaction was assessed using the Working Alliance Inventory (WAI). The WAI is a valid and reliable instrument with scores ranging from 12 to 84 (33). The median adolescent score was 78.0 (SD = 12.5, mean = 72.0) and the median parent was 81.0 (SD = 10.0, mean = 76.4), indicating that satisfaction was very high.

Assessments

A highly trained assessment team diverse in ethnicity and gender conducted computerized interviews. Audio computer-assisted self-interviewing (ACASI) was used for sensitive measures. Youth in the study completed assessments at baseline, and 3, 6 and 12 months after the baseline. All youth were paid for completing each assessment (e.g., \$30, \$35, and \$40). Of the 151 randomized adolescents, 77% (116) completed at least one follow-up; 71% (107) completed a 3-month; 58% (87) completed a 6-month; and 46% (69) completed a 12-month assessment. On average, the 3-month assessment occurred 16 days after the last intervention session, but 21 cases had the 3-month assessment before the last intervention session.

Demographic data (age, gender, sexual orientation, race/ethnicity, and birthplace) were collected at baseline, as well as information about running away. Lifetime history of sexual- and drug-related HIV risk behaviors was also assessed at baseline, as was history of delinquent behavior.

Measures of recent risky behavior were collected at each assessment. “Recent” behaviors were defined as in the past 3 months. Sex was defined as vaginal or anal intercourse and measures of sexual behavior included: a dummy indicator for having sex, an indicator for having unprotected sex (without using a condom), number of sexual partners, and number of times had sex. Measures of recent risky behavior also included substance use variables, again reflecting behavior in the 3 months prior to each assessment (indicators for using alcohol, marijuana, and hard drugs; number of times used alcohol, marijuana, and hard drugs).

Delinquent behaviors were measured by a count of conduct problems engaged in during the prior 3 months, from a list of 13 possible behaviors, identified in the DSM-IV as being symptomatic of conduct disorder (34). The list of behaviors is provided in Table 2; examples include starting fights or destroying property. Runaway behavior was not included, as it was an inclusion criterion for the study.

Statistical Methods

Baseline characteristics of participants assigned to the intervention were compared to those of participants assigned to the control condition. Measures compared included demographics, runaway history, and baseline values of outcome measures. Those completing at least one follow-up assessment were compared to those lost to follow-up. For both sets of comparisons, t-tests and chi square tests were performed for continuous and categorical measures, respectively.

The impact of the STRIVE intervention on adolescents’ risk-taking and delinquent behaviors was evaluated by intent-to-treat random-intercept regression models. Outcome variables were the measures of recent risky behavior and delinquent behaviors. In these models, a random intercept was included for each youth, to account for repeated measures taken from the same individual. Model predictors include the intercept, gender, a time measure for each assessment, an intervention status indicator, and an interaction between time and intervention status. The interaction term captures the impact of the intervention over time (i.e., the relative change across assessments for those in the intervention condition

as compared to control condition). To test for non-linear time trends in the outcome measures, the models were re-estimated with the addition of squared terms for the time variables. If the squared terms were significant, they were retained in the model; if non-significant, the more parsimonious model was used. We present results from significance tests of the time-intervention interaction terms, as well as observed and model-fitted mean values at each time point for youth in both arms of the study.

RESULTS

Baseline

Characteristics of randomized participants ($N = 151$) are shown in Table 2, overall and by intervention status. Adolescents were similar on all baseline characteristics except gender; the intervention condition had significantly more females (78%) compared to the control condition (57%) ($p = .006$). Thus, gender was controlled for in all analyses of the intervention effect. The average age of youth enrolled in the study was 15 years (range 12–17) and 90% were heterosexual. Almost all were U.S.-born (92%). Youth were predominantly Hispanic (62%) and African-American (21%), with fewer Whites (11%) and other race/ethnicities (7%). Few had ever run away for more than a month (13% away for 2–6 months), and the majority (63%) had never been away for more than two weeks. Over 70% were back with their families by the time the baseline assessment was administered.

Effects of the Intervention

For these analyses, we examined baseline and follow-up assessments administered 3, 6, and 12 months after the baseline. Baseline characteristics of those who did and did not complete any follow-ups were compared; no statistically significant differences were found.

We expected the newly homeless youth to be more engaged in risk behaviors than youth in general. Table 3 shows the frequencies of substance use and sexual risk behaviors among a nationally representative sample of high school students (35), compared to baseline estimates from the STRIVE study. Alcohol and inhalant use among STRIVE participants were comparable to national norms (72% vs. 73%, 11% vs. 12%, respectively; NS). Use of other substances (marijuana: 59% vs. 37%; cocaine: 17% vs. 6%; methamphetamines: 13% vs. 4%) was substantially greater among those in the STRIVE study ($p < .05$). STRIVE youth were not significantly more likely to have had sex (53% vs. 46%; NS) but were less likely to have used a condom if they did (45% vs. 61%; $p < .05$). Estimated use of alcohol or drugs prior to sex was lower among the STRIVE participants (14% vs. 22%), but the difference was not significant.

Intent-to-treat analyses were conducted to estimate the impact of the STRIVE intervention on risk behaviors in the 3 months prior to each assessment. With regard to sexual risk-taking, a significant effect was found for number of partners. No significant intervention effect was found for whether or not the adolescent had been sexually active, whether or not the adolescent had unprotected sex, or for the number of times the adolescent had sex. With regard to substance use, a significant effect was found for number of times alcohol, marijuana, and hard drugs were used. No significant intervention effect was found for whether the adolescent used these substances. A significant difference was found between intervention and control participants in number of delinquent behaviors. Observed and model-fitted means for measures in which there was a significant intervention effect are shown in Figure 1.

Figure 1A shows that at baseline the average number of sexual partners was less than one among both intervention and control participants, including zero values for the 47% of participants who had not yet had sexual intercourse. During a time of life when many

American adolescents are initiating sexual activity (36), the mean number of recent partners estimated by our model for the intervention group dropped from 0.84 at baseline to 0.46 at 12 months. Among controls, it increased from 0.72 to 1.72 over the same time period, resulting in a highly significant intervention effect ($p < .001$). These results imply that relatively few youth in the intervention arm initiated sexual activity, and those with one or more partners at baseline reduced that number over time.

Youth in both the intervention and control conditions reduced alcohol use over the study period, as shown in Figure 1B. Intervention participants reduced significantly more than controls, going from an estimated mean of 7.7 times in 3 months to 1.8 times, as compared to controls who decreased from 5.7 to 4.3 times ($p = .003$). Marijuana use, shown in Figure 1C, showed the opposite effect, with intervention participants increasing use over the time period (from 9 to 12 times in a 3-month period) while controls dropped (from 13 to 6 times, $p < .001$). Estimates of the amount of hard drug use, as shown in Figure 1D, indicate drops among youth in both arms of the study, with those in the intervention experiencing greater decreases in use, going from 2.8 to 0.3 times recently as compared to controls who drop from 2.7 to 1.2 ($p < .001$).

Delinquent behaviors (Fig. 1E) also declined more in the intervention condition (2.2 to 0.6) compared to the decreases among controls (2.6 to 1.5) ($p = .001$).

DISCUSSION

While other interventions have worked effectively with individual homeless youth to reduce risk-taking behaviors in the context of street life (15), a family-based program can also result in significant reductions in risky behavior. Antecedent problems of family conflict that lead to runaway incidents (1, 37) were the focus of the STRIVE intervention. We aimed to re-engage youth with their families. The intervention reduced the number of sex partners, and youth use of hard drugs and alcohol – activities associated with increased risk of HIV infection. Yet, youth in the intervention condition increased their marijuana use relative to youth in the control condition; perhaps substituting marijuana for alcohol or other hard drugs. The intervention also reduced delinquent behaviors. Increasingly family interventions are being suggested to address at-risk behaviors among youth (17, 20, 25, 27, 38). Moreover, the STRIVE intervention is only five sessions which is shorter than most family interventions (18, 19, 21, 22, 39). These findings suggest that STRIVE is an efficacious intervention that should be replicated with homeless youth in other urban settings and rural settings, and disseminated in the service sector to determine its effectiveness as a family intervention for homeless youth.

The primary challenge and limitation of this study was the difficulty in recruiting “whole” families to the intervention. Only 33% (151/442) of the families were successfully randomized because of the “research” mission of the project. The consenting and randomization process was complex, challenging and time-consuming. We had to obtain assent/consent from the youth, consent from the parent, conduct baseline assessments with both the youth and the parent before the completion of the recruitment process and randomization could occur. Youth were approached first and some did not want to be reunited with their families so we could not approach their parents who might have been more willing to take part in the study. Our assessment team was flexible in scheduling with families to streamline the consenting and randomization process (e.g., interviewers available to conduct baseline of youth and parent individually at the same time), but having to go through consenting and a baseline assessment, usually two home visits, was difficult from some families to integrate into their busy lives at the onset of the study before they were fully engaged in the study. A limitation that resulted from the challenge in recruiting

families was that some of the participants had their first follow up assessment before the last intervention session occurred. Another limitation is that protections for the ethical conduct of research prevented us from collecting any data from families that did not fully complete the recruitment process; as a result we cannot compare families from whom consent was not obtained. Although randomization eliminates many biases, we cannot evaluate selection effects for both less dysfunctional and conflicted or more dysfunctional and conflicted families. Last, the data are self-report. There may be a tendency for participants in the intervention condition to under report their risky behaviors. We acknowledge that this is a limitation but the use of ACSI does lessen some of this under reporting in studies of risky behaviors that rely on self-report data (40).

These challenges and limitations suggest an important avenue for future research in risk prevention with newly homeless and runaway youth. Innovative approaches for engaging parents of high-risk youth need to be developed. It was relatively easy to find youth interested in participating in the study, but much more difficult to enroll their parents, many of whom were extremely frustrated with their children. Recruitment strategies are needed that directly target parents, normalize potential sources of parent/adolescent conflict (e.g., adolescents' need for autonomy), and do not blame parents or adolescents for family conflict. This study highlights the importance of focusing public health efforts on identifying youth when they are newly homeless and targeting their families to stop the evolution of riskier behavior patterns.

Acknowledgments

This study was funded by the National Institute of Mental Health (NIMH R01-MH070322).

Abbreviations

ACASI	Audio computer-assisted self-interviewing
STRIVE	Support To Reunite, Involve and Value Each Other [intervention]
WAI	Working Alliance Inventory

REFERENCES

1. Ringwalt CL, Greene JM, Robertson MJ, McPeeters M. The prevalence of homelessness among adolescents in the United States. *Am J Public Health.* 1998; 88(9):1325–1329. [PubMed: 9736871]
2. Center for Disease Control and Prevention. [Accessed October 4, 2010] HIV prevalence among selected populations: youth populations [June 2006]. Available at: http://www.cdc.gov/hiv/topics/testing/resources/reports/hiv_prevalence/youth.htm
3. Pfeifer RW, Oliver J. A study of HIV seroprevalence in a group of homeless youth in Hollywood, California. *J Adolesc Health.* 1997; 20:339–342. [PubMed: 9168380]
4. Stricof RL, Kennedy JT, Nattell TC, Weisfuse IB, Novick LF. HIV seroprevalence in a facility for runaway and homeless adolescents. *Am J Public Health.* 1991; 81 Suppl.:50–53. [PubMed: 2014885]
5. Milburn N, Rotheram-Borus MJ, Rice E, Mallet S, Rosenthal D. Cross-national variations in behavioral profiles among homeless youth. *Am J Community Psychol.* 2006; 37:63–76. [PubMed: 16680537]
6. Solorio R, Milburn NG, Rotheram-Borus MJ, Higgins C, Gelberg L. Predictors of sexually transmitted infection testing among sexually active homeless youth. *AIDS Behav.* 2006; 10(2):179–184. [PubMed: 16479414]
7. Forney JC, Lombardo S, Toro PA. Diagnostic and correlates of HIV risk behaviors in a probability sample of homeless adults. *Psychiatr Serv.* 2007; 58(1):92–99. [PubMed: 17215418]

8. Tyler KA. Social network characteristics and risky sexual and drug-related behaviors among homeless young adults. *Soc Sci Res.* 2008; 37(2):673–685. [PubMed: 19069065]
9. Rice E, Milburn NG, Rotheram-Borus MJ, Mallett S, Rosenthal D. The effects of peer-group network properties on drug use among homeless youth. *Am Behav Sci.* 2005; 48(8):1102–1123. [PubMed: 20539820]
10. McMorris B, Tyler KA, Whitbeck LB, Hoyt DR. Familial and on the street risk factors associated with alcohol use among homeless and runaway adolescents. *J Stud Alcohol.* 2002; 63:34–43. [PubMed: 11925056]
11. Slesnick N, Meade M. System youth: a subgroup of substance-abusing homeless adolescents. *J Subst Abuse.* 2001; 13(3):367–384. [PubMed: 11693458]
12. Whitbeck LB, Hoyt DR, Bao WN. Depressive symptoms and co-occurring depressive symptoms, substance abuse, and conduct problems among runaway and homeless adolescents. *Child Dev.* 2000; 71(3):721–732. [PubMed: 10953939]
13. Milburn NG, Rosenthal D, Rotheram-Borus MJ, et al. Newly homeless youth typically return home. *J Adolesc Health.* 2007; 40(6):574–576. 2007. [PubMed: 17531769]
14. Milburn NG, Rotheram-Borus MJ, Rice E, et al. Adolescents exiting homelessness over two years: the risk amplification and abatement model. *J Res Adolesc.* In press.
15. Rotheram-Borus MJ, Song J, Gwadz M, Lee M, Van Rossem R, Koopman C. Reductions in HIV risk among runaway youth. *Prev Sci.* 1993; 4(3):173–187. [PubMed: 12940468]
16. Slesnick N, Prestopnik JL. Ecologically based family therapy outcome with substance abuse runaway adolescents. *J Adolesc.* 2005; 28:277–298. [PubMed: 15878048]
17. Taylor TK, Biglan A. Behavioral family interventions for improving child-rearing: a review of the literature for clinicians and policy makers. *Clin Child Fam Psychol Rev.* 1998; 1(1):41–60. [PubMed: 11324077]
18. Rotheram-Borus MJ, Piacentini J, Miller S, Graae F, Castro-Blanco F. Brief cognitive behavioral treatment for adolescent suicide attempts and their families. *J Am Acad Child Adolesc Psychiatr.* 1994; 33(4):508–517.
19. Henggeler, SW.; Schoenwald, SK.; Borduin, CM.; Rowland, MD.; Cunningham, PB. *Multisystemic Treatment of Antisocial Behavior in Children and Adolescents.* New York, NY: Guilford Press; 1998.
20. Spoth RL, Kavanagh KA, Dishion TJ. Family-centered prevention intervention science: toward benefits to larger populations of children, youth and families. *Prev Sci.* 2002; 3(3):145–152. [PubMed: 12387551]
21. Dishion TJ, Kavanagh K, Nelson S, Kaufman NK. Preventing early adolescent substance use. *Prev Sci.* 2002; 3(3):191–201. [PubMed: 12387554]
22. Ennett ST, Bauman KE, Pemberton M, Foshee VA, Chuang YC, King TS. Mediation in family-directed program for prevention of adolescent tobacco and alcohol use. *Prev Med.* 2001; 33(4):333–346. [PubMed: 11570838]
23. Catalano RF, Hawkins JD, Berglund ML, Pollard ML, Arthur MW. Prevention science and positive youth development: competitive or cooperative framework. *J Adolesc Health.* 2002; 31(6):230–239. [PubMed: 12470920]
24. Kumpfer KL, Alvarado R, Smith P, Bellamy N. Cultural sensitivity and adaptation in family-based prevention interventions. *Prev Sci.* 2002; 3(3):241–246. [PubMed: 12387558]
25. Rotheram-Borus, MJ.; Goldstein, AM.; Elkavitch, AS. Treatment of suicidality. In: Hoffman, S.; Thompson, M., editors. *Psychosocial Treatments of Severe Mental Disorders.* New York, NY: Guilford Press; 2002.
26. Bandura, A. *Social Foundations of Thought and Action: A Social-cognitive View.* Englewood Cliffs: Prentice Hall; 1986.
27. Dishion, TJ.; Kavanagh, K. *Intervening in Adolescent Problem Behavior: A Family-centered Approach.* New York, NY: Guilford Press; 2003.
28. Kagen, R.; Schlossberg, S. *Families in Perpetual Crisis.* New York, NY: Norton; 1989.
29. Rotheram-Borus MJ, Piacentini J, Cantwell C, Belin TR, Song J. Prevention of HIV among adolescents. *J Consult Clin Psychol.* 2000; 68(6):1081–1093. [PubMed: 11142542]

30. Walker LS, Greene JW. Negative life events, psychosocial resources, and psychophysiological symptoms in adolescents. *J Clin Child Psychol.* 1987; 16:29–36.
31. Patterson C. Children of lesbian and gay parents. *Child Dev.* 1992; 63:1025–1042. [PubMed: 1446541]
32. Stuart, R. *Helping Couples Change.* New York, NY: Guilford Press; 1980.
33. Horvath AO, Greenburg LS. Development and validation of the Working Alliance Inventory. *J Couns Psychol.* 1989; 36(2):223–233.
34. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition.* Washington, DC: American Psychiatric Association; 1994.
35. Centers for Disease Control, Division of Adolescent and School Health. [Accessed July 30, 2009] Data and statistics, YRBSS: Youth Risk Behavior Surveillance System. <http://www.cdc.gov/HealthyYouth/YRBS/index.htm>
36. Gavin, L.; MacKay, AP.; Brown, K., et al. [Accessed July 30, 2009] Sexual and reproductive health of persons aged 10–24 years—United States, 2002–2007; *MMWR Morb Mortal Wkly Rep.* 2009. p. 1–62. <http://www.cdc.gov/mmwr/PDF/ss/ss5806.pdf>
37. Rotheram-Borus MJ, Bradley J. Triage model for suicidal runaways. *Am J Orthopsychiatry.* 1991; 61:122–127. [PubMed: 2006668]
38. Sanders MR, Turner KMT, Markie-Dadds C. The development and dissemination of the Triple P —Positive Parenting Program: a multilevel, evidence-based system of parenting and family support. *Prev Sci.* 2002; 3:173–189. [PubMed: 12387553]
39. Bear JS, Garrett SB, Beadwell B, Wells EA, Peterson P. Brief motivational intervention with homeless adolescents: Evaluating effects on substance use and service utilization. *Psych Addict Behav.* 2007; 21:582–586.
40. Vanable PA, Carey MP, Brown MS, DiClemente RJ, Salzar LF, Brown LK, Romer D, Valois RF, Hennessy M, Stanton BF. Test-retest reliability of self-reported HIV/STD-related measures among African-American adolescents in four U.S. cities. *J Adolesc Health.* 2009:214–221. [PubMed: 19237106]

E. Number of delinquent behaviors

Figure 1.

Estimated Regression Lines and Observed Values for Outcome Measures: Adolescents in the Intervention Condition (solid lines, black triangles) and Control Condition (broken lines, white diamonds)*†

* Effect sizes are calculated from model estimated (ME) values as: $((ME_{\text{control, 12-month}} - ME_{\text{control, baseline}}) - (ME_{\text{intervention, 12-month}} - ME_{\text{intervention, baseline}})) / \text{standard deviation for the baseline value in the control condition}$.

† Significance testing is for the difference in slopes between the intervention and control conditions (time by treatment interaction). Analyses were conducted using SAS 9.1 software.³⁶ Continuous measures were modeled with the PROC MIXED procedure. Dichotomous and Poisson-distributed count measures were modeled with the PROC GLIMMIX procedure. No significant intervention effect was found for whether the adolescent had been sexually active, had unprotected sex, number of times had sex, whether used alcohol or marijuana, whether used hard drugs.

Table 1

Content of the STRIVE Intervention Sessions

Session	Description
1	Create a positive family atmosphere Establish facilitator's credibility Negotiation modeling Plan for and identify potential emergencies Connect feelings to behaviors Identify outside social supports
2	Identify and rank problem situations Relate feeling thermometer to each problem Assign priorities to each identified family problem Increase problem solving abilities Select a relatively easy family problem and practice problem-solving
3	Analyze a problem of moderate difficulty Uncover obstacles: rules, roles, assumptions, benefits of keeping the status quo Select a family problem of moderate difficulty and practice problem-solving Review HIV and street life educational materials
4	Decide how to cope with the problem(s) Select a family problem of medium to high difficulty and problem-solve Learn how to negotiate solutions
5	Select the family problem of the highest difficulty (i.e., the one with the highest probability of being the main source of family conflict) and practice problem-solving Evaluate solutions, and implement them through negotiation Review family's own strengths as problem solvers

TABLE 2

Observed Characteristics of STRIVE Participants at Baseline, by Intervention Status*

	Intervention (N = 68)	Control (N = 83)	Total (N = 151)
Age, Mean (SD)	14.7 (1.3)	14.9 (1.5)	14.8 (1.4)
Gender, N (%)			
Male	15 (22.1)	36 (43.4)	51 (33.8) [†]
Female	53 (77.9)	47 (56.6)	100 (66.2)
Sexual orientation, N (%)			
Heterosexual	60 (88.2)	76 (91.6)	136 (90.1)
Bisexual, Gay, Lesbian	8 (11.8)	7 (8.4)	15 (9.9)
Race/ethnicity, N (%)			
Hispanic	42 (61.8)	51 (61.4)	93 (61.6)
White	8 (11.8)	9 (10.8)	17 (11.3)
African American	12 (17.6)	19 (22.9)	31 (20.5)
Other, mixed	6 (8.8)	4 (4.8)	10 (6.6)
Born in this country, N (%)	62 (91.2)	77 (92.8)	139 (92.1)
Longest time ever away, N (%)			
2 weeks or less	43 (64.2)	52 (62.7)	95 (63.3)
3 weeks to 1 month	17 (25.4)	18 (21.7)	35 (23.3)
2 to 6 months	7 (10.4)	13 (15.7)	20 (13.3)
Where currently living, N (%)			
Birth or adoptive family	53 (77.9)	55 (66.3)	108 (71.5)
Other family or friends	9 (13.2)	11 (13.3)	20 (13.2)
Shelter, group home, other	6 (8.8)	17 (20.5)	23 (15.2)
In the 3 months prior to baseline:			
Had vaginal or anal sex, N (%)	26 (38.2)	33 (39.8)	59 (39.1)
Had unprotected sex (without a condom), N(%)	17 (25.0)	20 (24.1)	37 (24.5)
Number of times had sex, Mean (SD)	4.1 (12.3)	3.0 (7.7)	3.5 (10.0)
Number of partners, Mean (SD)	0.8 (1.5)	0.8 (1.4)	0.8 (1.4)
Used alcohol, N (%)	29 (43.3)	39 (47.0)	68 (45.3)
Times used alcohol, Mean (SD)	8.5 (25.9)	5.5 (11.9)	6.8 (19.4)

	Intervention (N = 68)	Control (N = 83)	Total (N = 151)
Used marijuana, N (%)	30 (44.1)	42 (50.6)	72 (47.7)
Times used marijuana, Mean (SD)	9.9 (29.0)	11.6 (25.1)	10.9 (26.9)
Used hard drugs, [‡] N (%)	14 (20.9)	22 (26.5)	36 (24.0)
Times used hard drugs, Mean (SD)	2.5 (9.4)	2.8 (6.6)	2.7 (7.9)
Number of delinquent behaviors, [§] Mean (SD)	2.4 (2.0)	2.8 (2.4)	2.6 (2.2)

* For characteristics expressed as means, t-tests were performed; for characteristics expressed as percents, chi square tests were performed.

[‡] $p < .01$.

[‡] Hard drugs defined as cocaine; crack; amphetamines; ice/smoked speed; heroin, non-prescription methadone; other opiates, narcotics or painkillers; barbiturates; tranquilizers; inhalants; party drugs; or other drugs.

[§] Out of 13 possible: bullied/threatened, started a fight, used a weapon, physically been cruel, tortured/hurt an animal, stole face-to-face, forced sex, set a fire, destroyed property, broke into a building, lied to get something, stole from someone not present, skipped class.

TABLE 3

Comparisons of STRIVE Participants with a Nationally Representative Sample of 9th–12th Graders (Youth Risk Behavior Survey, 2009)*

	<u>YRBS</u>	<u>STRIVE</u>
Sample size	16,410	151
Used alcohol, lifetime [†]	72.5 (70.6–74.3)	72.2 (68.4–79.2)
Used marijuana, lifetime	36.8 (34.8–38.8)	58.9 (51.1–66.8)
Used cocaine, lifetime	6.4 (5.7–7.1)	17.2 (11.2–23.2)
Used methamphetamines, lifetime	4.1 (3.6–4.6)	12.6 (7.3–17.9)
Used inhalants, lifetime	11.7 (10.6–12.8)	11.3 (6.2–16.3)
Had sex, lifetime	46.0 (42.9–49.2)	53.3 (45.4–61.3)
If had sex, used a condom during last sexual intercourse	61.1 (59.0–63.1)	45.3 (34.8–55.9)
If had sex, used alcohol or drugs before last sexual intercourse	21.6 (20.0–23.3)	14.0 (6.6–21.3)

* Source: CDC National Center for Chronic Disease Prevention and Health Promotion, Healthy Youth, YRBS National Trends in Risk Behaviors (www.cdc.gov/HealthyYouth/YRBS/trends.htm).

[†] Percent (95% confidence interval)