Published in final edited form as:

Pain. 2010 November ; 151(2): 257–265. doi:10.1016/j.pain.2010.05.022.

Reversal of pancreatitis-induced pain by an orally available, small molecule interleukin-6 receptor antagonist

Marina Vardanyan[#], Ohannes K Melemedjian,

Department of Pharmacology, College of Medicine, University of Arizona, Tucson, Arizona 85724

Theodore J Price,

Department of Pharmacology, College of Medicine, University of Arizona, Tucson, Arizona 85724

Michael H. Ossipov,

Department of Pharmacology, College of Medicine, University of Arizona, Tucson, Arizona 85724

Josephine Lai,

Department of Pharmacology, College of Medicine, University of Arizona, Tucson, Arizona 85724

Ed Roberts¹, Terrence L. Boos², Jeffrey R Deschamps³, Arthur E Jacobson², Kenner C Rice², and Frank Porreca

Department of Pharmacology, College of Medicine, University of Arizona, Tucson, Arizona 85724

¹Scripps Research Institute, La Jolla, CA 92037

²Drug Design and Synthesis Section, Chemical Biology Research Branch, National Institute on Drug Abuse and the National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda, MD 20892

³Laboratory for the Structure of Matter, Naval Research Laboratory, Washington DC 20375

Abstract

Pancreatic pain resulting from chronic inflammation of the pancreas is often intractable and clinically difficult to manage with available analgesics reflecting the need for more effective therapies. Mechanisms underlying pancreatitis pain are not well understood. Here, the possibility that interleukin-6 (IL-6) may promote pancreatitis pain was investigated with TB-2-081 (3-*O*-formyl-20R,21-epoxyresibufogenin, EBRF), a small molecule IL-6 receptor antagonist that was semi-synthetically derived from natural sources. The potential activity and mechanism of TB-2-081 was investigated following induction of persistent pancreatitis using dibutyltin dichloride (DBTC) in rats. TB-2-081 displaces binding of IL-6 to the human recombinant soluble IL-6 receptor with apparent high affinity and inhibits IL-6 mediated cell growth. Systemic or oral, but not intrathecal, administration of TB-2-081 reversed DBTC-induced abdominal hypersensitivity in a dose- and time-dependent manner. IL-6 levels were significantly upregulated in the dorsal root ganglia (DRG) of rats with pancreatitis on day 6 after DBTC injection. IL-6

© 2010 International Association for the Study of Pain. Published by Elsevier B.V. All rights reserved.

Correspondence: Frank Porreca, PhD Department of Pharmacology College of Medicine, University of Arizona Health Sciences Center Tucson, Arizona 85724 frankp@email.arizona.edu Phone: (520)-626-7421 Fax: (520) 626-4182 .

[#]Current address: Department of Surgery, College of Medicine, University of Arizona, Tucson, Arizona 85724

CONFLICT OF INTEREST. All authors declare that they have no conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Financial Disclosures: The authors have nothing to disclose

enhanced capsaicin-evoked release of calcitonin gene related peptide from cultured DRG neurons was blocked by TB-2-081. Our data demonstrate that TB-2-081 acts as a systemically available and orally active small molecule IL-6 receptor antagonist. TB-2-081 effectively reduces pancreatitis-induced pain through peripheral mechanisms that are likely due to (a) increased expression of IL-6 in the DRG and (b) IL-6-mediated sensitization of nociceptive neurons. The activity of TB-2-081 implicates an important role for IL-6 in sustaining pancreatitis pain. Strategies targeting IL-6 actions through small molecule antagonists may offer novel approaches to improve therapy of chronic pancreatitis and other chronic pain states.

Introduction

Chronic pancreatitis is a continuous inflammatory disease of the pancreas characterized by irreversible morphologic changes that typically cause pain and permanent loss of function. Pain is the most challenging and debilitating symptom associated with this condition. Initial treatment is generally conservative and includes non-opioid analgesics, followed by opioids [9]. However, a sub-group of patients ultimately develop severe, incapacitating epigastric pain that markedly reduces their quality of life [34]. Conservative pain control efforts in these patients can lead to a vicious cycle of increasing need for narcotic analgesics and significant unwanted side-effects including somnolence, mental confusion and gastrointestinal problems [3] as well as potential for addiction. Therefore, there is a high and unmet need for novel therapies for the treatment of pancreatitis-induced pain.

The pleiotropic cytokine, interleukin 6 (IL-6) is markedly up-regulated during various pathologic conditions and is generally associated with increased pain and hyperalgesia. Most experimental studies report pro-inflammatory and pro-nociceptive roles for IL-6. Intramuscular, intradermal, intracerebroventricular, or intrathecal injection of IL-6 induces allodynia or hyperalgesia in rats [6;30]. Several experimental models have shown that IL-6 is a mediator of hyperalgesia in inflamed tissues. Moreover, overproduction of IL-6 contributes to the pathogenesis of various autoimmune and inflammatory diseases, including rheumatoid arthritis (RA), systemic-onset juvenile chronic arthritis (JCA), osteoporosis, psoriasis, insulin-dependent diabetes mellitus, and inflammatory bowel disease [14;25]. IL-6 has been suggested to play a role in acute pancreatitis and pancreatic cancer [13;27;32].

IL-6 is produced by lymphoid and non-lymphoid cells, such as T cells, B cells, monocytes, pancreatic acinar cells, fibroblasts, keratinocytes, endothelial cells, mesangial cells and several tumor cells, including myeloma and hypernephroma cells [18]. When IL-6 reaches an IL-6-responsive cell, it binds to its specific receptor, IL-6R, that exists in both transmembrane and soluble forms [33]. Binding of IL-6 triggers an association of the IL-6R with the transducer protein, gp130 [12], initiating an intracellular cascade of phosphorylation of several signal proteins including Janus kinases, STAT factors [11] as well as the MAPKs [16;41]. Based on IL-6's known role in inflammation and pain promoting properties we hypothesized that IL-6 might be an important mediator of pancreatitis-induced pain. Inhibition of IL-6 signaling, therefore, may represent a potential non-narcotic therapeutic target for the treatment of this disorder.

TB-2-081(3-O-formyl-20R,21-epoxyresibufogenin) [17] (Fig. 1A; structure confirmed by X-ray crystallographic analysis, MW = 428.5) has been identified in extracts of Chinese toad skin [15] and has been reported to function as an IL-6R antagonist [7;17]. However, its potential activity in experimental models of pain is not known. Our results indicate that TB-2-081 is an orally active small molecule antagonist of the IL-6R, and is effective for alleviation of chronic pancreatitis pain in rats. Our data suggest that IL-6 may enhance nociceptive sensitivity in conditions of persistent pancreatitis.

Materials and Methods

Competition of IL-6 binding to human recombinant soluble IL-6R by TB-2-081

Ninety-six well-plates (Nunc) were coated with 500 ng/mL of human recombinant soluble IL-6 receptor (R&D Systems) overnight at 4°C. Blank wells were used as control. The plates were washed 3 times with 300 µL of phosphate buffered saline (PBS) and pre-blocked with 1% BSA (Sigma) in PBS at room temperature (RT) for 1 hr and washed 3 times with PBS. Twelve concentrations (50 pM – 100 nM) of human recombinant IL-6 (R&D Systems) were added to the wells, in triplicates, and incubated at RT with shaking for 2 hr. After washing to remove unbound IL-6, the wells were incubated with a biotinylated polyclonal anti-human IL-6 antibody (200 ng/mL, R&D Systems) at room temperature for 2 hr, washed and incubated with streptavidin-horseradish peroxidase (R&D Systems) for 20 min. Sample wells were washed and developed with tetramethylbenzidine (100 µL/well; 5 to 10 min at RT and stopped with 1M HCl). Optical density in each well was determined using a microplate reader (Multiskan Ascent, Thermo) set to 450 nm with a correction wavelength of 570 nm. Non-specific binding of IL-6 is defined by that in the absence of sIL-6R. Data were analyzed by non-linear regression analysis using GraphPad Prism4 (Graph Pad, San Diego, CA). Based on the titration analysis, competition of IL-6 binding to sIL-6R by TB-2-081 was carried out as follows. Ninety-six well-plates were prepared as described above. After pre-block with the BSA containing buffer, wells were incubated with 1.2 nM of recombinant human IL-6, or 1.2 nM IL-6 plus one of six concentrations of TB-2-081 (10^{-12} - 10^{-7} M), in triplicates. All incubation conditions and colorimetric development were as described above. Maximum binding of IL-6 (100% bound) is defined as the absorbance in the absence of TB-2-081. The absorbance in the presence of various concentrations of TB-2-081 is calculated as percent of maximum binding of IL-6. Data were analyzed by non-linear regression analysis to determine the IC₅₀ value of TB-2-081 using GraphPad Prism4 (Graph Pad, San Diego, CA).

Functional assay for IL-6

TF-1 is a factor-dependent human erythroleukemic cell line that proliferates in response to many cytokines including human IL-6 [5]. The cells were grown and maintained in RPMI 1640 / 10% fetal bovine serum/ 2mM L-glutamine / 100 units/mL penicillin / 100 µg/mL streptomycin / 2 ng/ml rhGM-CSF (R&D Systems). Cells were seeded at 50,000 cells/mL or higher in the medium and were passaged every 3 to 4 days with fresh medium. For the bioassay, cells were collected and washed with PBS and suspended at 200,000 cells/mL in the above medium but omitting rhGM-CSF (assay medium). Human recombinant IL-6 (2 ng/mL), or various doses of the drug with or without IL-6 were prepared in assay medium in a final volume of 50 µL in 96-well plates (BD Biosciences), and incubation was initiated by adding 50 µL of the cell suspension to each well. The plates were incubated for 72 hr at 37°C in humidified 95% air/5% CO₂. Cell proliferation was measured by CellTiter® 96 Aqueous One Solution Cell Proliferation Assay (Promega). Data were analyzed by non-linear regression analysis using GraphPad Prism4 (Graph Pad, San Diego, CA).

Animals

Male Sprague-Dawley rats (Harlan, Indianapolis, IN), weighing 150-200 g at the time of testing, were maintained in a climate-controlled room on a 12 hour light-dark cycle (lights on at 07:00 hours) and food and water were available *ad libitum*. All testing was performed in accordance with the policies and recommendations of the International Association for the Study of Pain (IASP) and National Institute of Health (NIH) and received approval from the Institutional Animal Care and Use Committee of the University of Arizona.

Induction of pancreatitis

Chronic pancreatic inflammation was induced by tail vein injection of dibutyltin dichloride (DBTC, Aldrich Milwaukee, WI) dissolved in 100% ethanol vehicle at a dose of 8 mg/kg under isoflurane anesthesia (2-3 liters/min of 4% isoflurane delivered in a mixture of 95% O₂ and 5% CO₂ to induce anesthesia and reduced to 2.5% isoflurane to maintain anesthesia) throughout the procedure [39;40]. Control rats received the same volume of ethanol vehicle solution only. This is a well established model of experimental pancreatitis that produces elevations of pancreatic enzymes and histological evidence of pancreatic inflammation without evidence of histological damage to other organs including the liver, heart, kidney or lungs [39;40]. Elevation in serum levels of amylase and lipase caused by DBTC injection have also been previously reported [38,39], consistent with what is observed in humans [10].

Intrathecal catheter implantation—Implantation of intrathecal catheters was performed as described by Yaksh and Rudy [44]. The rats were anesthetized with a mixture of ketamine (80 mg/kg, i.p.) and xylazine (12 mg/kg, i.p.) and a 5.5 cm length of PE-10 tubing was inserted through an incision made in the atlanto-occipital membrane, to the level of the lower thoracic levels (T₈-T₁₂). The catheter was then secured to the musculature at the site of incision, which was then closed. Drugs were dissolved in saline and administered in a volume of 5 µL through a length of PE20 tubing connecting the catheter with the injection syringe. The catheter was cleared by flushing with 9 µL saline after drug. The animals were allowed to recover for 6 days after surgery before pharmacological manipulations were made.

Behavioral testing: Assessment of pancreatitis-induced pain

Pancreatitis pain was assessed by referred abdominal hypersensitivity to probing the upper left abdominal quadrant of rats with calibrated von Frey filaments (4 g). Rats were allowed to acclimate in a suspended wire-mesh cage for 30 min before testing. A response was indicated by the sharp withdrawal of the abdomen, licking of abdominal area, or whole body withdrawal. A single trial consisted of 10 applications of a von Frey filament applied once every 10 s (intervals to allow the animal to return to a relatively inactive position). The mean occurrence of withdrawal events in each trial is expressed as the number of responses to 10 applications [39;40]. A significant increase in the number of abdominal withdrawals is interpreted as “pancreatitis-induced pain” [36,37].

Acute thermal nociception—The 52°C hot-plate test was employed to assess potential antinociceptive activity of TB-2-081. Rats were placed on a metallic hot-plate maintained at 52±0.5°C and latency to withdrawal, stamping or licking of the hindpaw was determined. A significant ($P < 0.05$) elevation in hot-plate latency from baseline values indicated antinociception.

ELISA assay for IL-6

Pancreas and the lower thoracic DRG (T₈-T₁₂) innervating the pancreas were collected and frozen in -80°C before use. The tissue was ultrasonicated in PBS (pH7.4) in the presence of protease inhibitor cocktail (Complete Protease Inhibitor Cocktail Tablets, Roche) after which the samples were centrifuged and the supernatant was collected. IL-6-like immunoreactivity was determined by ELISA (IL-6 ELISA kit, R&D systems) according to the manufacturer's instructions.

CGRP release from adult rat DRG culture

Male Sprague-Dawley rats (200-250 g) were used. Dorsal root ganglia (DRG) from all levels (~45/animal) were excised. After the ganglia were freed of capsular connective tissue

and the nerve trunk, DRG were dissociated enzymatically with collagenase A (1 mg/mL, 25 min) and collagenase D (1 mg/mL) with papain (30 U/mL) for 20 min at 37°C. Dissociated cells were seeded in 24 well titer plates and incubated at 37°C in a humidified 95% air/5% CO₂ incubator. On day 4 cells were incubated in media containing IL-6 only (10 ng/mL), TB-2-081 only (5 µM), or both drugs, for 30 min, followed by incubation with media containing the same drug(s) and capsaicin (30 nM) for an additional 15 min. Media were collected from the sample wells and measured for CGRP-like immunoreactivity by ELISA according to the manufacturer's instructions (CGRP (rat) ELISA, Cayman Chemicals).

Data analysis—The results of withdrawal events are presented as mean ± SEM. Statistical analysis was performed in FlashCalc (www.u.arizona.edu/~michaelo). One-way ANOVA was used to detect significant differences in behavioral outcomes within each experimental group over time. Two-factor ANOVA was employed to detect changes between treatment groups over time. In order to generate dose-response curves, data were converted to % inhibition by the formula: $100 \times (\text{maximum} - \text{response}) / \text{maximum}$, where maximum = 10 withdrawals. Linear regression analysis of the logdose-response curves was used to determine the A₅₀ (95% confidence limits), representing the dose producing a 50% response. The area under the curve (AUC) for each of the time-effect curves at each dose were calculated from % inhibition vs time by using the trapezoid rule. For all statistical comparisons, significance was established at $P = 0.05$.

Results

TB-2-081 competes with the binding of IL-6 to sIL-6R

IL-6 binds to the sIL-6R in a saturable, concentration dependent manner. Under the assay conditions, the half saturation concentration of IL-6 was 960 pM (Figure 1B). TB-2-081 displaced the binding of IL-6 (1.2 nM) to sIL-6R with an IC₅₀ value of 29.4 pM and a Hill slope of -1.0 suggesting a single binding site for TB-2-081 (Figure 1C).

TB-2-081 antagonizes the biological activity of IL-6

TB-2-081 inhibited IL-6 mediated growth of IL-6 dependent TF-1 cells in a dose dependent manner with an IC₅₀ of 250 pM (Figure 1D). Treatment with TB-2-081 alone had no effect on the growth of TF-1 cells, suggesting that TB-2-081 acts as an antagonist of IL-6R.

Treatment with TB-2-081 blocks pancreatitis-induced referred abdominal hypersensitivity

Pancreatitis-induced pain was evaluated by probing the abdomen with von Frey filaments on day 6 after injection of DBTC. The mean number of responses to 10 applications of von Frey filaments was 8.33 ± 0.18 in the DBTC treated rats compared with 0 ± 0.0 in ethanol vehicle treated control rats, indicating significant abdominal hypersensitivity when compared to pre-injection baselines. To investigate the potential role of IL-6 in pancreatitis-induced referred abdominal hypersensitivity, rats were given TB-2-081 (0.001, 0.01, 0.1 or 1 mg/kg) by s.c. injection on day 6 after induction of chronic pancreatitis. TB-2-081 produced a dose- and time-dependent reversal of referred abdominal hypersensitivity while injection of the vehicle (i.e.; 50% PBS and 50% ethanol) was without effect (Figure 2A). The peak inhibitory effect of TB-2-081 occurred rapidly, 15 min after subcutaneous administration and lasted for about 60 min. Injection of the ethanol vehicle (i.e.; DBTC vehicle) plus PBS/ethanol (TB-2-081 vehicle) did not produce increases in numbers of withdrawal (Figure 2A). The area under the time-effect curve analysis of the TB-2-081 effect suggests that the compound abolishes referred abdominal hypersensitivity in a dose- and time-dependent manner, consistent with a specific receptor mechanism (Figure 2B,C). The A₅₀ value for s.c. TB-2-081 was 0.006 mg/kg (0.0001-0.26 mg/kg; 95% confidence limits).

Based on the potent activity of systemically-administered TB-2-081, we explored the possibility that this molecule might also have oral activity. TB-2-081 was given by oral gavage at doses of 0.1 mg/kg or 1 mg/kg on day 6 after DBTC induction of pancreatitis. Oral TB-2-081 produced dose- and time-dependent reversal of referred abdominal hypersensitivity (Figure 3A,B). Analysis of the area under the time-effect curve for orally administered TB-2-081 shows that the compound reverses referred abdominal hypersensitivity induced by pancreatitis in a time- and dose-dependent manner (Figure 3B). The potency of orally administered TB-2-081 was approximately one-fifth that of s.c. administration. In contrast, the oral administration of up to 3 mg/kg of TB-2-081, representing 3 times the effective dose against pancreatitis-induced abdominal hypersensitivity, did not reverse acute nociception in the 52°C rat hot plate test (data not shown).

In order to determine if TB-2-081 acted through central sites, a dose of 5 µg was administered i.th. to rats with referred abdominal hypersensitivity on day 6 after DBTC administration. Intrathecal TB-2-081 did not reverse abdominal hypersensitivity in rats with DBTC-induced pancreatitis (data not shown).

IL-6 levels in the pancreas and DRG after the induction of pancreatitis

The results of the behavioral analysis suggested that endogenous IL-6 may drive pancreatitis pain. To determine if this may be due to increased synthesis of IL-6, the levels of IL-6 were quantified in pancreatic tissue and in thoracic DRGs from naïve rats and from treated rats 6 days following ethanol vehicle or DBTC injection.

A significant ($p < 0.05$) upregulation of IL-6 was seen in the pancreas in both ethanol vehicle and DBTC treated groups when compared with that from naïve rats (Figure 4A). Additional analyses over a period of 1 to 14 days after ethanol vehicle or DBTC treatment showed that increased levels of IL-6 were already evident one day after administration of either ethanol vehicle or DBTC and these levels remained elevated on day 6 post-injection. By day 14 post-injection, pancreatic levels of IL-6 returned to baseline levels for the ethanol vehicle group but remained elevated in the DBTC group (Figure 4A). There were no significant ($p > 0.05$) changes in IL-6 levels in liver, thoracic spinal cord or brain tissue on day 6 after DBTC (data not shown). This is consistent with our previous reports that DBTC did not affect morphologic changes in the liver, intestines, lungs, kidneys or heart [39;40].

Levels of IL-6 were also measured in the T8-T12 thoracic DRG on day 6 after DBTC or ethanol vehicle injection (Figure 4B). IL-6 was significantly elevated in the DRG of animals pretreated with DBTC ($p < 0.05$) compared with levels detected in naïve or ethanol vehicle pretreated rats. There was no significant difference in the IL-6 levels between ethanol vehicle control and naïve rats.

IL-6 potentiates capsaicin-evoked CGRP release from DRG neurons

As elevated levels of IL-6 in thoracic DRG might correlate with abdominal hypersensitivity after DBTC pretreatment, the possible effect of IL-6 on these cells was examined using cultured DRG neurons from adult rats by measuring the effect of IL-6 on capsaicin-evoked CGRP release. Stimulation of DRG cultures with capsaicin (30 nM) induced an increase in CGRP release that was further potentiated by IL-6 (20 ng/mL). The potentiating effect of IL-6 was blocked by TB-2-081 (5 µM), which had no effect on capsaicin evoked CGRP release (Figure 5A). IL-6 alone, TB-2-081 alone, or their combination did not activate CGRP release in the absence of capsaicin (Figure 5A).

As previous studies have proposed that exogenous soluble IL-6R (sIL-6R) administration is required to induce IL-6 competency [8;26], we tested whether sIL-6R administration would

further augment IL-6 effects on capsaicin-evoked CGRP release. Cultured DRG cells were incubated with IL-6 (20 ng/mL), sIL-6R (25 ng/mL), or both, for 30 min followed by 15 min incubation with capsaicin (30 nM). Neither IL-6 nor sIL-6R or the combination of both stimulated CGRP release (Figure 5B). IL-6, but not sIL-6R, potentiated capsaicin-evoked CGRP release (Figure 5B). Co-administration of IL-6 and sIL-6R did not further augment evoked CGRP release compared with IL-6 alone (Figure 5B). These experiments suggest that exogenous soluble IL-6R is not required for IL-6 augmentation of capsaicin-evoked CGRP release under these conditions.

Discussion

In the present study we used a novel, small molecule IL-6R antagonist TB-2-081 [8;15] to assess the role of IL-6 in pancreatitis-induced pain. TB-2-081 suppressed IL-6-induced mRNA expression of the hepatic acute phase protein α 1-antichymotrypsin (AACT) in human hepatoma HepG2[17] cells and inhibited IL-6 induced phosphorylation of STAT3, thereby antagonizing IL-6 signaling via its canonical signaling pathway [17;26]. Our data show that TB-2-081 displaces the binding of IL-6 from sIL-6R with apparent high affinity and inhibits IL-6 induced growth of TF-1 cells, which further support TB-2-081 as a natural product that exhibits characteristics of antagonist for IL-6R. The major finding of this study is that TB-2-081 effectively reverses pancreatitis-induced referred abdominal hypersensitivity; its action as an IL-6R antagonist further implies that IL-6R blockade, or IL-6 sequestration, may be therapeutic avenues for treatment of pancreatitis pain in humans. Furthermore, TB-2-081 was shown to be orally active.

We found that systemic or oral, but not intrathecal administration of TB-2-081 reversed pancreatitis-induced referred abdominal hypersensitivity, suggesting that this compound may have a peripheral mechanism of action. These data also suggest that peripheral IL-6R is likely to be critical in maintaining pancreatitis-induced hypersensitivity. This observation may be important in the development of novel therapeutics for pancreatitis pain. Furthermore, the rapid onset of reversal of referred abdominal hypersensitivity by TB-2-081 is also consistent with a direct interaction with IL-6R and disruption of a signaling pathway that modulates sensory input. Previous evidence suggests that the IL-6-IL-6R-gp130 complex stimulates MAPK activation and several MAPKs such as p38 and ERK. Activation of these pathways promotes the excitability of primary afferent neurons [16]; the relevance of these signaling pathways by IL-6R in pancreatitis induced pain states warrant further investigations.

Our findings on the alteration of IL-6 expression in the pancreas and DRG in the experimental model of pancreatitis also points to a possible role of IL-6 in the maintenance of the chronic inflammatory pain states. Both DBTC and ethanol vehicle caused a significant increase in IL-6 levels in the pancreas, compared to a very low expression of IL-6 in pancreata of naïve animals. However, as ethanol vehicle treated animals do not develop abdominal hypersensitivity or pancreatitis, it seems unlikely that pancreas-derived IL-6 alone is sufficient to promote the chronic pain state. One hypothesis that may explain the effects of alcohol on the pancreas is that the alcohol sensitizes the pancreas to the pathobiologic processes including inflammation, necrosis and fibrosis. Ethanol feeding alone, even at high doses, has minimal and inconsistent effects on morphologic findings in the pancreas in animals [20;38]. A 6-week, ethanol-containing diet increases the sensitivity of rats to acute pancreatitis induced by cholecystokinin (CCK)-8 [31]. An ethanol diet and CCK-8 infusion increases pancreatic NF κ B activation compared with injection of CCK-8 alone. Failure to produce experimental alcoholic pancreatitis and the fact that alcohol abuse causes pancreatic pathology in only a minority of patients suggests that ethanol may only

increase predisposition to pancreatitis. Therefore, elevated levels of IL-6 in vehicle treated group might be a necessary prerequisite for pancreatic injury induced by DBTC.

In contrast, we observed a large increase in IL-6 in the thoracic DRG after DBTC treatment but not in the ethanol vehicle treated controls. While the mechanisms underlying this increase in IL-6 expression in the DRG are presently unclear, IL-6 has been shown to be upregulated in injured nerves [1;2] and studies from IL-6 knockout mice indicate that the effects of IL-6 are critical for full expression of neuropathic pain in these models [24]. Thus, the upregulation of IL-6 in the thoracic DRG may be an autocrine or paracrine mechanism in the sensitization of primary afferents through activation of IL-6R in the peripheral terminals. Therefore, our data indicate that a possible mechanism of action for TB-2-081 in the DBTC model is an abrogation of IL-6 effects within the DRG of animals with chronic pancreatitis leading to a decrement in abdominal hypersensitivity. To determine if IL-6 promotes the excitability of primary afferents, particularly of nociceptors, the release of the excitatory neuropeptide, CGRP, from isolated DRG was used as a functional measure of the activation of sensory neurons in the absence or presence of capsaicin, a ligand for the TRPV1 channel [4;37]. Previous evidence suggests that IL-6 enhances the electrophysiological properties of TRPV1 [28]. IL-6 does not directly activate CGRP release, at least at the dose tested here, but significantly enhanced capsaicin-evoked CGRP release; the enhancing effect of IL-6 was blocked by TB-2-081. These data are consistent with a pronociceptive role of IL-6, and furthermore, suggest that the effect of IL-6 may be mediated by IL-6R present on the primary afferent terminals, where the signaling process through IL-6R activation modulates the function of TRPV1. Previous behavioral and electrophysiological studies suggested that exogenously applied soluble IL-6R was required for IL-6 mediated sensitization of TRPV1 currents or TRPV1-dependent thermal hyperalgesia [29;42]. However, we found that the enhanced release of CGRP by IL-6 does not require sIL-6R under our experimental conditions.

Thus, in the DBTC model of pancreatitis, the upregulation of IL-6 and the blockade of IL-6R by TB-2-081 implicate a critical role for IL-6 in inflammatory hyperalgesia. The mechanisms by which IL-6R activation potentiates the probable excitability of nociceptors remain to be elucidated. Previous work has suggested that TRPV1 plays a critical role in the etiology of acute and chronic pancreatitis [21;43]. It would be of interest to ascertain if potentiation of TRPV1 constitutes a critical mechanism for the pronociceptive effect of IL-6/IL-6R.

It is possible that TB-2-081 may block the effects of other members of IL-6 family of cytokines, including IL-11 and oncostatin M [17]. However these members of the IL-6 family of cytokines have been suggested to be involved in hematogenesis and oncogenesis and their role in nociception is not clearly established. Additionally, while oncostatin M also plays a role in the development of small TRPV1⁺ neurons, this is not likely to be relevant to our study [23]. Further studies are necessary to determine the antagonistic effects of TB-2-081 on IL-6 type cytokines *in vivo*.

A variety of studies have demonstrated that overproduction of IL-6 contributes to the pathogenesis of various autoimmune and inflammatory diseases, including rheumatoid arthritis (RA), systemic-onset juvenile chronic arthritis (JCA), degenerative lumbar spinal disorder. A large number of diseases and pathologic conditions in which IL-6 plays a key role, and existing clinical validation of mechanism, make the development of molecules which block the actions of IL-6 desirable targets for drug discovery and development.

Therapeutics that target IL-6R are currently limited to monoclonal antibodies. Blocking IL-6 actions by use of a humanized antibody, tocilizumab, has been shown to be therapeutically

effective for rheumatoid arthritis [22;35;36], systemic juvenile idiopathic arthritis and Crohn's disease [25]. While these are exciting developments, there remain some contraindications regarding the use of monoclonal antibodies in risks of infections and immunogenicity [19]. The promise of a small molecule, orally available IL-6R antagonist such as TB-2-081 and/or its derivatives, offers a viable alternative strategy to the development of therapeutics targeting the pathological actions of IL-6.

Acknowledgments

The research of the Drug Design and Synthesis Section, CBRB, NIDA & NIAAA, was supported by the NIH Intramural Research Programs of the National Institute on Drug Abuse (NIDA) and the National Institute on Alcohol Abuse and Alcoholism. We thank NIDA for support of the X-ray crystallographic studies (NIDA contract Y1-DA6002).

Abbreviations

CGRP	calcitonin gene related peptide
DBTC	dibutyltin dichloride
DRG	dorsal root ganglia
IL-6R	interleukin 6 receptor
TB-2-081	3- <i>O</i> -formyl-20 <i>R</i> ,21-epoxyresibufogenin

Reference List

- [1]. Arruda JL, Colburn RW, Rickman AJ, Rutkowski MD, DeLeo JA. Increase of interleukin-6 mRNA in the spinal cord following peripheral nerve injury in the rat: potential role of IL-6 in neuropathic pain. *Brain Res Mol Brain Res*. 1998; 62:228–235. [PubMed: 9813345]
- [2]. Arruda JL, Sweitzer S, Rutkowski MD, DeLeo JA. Intrathecal anti-IL-6 antibody and IgG attenuates peripheral nerve injury-induced mechanical allodynia in the rat: possible immune modulation in neuropathic pain. *Brain Res*. 2000; 879:216–225. [PubMed: 11011025]
- [3]. Ballantyne JC, LaForge KS. Opioid dependence and addiction during opioid treatment of chronic pain. *Pain*. 2007; 129:235–255. [PubMed: 17482363]
- [4]. Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, Julius D. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature*. 1997; 389:816–824. [PubMed: 9349813]
- [5]. Chaudhry MA, Bowen BD, Eaves CJ, Piret JM. Empirical models of the proliferative response of cytokine-dependent hematopoietic cell lines. *Biotechnol Bioeng*. 2004; 88:348–358. [PubMed: 15486941]
- [6]. Dina OA, Green PG, Levine JD. Role of interleukin-6 in chronic muscle hyperalgesic priming. *Neuroscience*. 2008; 152:521–525. [PubMed: 18280048]
- [7]. Enomoto A, Rho MC, Fukami A, Hiraku O, Komiyama K, Hayashi M. Suppression of cancer cachexia by 20*S*,21-epoxy-resibufogenin-3-acetate-a novel nonpeptide IL-6 receptor antagonist. *Biochem Biophys Res Commun*. 2004; 323:1096–1102. [PubMed: 15381112]
- [8]. Enomoto A, Rho MC, Komiyama K, Hayashi M. Inhibitory effects of bufadienolides on interleukin-6 in MH-60 cells. *J Nat Prod*. 2004; 67:2070–2072. [PubMed: 15620253]
- [9]. Gachago C, Draganov PV. Pain management in chronic pancreatitis. *World J Gastroenterol*. 2008; 14:3137–3148. [PubMed: 18506917]
- [10]. Graham, DD.; Bonica, JJ. Painful diseases of the liver, biliary system and pancreas. In: Loeser, JD.; Butler, SH.; Chapman, R.; Turk, DC., editors. *Bonica's Management of Pain*. Lippincott Williams&Williman; 2001. p. 1293-1308.

- [11]. Heinrich PC, Behrmann I, Muller-Newen G, Schaper F, Graeve L. Interleukin-6-type cytokine signalling through the gp130/Jak/STAT pathway. *Biochem J.* 1998; 334(Pt 2):297–314. [PubMed: 9716487]
- [12]. Hibi M, Murakami M, Saito M, Hirano T, Taga T, Kishimoto T. Molecular cloning and expression of an IL-6 signal transducer, gp130. *Cell.* 1990; 63:1149–1157. [PubMed: 2261637]
- [13]. Inagaki T, Hoshino M, Hayakawa T, Ohara H, Yamada T, Yamada H, Iida M, Nakazawa T, Ogasawara T, Uchida A, Hasegawa C, Miyaji M, Takeuchi T. Interleukin-6 is a useful marker for early prediction of the severity of acute pancreatitis. *Pancreas.* 1997; 14:1–8. [PubMed: 8981500]
- [14]. Ishihara K, Hirano T. IL-6 in autoimmune disease and chronic inflammatory proliferative disease. *Cytokine Growth Factor Rev.* 2002; 13:357–368. [PubMed: 12220549]
- [15]. Kamano Y, Nogawa T, Yamashita A, Hayashi M, Inoue M, Drasar P, Pettit GR. Isolation and structure of a 20,21-epoxybufenolide series from “Ch’an Su”. *J Nat Prod.* 2002; 65:1001–1005. [PubMed: 12141860]
- [16]. Kang KW, Wagley Y, Kim HW, Pokharel YR, Chung YY, Chang IY, Kim JJ, Moon JS, Kim YK, Nah SY, Kang HS, Oh JW. Novel role of IL-6/SIL-6R signaling in the expression of inducible nitric oxide synthase (iNOS) in murine B16, metastatic melanoma clone F10.9, cells. *Free Radic Biol Med.* 2007; 42:215–227. [PubMed: 17189827]
- [17]. Kino T, Boos TL, Sulima A, Siegel EM, Gold PW, Rice KC, Chrousos GP. 3-O-Formyl-20R,21-epoxyresibufogenin suppresses IL-6-type cytokine actions by targeting the glycoprotein 130 subunit: potential clinical implications. *J Allergy Clin Immunol.* 2007; 120:437–444. [PubMed: 17451794]
- [18]. Kishimoto T. The biology of interleukin-6. *Blood.* 1989; 74:1–10. [PubMed: 2473791]
- [19]. Kong JS, Teuber SS, Gershwin ME. Potential adverse events with biologic response modifiers. *Autoimmun Rev.* 2006; 5:471–485. [PubMed: 16920574]
- [20]. Kono H, Nakagami M, Rusyn I, Connor HD, Stefanovic B, Brenner DA, Mason RP, Arteel GE, Thurman RG. Development of an animal model of chronic alcohol-induced pancreatitis in the rat. *Am J Physiol Gastrointest Liver Physiol.* 2001; 280:G1178–G1186. [PubMed: 11352811]
- [21]. Liddle RA. The role of Transient Receptor Potential Vanilloid 1 (TRPV1) channels in pancreatitis. *Biochim Biophys Acta.* 2007; 1772:869–878. [PubMed: 17428642]
- [22]. Maini RN, Taylor PC, Szechinski J, Pavelka K, Broll J, Balint G, Emery P, Raemen F, Petersen J, Smolen J, Thomson D, Kishimoto T. Double-blind randomized controlled clinical trial of the interleukin-6 receptor antagonist, tocilizumab, in European patients with rheumatoid arthritis who had an incomplete response to methotrexate. *Arthritis Rheum.* 2006; 54:2817–2829. [PubMed: 16947782]
- [23]. Morikawa Y, Tamura S, Minehata K, Donovan PJ, Miyajima A, Senba E. Essential function of oncostatin m in nociceptive neurons of dorsal root ganglia. *J Neurosci.* 2004; 24:1941–1947. [PubMed: 14985435]
- [24]. Murphy PG, Ramer MS, Borthwick L, Gauldie J, Richardson PM, Bisby MA. Endogenous interleukin-6 contributes to hypersensitivity to cutaneous stimuli and changes in neuropeptides associated with chronic nerve constriction in mice. *Eur J Neurosci.* 1999; 11:2243–2253. [PubMed: 10383613]
- [25]. Nishimoto N, Kishimoto T. Interleukin 6: from bench to bedside. *Nat Clin Pract Rheumatol.* 2006; 2:619–626. [PubMed: 17075601]
- [26]. Nogawa T, Kamano Y, Yamashita A, Pettit GR. Isolation and structure of five new cancer cell growth inhibitory bufadienolides from the Chinese traditional drug Ch’an Su. *J Nat Prod.* 2001; 64:1148–1152. [PubMed: 11575946]
- [27]. Noh KW, Pungpapong S, Wallace MB, Woodward TA, Raimondo M. Do cytokine concentrations in pancreatic juice predict the presence of pancreatic diseases? *Clin Gastroenterol Hepatol.* 2006; 4:782–789. [PubMed: 16713745]
- [28]. Obreja O, Biasio W, Andratsch M, Lips KS, Rathee PK, Ludwig A, Rose-John S, Kress M. Fast modulation of heat-activated ionic current by proinflammatory interleukin 6 in rat sensory neurons. *Brain.* 2005; 128:1634–1641. [PubMed: 15817518]

- [29]. Obreja O, Schmelz M, Poole S, Kress M. Interleukin-6 in combination with its soluble IL-6 receptor sensitises rat skin nociceptors to heat, in vivo. *Pain*. 2002; 96:57–62. [PubMed: 11932061]
- [30]. Oka T, Oka K, Hosoi M, Hori T. Intracerebroventricular injection of interleukin-6 induces thermal hyperalgesia in rats. *Brain Res*. 1995; 692:123–128. [PubMed: 8548295]
- [31]. Pandol SJ, Periskic S, Gukovsky I, Zaninovic V, Jung Y, Zong Y, Solomon TE, Gukovskaya AS, Tsukamoto H. Ethanol diet increases the sensitivity of rats to pancreatitis induced by cholecystokinin octapeptide. *Gastroenterology*. 1999; 117:706–716. [PubMed: 10464148]
- [32]. Reyes-Gibby CC, Shete S, Yennurajalingam S, Frazier M, Bruera E, Kurzrock R, Crane CH, Abbruzzese J, Evans D, Spitz MR. Genetic and nongenetic covariates of pain severity in patients with adenocarcinoma of the pancreas: assessing the influence of cytokine genes. *J Pain Symptom Manage*. 2009; 38:894–902. [PubMed: 19692203]
- [33]. Rose-John S, Scheller J, Elson G, Jones SA. Interleukin-6 biology is coordinated by membrane-bound and soluble receptors: role in inflammation and cancer. *J Leukoc Biol*. 2006; 80:227–236. [PubMed: 16707558]
- [34]. Sarr MG, Sakorafas GH. Incapacitating pain of chronic pancreatitis: a surgical perspective of what is known and what needs to be known. *Gastrointest Endosc*. 1999; 49:S85–S89. [PubMed: 10049457]
- [35]. Smolen JS, Beaulieu A, Rubbert-Roth A, Ramos-Remus C, Rovinsky J, Alecock E, Woodworth T, Alten R. Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomised trial. *Lancet*. 2008; 371:987–997. [PubMed: 18358926]
- [36]. Strand V, Singh JA. Newer biological agents in rheumatoid arthritis: impact on health-related quality of life and productivity. *Drugs*. 2010; 70:121–145. [PubMed: 20108988]
- [37]. Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, Skinner K, Raumann BE, Basbaum AI, Julius D. The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron*. 1998; 21:531–543. [PubMed: 9768840]
- [38]. Tsukamoto H, Towner SJ, Yu GS, French SW. Potentiation of ethanol-induced pancreatic injury by dietary fat. Induction of chronic pancreatitis by alcohol in rats. *Am J Pathol*. 1988; 131:246–257. [PubMed: 3358454]
- [39]. Vera-Portocarrero LP, Lu Y, Westlund KN. Nociception in persistent pancreatitis in rats: effects of morphine and neuropeptide alterations. *Anesthesiology*. 2003; 98:474–484. [PubMed: 12552208]
- [40]. Vera-Portocarrero LP, Xie JY, Kowal J, Ossipov MH, King T, Porreca F. Descending facilitation from the rostral ventromedial medulla maintains visceral pain in rats with experimental pancreatitis. *Gastroenterology*. 2006; 130:2155–2164. [PubMed: 16762636]
- [41]. von Banchet GS, Kiehl M, Schaible HG. Acute and long-term effects of IL-6 on cultured dorsal root ganglion neurones from adult rat. *J Neurochem*. 2005; 94:238–248. [PubMed: 15953366]
- [42]. Wick EC, Hoge SG, Grahn SW, Kim E, Divino LA, Grady EF, Bunnett NW, Kirkwood KS. Transient receptor potential vanilloid 1, calcitonin gene-related peptide, and substance P mediate nociception in acute pancreatitis. *Am J Physiol Gastrointest Liver Physiol*. 2006; 290:G959–G969. [PubMed: 16399878]
- [43]. Xu GY, Winston JH, Shenoy M, Yin H, Pendyala S, Pasricha PJ. Transient receptor potential vanilloid 1 mediates hyperalgesia and is up-regulated in rats with chronic pancreatitis. *Gastroenterology*. 2007; 133:1282–1292. [PubMed: 17698068]
- [44]. Yaksh TL, Rudy TA. Chronic catheterization of the spinal subarachnoid space. *Physiol Behav*. 1976; 17:1031–1036. [PubMed: 14677603]

Figure 1.

A. Structure of TB-2-081. **B** Saturation of IL-6 at human recombinant sIL-6R. Half saturation concentration is 960 pM. **C.** Competitive binding of TB-2-081 to IL-6R ($IC_{50} = 29.4$ pM). **D.** TB-2-081 inhibits the growth of IL-6 dependent cell line TF-1 ($IC_{50} = 250$ pM).

Figure 2. Effects of systemic TB-2-081 in the DBTC pancreatitis model. **A.** Time course of the effects of TB-2-081 in rats with pancreatitis (DBTC) or without pancreatitis (ethanol vehicle). The doses of TB-2-081 are represented as mg/kg. PBS/ethanol indicates the vehicle for TB-2-081. N=6-10 animals in all groups. (* $P < 0.05$). **B.** Dose-response curve for TB-2-081 induced reversal of abdominal hypersensitivity 15 min after subcutaneous (s.c.) administration in animals with experimental pancreatitis. **C.** Area under the time-effect curve for subcutaneously administered TB-2-081 against referred abdominal hypersensitivity in rats injected with DBTC. N=6-10 animals in all groups. *indicates significant difference from the control group (ethanol vehicle) (* $P < 0.01$).

Figure 3.

Effects of orally administered TB-2-081 in the DBTC pancreatitis model. **A.** Time course of the effects of TB-2-081 in rats with pancreatitis (DBTC) or without pancreatitis (ethanol vehicle). The doses of TB-2-081 are represented as mg/kg. PBS/ethanol indicates the vehicle for TB-2-081. N=5-9 animals per group. **B.** Area under the time-effect curve for orally administered TB-2-081 against referred abdominal hypersensitivity in rats injected with DBTC. *indicates significant difference from the control group (ethanol vehicle). (*P < 0.01).

Figure 4.

Pancreatitis-induced changes in IL-6 expression in the pancreas and DRG. **A.** Time-course of changes in levels of IL-6 in pancreata of naïve, ethanol vehicle or DBTC-treated animals (*indicates significant difference from naïve group, $P < 0.01$). **B.** The levels of IL-6 in the T₈-T₁₂ DRG obtained from naïve animals or animals treated with ethanol vehicle or with DBTC-induced visceral inflammation at 6 day after injection are shown (* $P < 0.01$, N= 6 rats per group in all panels).

Figure 5.

Effects of IL-6 on capsaicin-evoked CGRP release from adult DRG neurons in culture. **A.** Adult cultured DRGs were treated with capsaicin (30 nM) alone (i.e.; Control), TB-2-081 (5 μ M) followed by capsaicin (TB-2-081), IL-6 (20 ng/ml) followed by capsaicin (IL-6), or IL-6 with TB-2-081 followed by capsaicin (IL-6+TB-2-081). Treatment with IL-6 increased the levels of capsaicin-evoked CGRP release by 30 % (*P = 0.01). **B.** Application of IL-6, soluble IL-6R (20 ng/ml) or a combination of both on the basal release of CGRP (*P = 0.02, n=6 per group in all panels).