CORRESPONDENCE

Hysterectomy in Germany: A DRG-Based Nationwide Analysis, 2005–2006

by Prof. Dr. med. Andreas Stang MD, Prof. Dr. Ray M. Merrill, PhD, PD Dr. sc. hum. Oliver Kuss, PhD in volume 30/2011

3. www.ClinicalTrials.gov Identifier: NCT00839722: Fertility after UAE for the Treatment of Leiomyomas. Last accessed: 21 September 2009
4. Stang A, Merrill RM, Kuss O: Hysterectomy in Germany: a DRG-based nationwide analysis, 2005–2006. *Dtsch Arztebl Int* 2011; 108(30): 508–14.

Dr. med. Gernot Rott
Klinik für diagnostische und interventionelle Radiologie
BETHESDA-Krankenhaus Duisburg
info@myom-therapie.de

Conflict of interest statement

Dr Rott has received conference delegate fees, travel expenses, and hotel expenses from Terumo.

Hysterectomy Is Mostly Unnecessary

As a radiologist who has been performing uterine artery embolizations for more than 10 years I have been receiving inquiries from women with asymptomatic fibroids for exactly the same period of time. The women want to know whether the hysterectomy their gynecologist in private practice recommended—for which they usually already have an inpatient appointment—might be avoidable if they underwent uterine artery embolization for their fibroids. As a rule, these women obviously require neither a hysterectomy nor other surgery nor uterine artery embolization nor any other treatment.

The number of unnecessary hysterectomies in Germany, which are performed without any indication, is likely to be shockingly large. Women only find out about the uterine artery embolization option through their own internet searches.

The mention of interventional radiological procedures, such as uterine artery embolization and high intensity focused ultrasound (HIFUS; MRI-guided focused ultrasonographic ablation, to use the authors' term), in the article is therefore particularly welcome.

Whether uterine artery embolization negatively affects fertility is still the subject of controversial discussion. The study reported by Mara (1), which is usually cited in this context, favors myomectomy over uterine artery embolization in this regard, while pointing out limitations and weaknesses. The results of other studies (unfortunately of a lower evidence level) point into the opposite direction (2) or remain to be evaluated (3).

Quite apart from all these facts, whether uterine artery embolization has a questionable negative effect on fertility should not be used as an argument against the procedure in an article about hysterectomy, after which fertility is not an issue anyway.

DOI: 10.3238/arztebl.2012.0158a

REFERENCES

1. Mara M, Maskova J, Fucikova Z, Kuzel D, Belsan T: Midterm clinical and first reproductive results of a randomized controlled trial comparing uterine fibroid embolization and myomectomy. *Cardiovasc Intervent Radiol* 2008; 31: 73–85.
2. Pisco JM, Duarte M, Bilhim T, Cirurgiao F, Oliveira A: Pregnancy after uterine fibroid embolization. *Fertility and Sterility* 2011; 95: 1121.e5–8.

Data Not Correctly Calculated

Stang et al have provided an important impulse for further discussion with their much-welcomed article on hysterectomy in Germany. All crucial statements in the article relate to hysterectomy undertaken for benign indications. The data on hysterectomy given in the first sentences of the results and discussion sections are contradictory: the hysterectomy rate in benign diseases of the female genital organs in Germany was not 3.6 per 1000 person-years; this is obviously the total hysterectomy rate for the whole of Germany. A more detailed explanation of the hysterectomy rate per 100 000 person-years, which was mentioned several times, would have been desirable. One critical comment: the authors obviously used the entire female population on their selected date of 2005 in their calculations. However, it does seem sensible to exclude at least girls younger than 10 years and women older than 80 years. The statement relating to the use of MRI-guided focused ultrasonographic ablation and uterine artery embolization for women wishing to keep their uterus requires a correction: these treatments for fibroids have been routinely used for many years. An alternative to hysterectomy in menorrhagia is hysteroscopy, a minimally invasive procedure, which was not mentioned in the discussion section (2).

The hysterectomy rate, but also the way in which hysterectomies are performed in benign indications, gives a clue about the respective surgeon's school of thought, the apparatus and other options available in the respective hospital, and flows—sometimes extending beyond state boundaries.

A self-critical look at the hysterectomy rate in Germany on the part of the gynecological community is urgently required. This relates to defining indications in case of benign disease, the surgical *modus operandi*, and the question of whether the cervix should be spared. In conclusion, a large, nationwide, prospective health services research study is more than overdue in Germany.

DOI: 10.3238/arztebl.2012.0158b

REFERENCES

1. Stang A, Merrill RM, Kuss O: Hysterectomy in Germany: a DRG-based nationwide analysis, 2005–2006. *Dtsch Arztebl Int* 2011; 108(30): 508–14.
2. Stovall DW: Alternatives to hysterectomy: focus on global endometrial ablation, uterine fibroid embolization, and magnetic resonance-guided focused ultrasound. *Menopause* 2011; 18: 443–50.

Prof. Dr. med. Matthias David
Charité – Universitätsmedizin Berlin
Campus Virchow-Klinikum
matthias.david@charite.de

Prof. Dr. med. Heribert Kantenich
Fertility Center Berlin

Conflict of interest statement

The authors declare that no conflict of interest exists.

Outpatient Hysterectomies Were Not Included

Sometimes I wonder just what the point of certain statistical analyses might be. In the above article, the authors present total hysterectomy rates and the different approaches to the procedure in Germany.

The authors based their article on data from nationwide diagnosis-related groups (DRG) from 2005 and 2006. But these data reflect only the services provided by hospitals. The authors know this because they say that outpatient hysterectomies can be ignored as “day-surgery hysterectomy is rare in Germany.” In this context they cite a study from 2007. However, the very study reported by Salfelder et al. says that the rates of day-surgery hysterectomies are rising. In the „Tagesklinik Altonaer Straße“, a Hamburg-based day clinic, some 150 outpatient hysterectomies were performed in 2005 and 200 in 2006. The study reported by Salfelder et al. also points out the day surgery centers of the Velen Working Group for Outpatient Surgery, which also deliver hysterectomy surgery on an outpatient basis.

Add to this the multitude of gynecological practices that conducted considerable numbers of hysterectomies on an outpatient basis as early as in 2005 and 2006.

In our institution alone, 40 outpatient hysterectomies were performed in 2005 and 50 in 2006.

According to statistical data, 9222 hysterectomies were performed in 2005 in the federal state of Saxony. Bearing in mind that Saxony has several day surgery centers that also perform hysterectomies on an outpatient basis, then it is highly questionable whether the statistical analysis of the number of hysterectomies performed in Germany that is presented in the article is actually in any way meaningful.

Furthermore, I think that the negative consequences of bilateral ovariectomy that were mentioned in the conclusion fall outside the scope of the article.

DOI: 10.3238/arztebl.2012.0159a

REFERENCES

1. Salfelder A, Lueken RP, Gallinat A, et al.: Hysterektomie als Standard-eingriff in der Tagesklinik – ein Wagnis? *Frauenarzt* 2007; 48: 954–8.

2. Stang A, Merrill RM, Kuss O: Hysterectomy in Germany: a DRG-based nationwide analysis, 2005–2006. *Dtsch Arztebl Int* 2011; 108(30): 508–14.

Dr. med. Jost Klippel
Frauenarzt/Ambulante Operationen
Hohe Straße 21
02829 Holtendorf

Conflict of interest statement

Dr Klippel has received conference delegate fees, travel expenses, and hotel expenses from Takeda.

In Reply:

David and Kantenich are correct in their criticism of the fact that the hysterectomy rates reported in the first sentences of the results and discussion sections (1) are contradictory. The reason for this is as follows: in the first sentence of the results section, we reported the hysterectomy rates for any indication (362/100 000 person-years) for the entire age range. In the first sentence of the discussion section, we reported international comparisons of hysterectomy rates for benign diseases of the female genital organs (Sweden, US, Australia). These rates were reported for women aged 20 years or older (2).

David and Kantenich further criticize that we included girls younger than 10 years and women older than 80 years in our calculations. This criticism is not justified. Excluding persons from rate calculations makes sense only if persons are not “at risk” for hysterectomy. In 2005–2006 in Germany, hysterectomies were performed in eight girls younger than 10 and 8180 women older than 80 years (2.7% of all hysterectomies).

David and Kantenich asked for a correction relating to our mention of focused ultrasonographic ablation and uterine artery embolization. This statement related to a recommendation of an S2 guideline for the diagnostic evaluation and therapy of endometriosis (3), which we clarified by citing a reference. David and Kantenich are thus asking for a discussion of the S2 guideline, which obviously cannot take place here.

Klippel mentions the study of Salfelder et al. (4), which we cited in our article, and notes that hysterectomies performed outside the inpatient setting should not be ignored. According to Salfelder et al, a total of 193 laparoscopic hysterectomies were performed in the „Tagesklinik Altonaer Straße“ in 2006. According to DRG data, a total of 2335 hysterectomies were performed among Hamburg’s female population in 2006. If we assumed (which is unlikely) that the patients operated on by Salfelder et al. were Hamburg residents, then the DRG data would include 92% of hysterectomies performed in this population. Since it may safely be assumed that only some of the patients operated on by Salfelder et al were Hamburg residents, we can assume that the proportion is higher than 92%.

Differences in DRG based hysterectomy rates for benign diseases of the female genital organs that are specific to federal states (Hamburg: 213.8;

Mecklenburg-Western Pomerania: 361.9 per 100 000 person-years) can obviously not be “debated away” by using the argument of differently high hysterectomy rates outside inpatient settings. Why Klippel has to cast considerable doubt on our findings is not clear, especially when remembering that no better hysterectomy data for Germany are currently available. We have nothing to add to the comments made by Rott.

DOI: 10.3238/arztebl.2012.0159b

REFERENCES

1. Stang A, Merrill RM, Kuss O: Hysterectomy in Germany: a DRG-based nationwide analysis, 2005–2006. *Dtsch Arztebl Int* 2011; 108(30): 508–14.
2. Stang A, Merrill RM, Kuss O: Nationwide rates of conversion from laparoscopic or vaginal hysterectomy to open abdominal hysterectomy in Germany. *Eur J Epidemiol* 2011; 26: 125–33.
3. Deutsche Gesellschaft für Gynäkologie und Geburtshilfe (DGGG), Schweizerische Gesellschaft für Gynäkologie und Geburtshilfe (SGGG), Österreichische Gesellschaft für Gynäkologie und Geburtshilfe e. V. (OEGGG), Stiftung Endometriose-Forschung (SEF), Deutsche Gesellschaft für Allgemein- und Viszeralchirurgie e. V., Deutsche Gesellschaft für Urologie e. V., Deutsche Gesellschaft für Gynäkologische Endokrinologie und Fortpflanzungsmedizin e. V., Arbeitsge-

meinschaft für Gynäkologische Onkologie (AGO) e. V. der DGGG, Endometriose-Vereinigung Deutschland e. V., Österreichische Endometriose Vereinigung e. V. Diagnostik und Therapie der Endometriose: AWMF-Leitlinien-Register Nr. 015/045. Stand 1. 5. 2010. Link: www.awmf.org/uploads/tx_szleitlinien/015-045_S1_Diagnostik_und_Therapie_der_Endometriose_05-2010_05-2015.pdf. Last accessed on 2 May 2011.

4. Salfelder A, Lueken RP, Gallinat A, et al.: Hysterektomie als Standard-eingriff in der Tagesklinik – ein Wagnis? *Frauenarzt* 2007; 48: 954–8.

Prof. Dr. med. Andreas Stang

Institut für Klinische Epidemiologie
Medizinische Fakultät
Martin-Luther-Universität Halle-Wittenberg, Halle (Saale)
andreas.stang@medizin.uni-halle.de

Prof. Dr. Ray M. Merrill, PhD

Department of Health Science, College of Life Sciences,
Brigham Young University, Provo, UT 84602, U.S.A.

PD Dr. sc. hum. Oliver Kuss

Institut für Medizinische Epidemiologie, Biometrie und Informatik
Universitätsklinikum Halle und Medizinische Fakultät
Martin-Luther-Universität Halle-Wittenberg, Halle (Saale)

Conflict of interest statement

The authors declare that no conflict of interest exists.