Institute of Microbiology, Vídeňská 1083, 142 20 Prague 4-Krč, Czech Republic

EMBL accession no. X69872

4. Mandel, M. and Marmur, J. (1968) In Grossman, L. and Moldave, K. (eds), *Methods in Enzymology*. Academic Press, New York, London. Vol. **12B**, pp. 195–206.
5. Bolotin, A. P., Sorokin, A. V., Aleksandrov, N. N., Danilenko, V. N. and Kozlov, Y. I. (1985) *Dokl. Akad. Nauk USSR* **283**, 1014–1017.
6. Kendall, K. J. and Cohen, S. N. (1988) *J. Bacteriol.* **170**, 4634–4651.
7. Wu, X. and Rov, K. (1993) *J. Bacteriol.* **175**, 37–52.

After the circular plasmids pBS24.2 from *S. cyanogenes* (5) and pIJ101 from *S. lividans* (6), and the linear plasmid pSCL1 from *S. clavuligerus* (7), pSLG33 is another completely sequenced native plasmid from the genus *Streptomyces*. Analysis of all up to currently accumulated sequence data and their comprehensive comparison should bring new informations for deeper and general understanding of streptomycetes plasmid composition and plasmid replication.

We are indebted to P. Formánek from the Institute of Molecular Genetics, Prague for the determinations of T_m 's.

1. Tichy, P., Moskalenko, L. N., Smrčková, I. and Krumphanzl, V. (1985) *FEMS Microbiol. Lett.* **27**, 65–68.
2. Petříček, M. and Tichy, P. (1989) *FEMS Microbiol. Lett.* **61**, 109–113.
3. Locci, R. (1989) In Williams, S. T., Sharpe, M. E. and Holt, J. G. (eds), *Bergey's Manual of Systematic Bacteriology*. Williams and Wilkins, Baltimore, MD. Vol. 4, pp. 2451–2508.

[illegible]

Figure 1. Nucleotide sequence of the mini-plasmid pSLG33 from *Streptomyces lavendulae-grasseri* RIA746, linearized with the endonuclease *Pst*I.

* To whom correspondence should be addressed