Published in final edited form as:

J Am Chem Soc. 2012 March 28; 134(12): 5496–5499. doi:10.1021/ja3004733.

Asymmetric Hydrovinylation of Vinylindoles. A Facile Route to Cyclo-penta[g]indole Natural Products (+)-*cis*-Triketrin A and (+)-*cis*-Triketrin B

Wang Liu, Hwan Jung Lim⁺, and T. V. RajanBabu^{*}

Department of Chemistry, The Ohio State University, 100 West 18th Avenue, Columbus, Ohio 43210, United States

Abstract

Vinylindoles undergo Ni(II)-catalyzed asymmetric hydrovinylation under very mild conditions (−78 °C, 1 atmosphere ethylene, 4 mol% catalyst) to give the corresponding 2-but-3-enyl derivatives in excellent yields and enantioselectivities. Hydroboration of the alkene, oxidation to an acid, followed by Friedel-Crafts annulation gives an indole-annulated cyclopentanone that is a suitable precursor for the syntheses of *cis*-trikettrins and all known herbindoles. For example, the cyclopentanone from 4-ethyl-7-vinylindole is converted into (+)-*cis*-trikettrins A in four steps (Wittig reaction, alkene isomerization, diastereoselective hydrogenation and nitrogen deprotection). The previous synthesis of this molecule from (*S*)-(-)-malic acid involved more than 20 steps and a preparative HPLC separation of diastereomeric intermediates.

Many biologically important indole derivatives are adorned with methyl-bearing secondary stereogenic centers at one or more of their peripheral carbons. Examples include the potent neuroprotective agent indole-3-propionic acid (**1**),¹ antibacterial (-)-indolmycin (**2**),² protein phosphatase inhibitors such as drag-macidin D (**3**),³ anti-bacterial trikettrins (**4**, **5**)^{4a} and cytotoxic herbindoles (**6**).^{4b}

While many efficient catalytic procedures have been developed for the stereoselective attachment of a chiral side-chain to the reactive C₃ position, there is a paucity of such methods if this chain is appended to any of the other carbons of the indole nucleus.⁵ Apart from notable addition of an indole-2-trifluoro(organo)borate to α,β-unsaturated aldehydes catalyzed by an organocatalyst,⁶ no general procedures for the installation of chiral appendages at other positions of indole have been reported. While considering a broadly applicable enantioselective approach to the cyclopenta[g]indole natural products, trikettrins, herb-indoles and related molecules,⁷ we wondered whether asymmetric hydrovinylation⁸ (Scheme 1) of an appropriately substituted vinyl-indole might provide an expeditious entry into this important class of compounds.

In this paper we report that Ni(II)-catalyzed asymmetric hydrovinylation of C₂, C₃, C₄, C₅, C₆ and C₇ vinylindoles proceed with surprising efficiency and excellent selectivity even when the indole nitrogen is unprotected. Details of this study along with applications of the resulting (2-but-3-enyl)indole derivatives for the syntheses of (+)-*cis*-trikettrins A (**4a**) and (+)-*cis*-trikettrins-B (**5a**) are documented here. Also reported is a novel application of a

rajanbabu.1@osu.edu.

⁺Current address: Korea Research Institute of Chemical Technology, 141 Gajeongro, Daejeon 305-600, Korea

Supporting Information Available: Full experimental details for the preparation of the intermediates and hydrovinylation reactions, spectroscopic and chromatographic data for characterization of all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>

recently disclosed⁹ [LPd(II)-H]⁺-catalyzed isomerization of a double bond to set up a highly diastereoselective reduction of a functionalized 1,3-dimethylcyclopent-1-ene.

Our initial studies directed towards the optimization of the reaction conditions for the asymmetric hydrovinylation of prototypical vinylindoles revealed that these compounds were among the most reactive substrates, with the reactions proceeding at -78°C under 1 atmosphere of ethylene, even when the nitrogen was left unprotected. Of the numerous ways of synthesizing indole derivatives, we resorted to the operationally simple, yet broadly applicable Bartoli method,¹⁰ which is illustrated for 7-vinylindole in Scheme 2. The asymmetric hydrovinylation of this substrate (Scheme 2) shows the subtle effects of the highly tunable phosphoramidite ligands (Figure 2)¹¹ on the selectivity of this reaction.¹² While each of the ligands **L1**–**L3** gave excellent yield (>97%) of the product, the *C*₁-symmetric ligand **L3** gave the best enantioselectivity (99% ee) as determined by chiral stationary phase GC. A racemic sample of the product was prepared using the ligand **L4**, or using racemic **L2**.

Results of asymmetric hydrovinylation of other vinylindoles are shown in Table 1.¹² In general, excellent yields and very good to excellent ee's were obtained for the hydrovinylation of a variety of substrates with as little as 4 mol% of the catalyst. For 2- and 7-vinylindoles (entries 1, 7 and 8), the ligand **L3** appears to be significantly better than **L1** and **L2**, whereas for the others, there is little perceptible difference. The enantioselectivity in the reaction of 5-vinylindole (entry 4) was determined after conversion of the terminal alkene into an alcohol and analyzing the Mosher ester derived from this compound by NMR.¹² The uncertainty in the ee measurements reflects the limits of this method rather than a less selective reaction. An *N*-Ts derivative (entry 6), which required a higher catalyst loading for comparable conversions, gave slightly lower ee's than the corresponding unprotected derivative (entry 5). Finally, the 7-vinyl derivatives shown in entries 7 and 8, gave excellent yields and selectivities of products that are potentially useful for enantioselective syntheses of trikentrin and herb-indoles.

The starting material for a short synthesis of (+)-*cis*-trikentrin A (Scheme 3) is the HV product **7a**, which is formed in 99% yield and 96% ee from 4-ethyl-7-vinylindole (entry 8, Table 1, using **L3**). The *N*-tosylated alkene **8a** was subjected to hydroboration followed by oxidation to prepare the carboxylic acid **10a**. An intramolecular Friedel-Crafts reaction, followed by a Wittig reaction gave the exo-methylene compound **12a** in ~70% yield from the acid **10a**. Apart from the asymmetric hydrovinylation, the major challenge in this synthesis turned out to be the *direct* stereoselective hydrogenation of **12a** to a trikentrin derivative. Repeated attempts under a variety of homogeneous and heterogeneous hydrogenation conditions¹⁵ led to a mixture of *cis*- and *trans*-trikentrin derivatives (both useful for syntheses of the corresponding natural products, albeit in a less selective fashion). A satisfactory solution to the problem involved the use of our recently published⁹ Pd(II)-catalyzed isomerization of the terminal **12a** into an internal alkene **13a**, followed by Crabtree hydrogenation,¹⁶ which gave exclusively the *N*-Ts derivative **14a** as a single diastereomer. Removal of the *N*-Ts derivative using TBAF in refluxing THF¹⁷ gave the natural product (+)-*cis*-trikentrin (**4a**) identical in all respects with the reported natural product.^{4a,14}

Synthesis of a more complex natural product, (+)-*cis*-trikentrin B, shown in Scheme 3 illustrates the use of the HV product, **7b** (entry 7, Table 1), from 7-vinylindole, and a late-stage functionalization of a 6,8-dimethyl-cyclopent[*g*]indole nucleus (**15**). Using the chemistry described earlier, **7b** was converted into **14b** in nearly the same yields and selectivities. Not unexpectedly, the direct electrophilic reactions at the C₅-peri-position of **14b** was not feasible in the presence of the more reactive indole C₃ position, even with the

deactivating *N*-Ts moiety. We resorted to hydrogenation of the C₂-C₃ bond before iodination of the substrate using I₂ and PhI(OCOCF₃)₂. A highly regioselective reaction ensued giving **16** as the exclusive product. Subsequent dehydrogenation using Mn(OAc)₃, followed by Stille reaction to install the (*E*)-1-butenyl side-chain gave a good yield of *N*-Ts (+)-*cis*-trikentrin-B. Deprotection of the *N*-Ts group using TBAF gave the natural product.^{4a,14b}

Finally it should be noted that the chemistry outlined here is compatible with the substitution patterns in other known trikentrins and herbindoles,⁷ and it should be possible to prepare any of the *cis*-congeners with complete stereoselectivity using a similar route with only slight modifications. Preparation of the compounds in the *trans*-series can be envisioned using alternate chemistry of the key ketone **11a** or **11b**.¹⁸ Such studies and application of this chemistry to prepare other complex indole alkaloids are in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial assistance for this research by NIH (General Medical Sciences, R01 GM075107) is gratefully acknowledged.

References

- (a) Chyan YJ, Poeggeler B, Omar RA, Chain DG, Frangione B, Ghiso J, Pappolla MA. *J Biol Chem.* 1999; 274:21937. [PubMed: 10419516] (b) Bendheim PE, Poeggeler B, Neria E, Ziv V, Pappolla MA, Chain DG. *J Mol Neurosci.* 2002; 19:213. [PubMed: 12212784] (c) Hwang IK, Yoo KY, Li H, Park OK, Lee CH, Choi JH, Jeong YG, Lee YL, Kim YM, Kwon YG, Won MH. *J Neurosci Res.* 2009; 87:2126. [PubMed: 19235887]
- (a) Rao KV. *Antibiot Chemother.* 1960; 10:312. [PubMed: 14436229] (b) Marsh WS, Garretson AL, Wesel EM. *Antibiot Chemother.* 1960; 10:316. [PubMed: 14421508] For a synthesis, see: (c) Sutou N, Kato K, Akita H. *Tetrahedron: Asymm.* 2008; 19:1833.
- (a) Capon RJ, Rooney F, Murray LM, Collins E, Sim ATR, Rostas JAP, Butler MS, Carroll AR. *J Nat Prod.* 1998; 61:660. [PubMed: 9599272] (b) Wright AE, Pomponi SA, Cross SS, McCarthy P. *J Org Chem.* 1992; 57:4772. For recent synthetic studies of related molecules see: (c) Feldman KS, Ngermeesri P. *Org Lett.* 2011; 13:5704. [PubMed: 21958431] (d) Garg NK, Sarpong R, Stoltz BM. *J Am Chem Soc.* 2002; 124:13179. [PubMed: 12405846]
- (a) Capon RJ, Macleod JK, Scammells PJ. *Tetrahedron.* 1986; 42:6545. (b) Herb R, Carroll AR, Yoshida WY, Scheuer PJ, Paul VJ. *Tetrahedron.* 1990; 46:3089. For examples of other related indole natural products, see: raputindoles: (c) Vougiannopoulou K, Fokialakis N, Aliannis N, Cantrell C, Skaltsounis AL. *Org Lett.* 2010; 12:1908. [PubMed: 20394392] antimalarial flinderroles: (d) Fernandez LS, Buchanan MS, Carroll AR, Feng YJ, Quinn RJ, Avery VM. *Org Lett.* 2009; 11:329. [PubMed: 19090698]
- For recent reviews of functionalization of the indole nucleus, see: (a) Cacchi S, Fabrizi G. *Chem Rev.* 2011; 111:PR215. [PubMed: 21557620] (b) Bandini M, Eichholzer A. *Angew Chem Int Ed.* 2009; 48:9608. (c) Poulsen TB, Jørgensen KA. *Chem Rev.* 2008; 108:2903. [PubMed: 18500844]
- Lee S, MacMillan DWC. *J Am Chem Soc.* 2007; 129:15438. [PubMed: 18031044]
- For recent reviews, see: (a) Silva LF Jr, Craveiro MV, Tebeka IRM. *Tetrahedron.* 2010; 66:3875. (b) Sapeta K, Lebold TP, Kerr MA. *Synlett.* 2011:1495.
- Recent reviews: (a) RajanBabu TV. *Synlett.* 2009:853. [PubMed: 19606231] (b) RajanBabu TV. *Chem Rev.* 2003; 103:2845. [PubMed: 12914483] (c) Jolly PW, Wilke G, Cornils B, Herrmann WA. *Hydrovinylolation. Applied Homogeneous Catalysis with Organometallic Compounds.* VCH New York 1996; 2:1024–1048. For applications in synthesis, see: (d) Lim HJ, RajanBabu TV. *Org Lett.* 2011; 13:6596. [PubMed: 22103775] (e) Mans DJ, Cox GA, RajanBabu TV. *J Am Chem Soc.*

- 2011; 133:5776. [PubMed: 21449569] (f) Smith CR, RajanBabu TV. J Org Chem. 2009; 74:4896. (g) Saha B, Smith CR, RajanBabu TV. J Am Chem Soc. 2008; 130:9000. [PubMed: 18570419] For a detailed procedure of a typical asymmetric hydrovinylation, see: (h) Smith CR, Zhang A, Mans DJ, RajanBabu TV. Org Synth. 2008; 85:248. [PubMed: 19672483] For a Co(II)-catalyzed asymmetric hydrovinylations, see: Sharma RK, RajanBabu TV. J Am Chem Soc. 2010; 132:3295. [PubMed: 20163120] Grut-ter's MMP, Müller C, Vogt D. J Am Chem Soc. 2006; 128:7414. [PubMed: 16756275]
9. Lim HJ, Smith CR, RajanBabu TV. J Org Chem. 2009; 74:4565. [PubMed: 19441793]
10. (a) Bartoli, G.; Bosco, M.; Dalpozzo, R.; Palmieri, G.; Marcantoni, E. J Chem Soc, Perkin Trans. Vol. 1. 1991. p. 2757 (b) Bartoli G, Palmieri G, Bosco M, Dalpozzo R. Tetrahedron Lett. 1989; 30:2129. (c) Dobbs A. J Org Chem. 2001; 66:638. [PubMed: 11429846]
11. (a) Smith CR, Mans DJ, RajanBabu TV. Org Synth. 2008; 85:238. [PubMed: 19655040] (b) Smith CR, RajanBabu TV. Org Lett. 2008; 10:1657. [PubMed: 18351772] For a leading reference to the seminal contributions of the Leitner group, see: (c) Lassauque N, Francio G, Leitner W. Adv Synth Catal. 2009; 351:3133. and references cited therein.
12. See Supporting Information for details of the syntheses of the vinylindoles, hydrovinylation reactions, and spectroscopic and analytical data for all new compounds.
13. For representative examples of recent syntheses of racemic cyclopenta[g]indoles, see: (a) Brown N, Luo DH, Velde DV, Yang SR, Brassfield A, Buszek KR. Tetrahedron Lett. 2009; 50:63. [PubMed: 21057588] (b) Silva LF, Craveiro MV. Org Lett. 2008; 10:5417. [PubMed: 18973334] (c) Jackson SK, Kerr MA. J Org Chem. 2007; 72:1405. [PubMed: 17256909] (d) Huntley RJ, Funk RL. Org Lett. 2006; 8:3403. [PubMed: 16836416] (e) MacLeod JK, Ward A, Willis AC. Austr J Chem. 1998; 51:177. (f) Wiedenau P, Monse B, Blechert S. Tetrahedron. 1995; 51:1167. (g) Boger DL, Zhang MS. J Am Chem Soc. 1991; 113:4230.
14. The reported syntheses of these deceptively looking compounds are lengthy, low-yielding, stereorandom, and involve difficult separations of diastereomeric intermediates. For example, (–)-cis-trikentrin A (ent-**4a**) and (–)-trans-trikentrin A (ent-**4b**) have been synthesized in very low yields (<1%) in a stereorandom fashion starting from (–)-pulegone. (a) Muratake H, Natsume M. Tetrahedron Lett. 1989; 30:5771. cis-Trikentrin-A (**4a**) and cis-trikentrin-B (**5a**) were synthesized in more than 20 steps starting with (–)-malic acid through diastereomeric intermediates that were separated by HPLC (b) Muratake H, Mikawa A, Seino T, Natsume M. Chem Pharm Bull. 1994; 42:854. and references cited therein. For another synthesis of cis-trikentrin-B starting with (+)-menthol (30 steps), see: (c) Lee M, Ikeda I, Kawabe T, Mori S, Kanematsu K. J Org Chem. 1996; 61:3406.
15. This included use of Crabtree's¹⁶ and Pfaltz's Ir-catalysts,^{15b} and various heterogeneous hydrogenation conditions. For Pd/C-catalyzed hydrogenation of a related intermediate see: (a) ref. 13(f). Bell S, Wustenberg B, Kaiser S, Menges F, Netscher T, Pfaltz A. Science. 2006; 311:642. [PubMed: 16339409] We thank Professor Pfaltz for a generous gift of samples of his catalysts.
16. (a) Crabtree RH, Felkin H, Morris GE. J Organomet Chem. 1977; 141:205. (b) Crabtree RH, Felkin H, Fillebeenkhan T, Morris GE. J Organomet Chem. 1979; 168:183. See Supporting Information for details.
17. Yasuhara A, Sakamoto T. Tetrahedron Lett. 1998; 39:595.
18. One approach might be to prepare the corresponding benzyl bromide (NaBH₄, then PBr₃) followed by diastereoselective Negishi reaction (MeZnI, NiBr₂, (S)-i-Pr-PyBOX) to install the trans-1,3-dimethyl pattern as has been demonstrated in a model system (trans:cis = >30:1). See: Arp FO, Fu GC. J Am Chem Soc. 2005; 127:10482. [PubMed: 16045323]

Diastereoselective Synthesis of *cis* or *trans*-Trikentrins and Herbindoles

(see ref. 7a for highly readable graphical summaries of various syntheses)

(-)-*cis*-A: stereorandom from (-) pulegone 10 steps, Natsume, ref. 14a.

(+)-*cis*-A: from (-)-malic acid 21 steps, Natsume, ref. 14b.

(-)-*trans*-A: stereorandom from (-) pulegone 10 steps, Natsume, ref. 14a.

(+)-*cis*-B: from (-)-malic acid 21 steps, Natsume, ref. 14b.

(+)-*cis*-B: from (+)-menthol 30 steps, Kanematsu, ref. 14c.

(+)-*trans*-B: from (-)-malic acid 21 steps, Natsume, ref. 14b.

Herbindoles A, B or C: from (-)-malic acid >20 steps Natsume, ref. 14b.

Figure 1.
Biologically Active Indole Derivatives.

Figure 2.
Ligands for Asymmetric Hydrovinylation of Vinylindoles.

Figure 3.
Major Products from HV of Vinylindoles Using Ligands **L1-L3**

Scheme 1. Selected Examples of Asymmetric Hydrovinylation of Alkenes⁸.

Scheme 2.

Synthesis and Asymmetric Hydrovinylation of a Prototypical Vinylindole

Scheme 3.
Enantioselective Synthesis of (+)-*cis*-Trikentrin A

Scheme 4.
Enantioselective Synthesis of *cis*-Trikentrin B

Table 1

Asymmetric Hydrovinylation of Vinylindoles (VI)^a

entry	substrate	yield (%) / ee (%) ^b			
		L1	L2	L3	L4 ^c
1.	2-VI	96/80	98/82	99/94	98
2.	3-VI	93/91	98/90	96/92	98
3.	7-Et-4-VI	96/89	97/93	98/94	91
4.	5-VI	96 > 90 ^d	98 / > 96 ^d	96 / > 92 ^d	98 ^e
5.	6-VI	95/94	96/95	97/93	97 ^e
6.	1-Ts-6-VI ^f	92/91	93/90	93/89	--
7.	7-VI	97/91	98/93	98/99	90
8.	4-Et-7-VI	96/86	95/87	99/96	90

^a see Scheme 2 and Supporting Information for details.^b determined by chiral stationary phase GC on cyclodex-B column except for entry 4.^c yield of racemic product.^d determined by Mosher method.^e using [L2+*ent*-L2].^f 8 mol% catalyst. %ee of the product determined by GC after detosylation.