Published in final edited form as:

Curr Opin Neurobiol. 2009 August ; 19(4): 439–444. doi:10.1016/j.conb.2009.07.005.

Spontaneous and driven cortical activity: implications for computation

Dario L. Ringach

Department of Neurobiology and Psychology, Jules Stein Eye Institute, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA 90095-1563

Dario L. Ringach: dario@ucla.edu

Abstract

The traditional view of spontaneous neural activity as “noise” has been challenged by recent findings suggesting that: (a) spontaneous activity in cortical populations is highly structured in both space and time, (b) the spatio-temporal structure of spontaneous activity is linked to the underlying connectivity of the cortical network, (c) spontaneous cortical activity interacts with external stimulation to generate responses to the individual presentations of a stimulus, (d) network connectivity is shaped in part by the statistics of natural signals, and (e) ongoing cortical activity represents a continuous top-down prediction/expectation signal that interacts with incoming input to generate an updated representation of the world. These results can be integrated to provide a new framework for the study of cortical computation.

Introduction

Spontaneous activity is defined as the firing of neurons measured in the absence of sensory input. Yet, even under conditions where one attempts to minimize external stimulation the sensory end-organs remain active and generate a weak but continual input to central cortical structures. It has been known that such weak input can lead to perceptual outcomes (hallucinations), as it occurs in cases of sensory deprivation. An example is the Charles Bonnet syndrome, a condition where patients experience various sorts of visual hallucinations due to vision loss, from simple straight lines to detailed and complex pictures [1]. A lesser known but more amusing instance is the recount by Richard Feynman of his experiences undergoing sensory deprivation in one of John Lilly’s isolation tanks, after which he pondered: “[...] whether hallucinations, like dreams, are influenced by what you already have in your mind – from other experiences during the day or before, or from things you are expecting to see” [2]. Indeed, recent findings show that spontaneous activity can have perceptual consequences that reflect “what you already have in your mind”, as it appears directly linked to the intrinsic connectivity of cortical networks and top-down expectations. These studies highlight that there is much to learn about brain function even in its quiescent state, and underlies the importance of understanding cortical responses as the fusion of ongoing activity and sensory input.

© 2009 Elsevier Ltd. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Spontaneous activity is structured in space and time

The relative low rates and apparent randomness in the spontaneous firing pattern of individual cells did not prompt early pioneers of cortical electrophysiology to suspect that they reflected anything else but noise [3]. This picture changed with the first measurements of population activity across large cortical areas, which initially relied on the use of voltage-sensitive dyes (VSDI) [4] and recently augmented by the proliferation of multi-electrode array recordings [5] and *in-vivo*, two-photon imaging [6]. These new measurements reveal a number of important characteristics of spontaneous activity in cortical populations: correlated neural activity can be observed across millimeters of cortex [7,8] and across different time scales [9,10], the firing of individual cells can be related to the patterns of ongoing activity in its neighborhood [11,12], and the size of the fluctuations observed in the spontaneous population activity are of similar magnitude to those the mean response to a high-contrast stimulus [7,13]. These data point to a potential role of spontaneous activity in normal cortical processing, which has motivated investigators to take a fresh look at the role of spontaneous activity in normal cortical processing.

The structure of spontaneous activity reflects the underlying connectivity

The first question is what mechanism might be responsible for the generation of structured patterns of ongoing activity. A promising working hypothesis is that the structure of spontaneous activity reflects the connectivity of the cortical network, and the rules by which neurons at different cortical locations and with different preferences for stimulus attributes connect to each other [14–17]. The presence of functional maps in primary visual cortex provides an ideal model to test this idea. It has been observed that when the spikes of a cell tuned to a particular orientation are used to trigger the averaging of the VSDI signal in its neighborhood, a pattern emerges highlighting cortical columns matching the orientation preference of the cell [11]. This result is expected from a similar bias in lateral connectivity inferred from anatomical and spike cross-correlation data [8,18]. Interestingly, such patterns also emerge in the total absence of an external visual stimulus [19], as if the cortex were to be spontaneously “hallucinating” a physical stimulus with a particular orientation [16]. Theoretical studies have demonstrated that this behavior is expected from an underlying connectivity scheme where nearby cells with similar orientation preference excite each other, while those with different orientation preferences inhibit each other [15,17,20]. Similar connectivity analyses are now being attempted across cortical areas using BOLD signals [21]. The key result from these studies is that the activity of cortical networks under weak input may be used to reveal their intrinsic connectivity, one that may in fact be masked under conditions of strong sensory stimulation.

Spontaneous activity interacts with external stimulation

Spontaneous activity can then offer a handle into brain connectivity, but does it play a functional role in cortical computation? It could be, for example, that external stimulation is so strong that effectively erases any traces of ongoing activity just prior to the presentation of a stimulus, a situation that would weaken the notion of spontaneous activity as having a significant role. All indications are that this is not the case. First, it has been shown that the variability of responses to repeated presentations of the same stimulus can be largely accounted by the ongoing activity just prior to the onset of the stimulus [22]. The simple addition of the spontaneous state of the cortex at the time of stimulus presentation to the mean response provided a reasonable model to explain the responses of the cortex to individual presentations. This result implies that external stimulation does not overwhelm the ongoing cortical activity but interacts with it. Second, external stimulation can lead to response patterns across the cortical population that are similar to those observed in the

spontaneous regime, as if thalamic input triggers a set of stereotyped response patterns [23–25]. Consistent with this notion of attractors, the cortical responses evoked by natural stimulation appear to be very similar to those observed spontaneously [26–28]. Such states may be temporarily protected from being disrupted by external when the network is in a so-called UP state [29], providing a type of short term memory mechanism.

The interaction between the input and the ongoing cortical state depends on the strength of the feed-forward thalamic signals [9,12,30,31]. Strong sensory stimulation drives the cells in the network to de-correlate and respond largely to their local thalamic input. When the stimulus is weak, lateral interactions in the cortex have a substantial influence on the cortical responses [12]. Cortical computation operates in a feed-forward regime for strong stimuli, relies on feedback and top-down signals for weak stimuli, and adopts an intermediate regime for moderate inputs [12,32]. Such adaptations occur locally at each retinotopic location, so that areas of the image with high contrast are processed in a feed-forward fashion, while areas of low contrast rely more on contextual influences (driven by lateral interactions) and top-down predictions. The fact that large areas of natural images contain low-contrast signals [33] suggests that vast areas of the cortex may rely on contextual information to generate an accurate representation of the sensory input.

Network connectivity may be shaped by the statistics of natural stimuli

It is easy to envision how patterned spontaneous activity might be linked to the wiring of the network, but what dictates the wiring of the network itself, how do attractors emerge, and what do they mean? One idea is that evolution and developmental rules during the critical period, where cells that ‘fire-together wire-together’ [34] may shape cortical connectivity. Under some condition, one might expect such connectivity to create spatially localized ‘memories’ (or ‘attractors’) [35] representing the manifold of naturally occurring stimuli [36–38]. In support of such a scenario, the repetitive presentation of natural image sequences leads to activation patterns that are seen to repeat during spontaneous activity [39,40], therefore leaving a memory trace in the network. The statistics of natural images may thus contribute to shape the local network interactions which, in turn, govern the intrinsic dynamics of cortical responses. From a signal processing point projecting the input data into the manifold of natural signals may prove useful for a variety of operations, such as de-noising and perceptual filling-in [36].

Top-down expectation/prediction/attention modulates ongoing activity

Some studies have asked directly if spontaneous activity has a demonstrable perceptual consequence for the processing of external stimuli. In one such case, Super and colleagues presented monkeys with a difficult figure/ground detection task [41]. In this task the same physical stimulus may be detected in some trials, but not perceived in others. They discovered that the likelihood of the animal detecting the stimulus correlated with the spontaneous level of activity in V1 *prior* to the stimulus presentation. In other words, the ability of the animal to detect the stimulus depended critically on the state of the cortex at the time of the presentation. In another study, monkeys’ decisions in a motion detection task were correlated with the spontaneous activity of a neurons in area LIP at the time of stimulus onset, and this correlation was present even in cases where the stimulus had no net motion [42]. Direct contribution of spontaneous activity to inter-trial variability in human behavior has also been shown by means of fMRI [43].

While it was originally proposed that neuronal noise in sensory neurons could provide a “feed-forward” account of these phenomena, it is now becoming increasingly clear that such activity reflects a top-down expectation or prediction signal [44]. A recent experiment has

taken advantage of reverse correlation methods in the binocular disparity domain to show very different temporal courses for stimulus-related and choice-related signals in V2 neurons [45]. During the trial, choice-related activity increased over time while stimulus-related signals decreased, arguing for a top-down modulatory signal of choice. Interestingly, the choice-related activity was significant at the time of stimulus presentation, consistent with prior studies and the notion of top-down expectation.

Conclusions

These results can be integrated to provide a new framework for the study of cortical computation. To summarize the concepts in a simplified setting, consider the case of coding the orientation of a sinusoidal grating by V1 neurons. Each stimulus (Fig 1a, *left*) produces an activity profile across the population that translates along the periodic x -axis as its orientation changes (Fig 1b, *right*, blue curves). The manifold of this space is a circle (Fig 1b). The intrinsic connectivity of the cortex would work continuously to push the state towards the circle, meaning the circle is an attractor of its dynamics (Fig 2c, *middle left*). If, for whatever reason, the cortical state is found outside the manifold it is pushed back into it, as indicated by the blue arrows. Feedforward and feedback signals can be viewed as external forces acting on the state of the cortex. Feedforward input would push the state to the orientation signaled by the LGN (Fig 1c, *bottom*), while feedback input will push the state to one that would represent an expected or predicted one (Fig 1c, *top*). The relative strength of feed-forward drive would depend on the reliability of LGN input signaling a particular orientation, while the strength of top-down signals will depend on reliability of the predictive signal. The final state will be driven by all these forces combined together (Fig 1c, *right*).

In this scenario, the lack of a stimulus generates a weak feedforward drive and the cortical state is determined largely by its own intrinsic dynamics and top-down expectations. This explains why spontaneous activity is not noise. Further, if the attractor has been shaped (via evolution or developmental rules) to represent the manifold of natural signals, it is easy to see how sensory deprivation or weak input can lead to “hallucinations” during deprivation [19,40] and situations where the cortex representing signals related to expectation or choice bias [44,45]. In general, however, the manifold of natural signals is not simply a circle but could be much more complicated (Fig 1d) [36–38], yet all the same ideas apply. It should be mentioned that theoretical versions of these ideas have been put forward before [46–50], and are now finding growing acceptance as general models of signal processing [36].

The above discussion highlights the increasing need to study cortical computation in behaving animals, as it is not possible to study the effects of top-down expectations and context in anesthetized preparations. Further, measurements of population activity, such as VSDI, multi-electrode arrays, or two-photon imaging will be needed to be able to detect the presence of manifold-attractors in cortical activity. Finally, analyses should concentrate on single-trial data that explains how sensory stimulation combines with ongoing activity to determine behavior.

Acknowledgments

I thank David Jentsch and members of Pro-Test @ UCLA for their continuing support. D. L. Ringach is supported by NIH-EY-12816 and NIH-EY-18322 and DARPA FA-8650-06-C-7633.

References

1. Schultz G, Melzack R. The Charles Bonnet syndrome: ‘phantom visual images’. *Perception*. 1991; 20:809–825. [PubMed: 1816537]

2. Feynman, RP. Surely you're joking, Mr Feynman!. W. W. Norton & Company, Inc; New York, N. Y: 1985.
3. Tolhurst DJ, Movshon JA, Dean AF. The Statistical Reliability of Signals in Single Neurons in Cat and Monkey Visual-Cortex. *Vision Research*. 1983; 23:775–785. [PubMed: 6623937]
4. Shoham D, Glaser DE, Arieli A, Kenet T, Wijnbergen C, Toledo Y, Hildesheim R, Grinvald A. Imaging cortical dynamics at high spatial and temporal resolution with novel blue voltage-sensitive dyes. *Neuron*. 1999; 24:791–802. [PubMed: 10624943]
5. Kelly RC, Smith MA, Samonds JM, Kohn A, Bonds AB, Movshon JA, Lee TS. Comparison of recordings from microelectrode arrays and single electrodes in the visual cortex. *J Neurosci*. 2007; 27:261–264. [PubMed: 17215384]
6. Ohki K, Chung S, Ch'ng YH, Kara P, Reid RC. Functional imaging with cellular resolution reveals precise micro-architecture in visual cortex. *Nature*. 2005; 433:597–603. [PubMed: 15660108]
7. Arieli A, Shoham D, Hildesheim R, Grinvald A. Coherent Spatiotemporal Patterns of Ongoing Activity Revealed by Real-Time Optical Imaging Coupled with Single-Unit Recording in the Cat Visual-Cortex. *Journal of Neurophysiology*. 1995; 73:2072–2093. [PubMed: 7623099]
8. Ts'o DY, Gilbert CD, Wiesel TN. Relationships between horizontal interactions and functional architecture in cat striate cortex as revealed by cross-correlation analysis. *J Neurosci*. 1986; 6:1160–1170. [PubMed: 3701413]
9. Smith MA, Kohn A. Spatial and temporal scales of neuronal correlation in primary visual cortex. *J Neurosci*. 2008; 28:12591–12603. [PubMed: 19036953]
- 10. Kohn A, Smith MA. Stimulus dependence of neuronal correlation in primary visual cortex of the macaque. *J Neurosci*. 2005; 25:3661–3673. A careful and detailed study of the different spatial and temporal scales at which correlations in neuronal activity can be seen. The authors propose two different mechanisms linking cells with similar orientation preferences: one with a small spatial scale (~3mm) and high temporal precision, and another with larger spatial scales (>10mm) but slower time constants. [PubMed: 15814797]
11. Tsodyks M, Kenet T, Grinvald A, Arieli A. Linking spontaneous activity of single cortical neurons and the underlying functional architecture. *Science*. 1999; 286:1943–1946. [PubMed: 10583955]
- 12. Nauhaus I, Busse L, Carandini M, Ringach DL. Stimulus contrast modulates functional connectivity in visual cortex. *Nat Neurosci*. 2009; 12:70–76. This study used spike-triggered local field potentials across a micro-electrode array to investigate the influence of lateral connections on the responses of cortical cells at different contrast levels. They found that cortical spikes initiate outward travelling waves, but that the amplitude and distance these waves travel decreases as the strength of the stimulus is increased. The study lends support to the notion that the cortex is in a feed-forward regime when the input is strong, and gradually shifts to a feed-back regime when the stimulus is weak. [PubMed: 19029885]
13. Chen Y, Geisler WS, Seidemann E. Optimal decoding of correlated neural population responses in the primate visual cortex. *Nat Neurosci*. 2006; 9:1412–1420. [PubMed: 17057706]
14. Sporns O, Kotter R. Motifs in brain networks. *PLoS Biol*. 2004; 2:e369. [PubMed: 15510229]
15. Blumenfeld B, Bibitchkov D, Tsodyks M. Neural network model of the primary visual cortex: From functional architecture to lateral connectivity and back. *Journal of Computational Neuroscience*. 2006; 20:219–241. [PubMed: 16699843]
16. Ringach DL. Neuroscience - States of mind. *Nature*. 2003; 425:912–913. [PubMed: 14586455]
17. Goldberg JA, Rokni U, Sompolinsky H. Patterns of ongoing activity and the functional architecture of the primary visual cortex. *Neuron*. 2004; 42:489–500. [PubMed: 15134644]
18. Bosking WH, Zhang Y, Schofield B, Fitzpatrick D. Orientation selectivity and the arrangement of horizontal connections in tree shrew striate cortex. *Journal of Neuroscience*. 1997; 17:2112–2127. [PubMed: 9045738]
19. Kenet T, Bibitchkov D, Tsodyks M, Grinvald A, Arieli A. Spontaneously emerging cortical representations of visual attributes. *Nature*. 2003; 425:954–956. [PubMed: 14586468]
20. Carandini M, Ringach DL. Predictions of a recurrent model of orientation selectivity. *Vision Research*. 1997; 37:3061–3071. [PubMed: 9425519]

21. Vincent JL, Patel GH, Fox MD, Snyder AZ, Baker JT, Van Essen DC, Zempel JM, Snyder LH, Corbetta M, Raichle ME. Intrinsic functional architecture in the anaesthetized monkey brain. *Nature*. 2007; 447:83–86. [PubMed: 17476267]
22. Arieli A, Sterkin A, Grinvald A, Aertsen A. Dynamics of ongoing activity: Explanation of the large variability in evoked cortical responses. *Science*. 1996; 273:1868–1871. [PubMed: 8791593]
23. MacLean JN, Watson BO, Aaron GB, Yuste R. Internal dynamics determine the cortical response to thalamic stimulation. *Neuron*. 2005; 48:811–823. [PubMed: 16337918]
24. Cossart R, Aronov D, Yuste R. Attractor dynamics of network UP states in the neocortex. *Nature*. 2003; 423:283–288. [PubMed: 12748641]
25. Ikegaya Y, Aaron G, Cossart R, Aronov D, Lampl I, Ferster D, Yuste R. Synfire chains and cortical songs: temporal modules of cortical activity. *Science*. 2004; 304:559–564. [PubMed: 15105494]
- 26. Luczak A, Bartho P, Harris KD. Spontaneous events outline the realm of possible sensory responses in neocortical populations. *Neuron*. 2009; 62:413–425. The latest of a series of studies from various groups showing stereotyped patterns of responses in the cortex. This time, the recordings were obtained from rat auditory and somatosensory cortices. Furthermore, these patterns were related to those observed spontaneously, consistent with the idea of attractors of cortical activity. [PubMed: 19447096]
27. Fiser J, Chiu CY, Weliky M. Small modulation of ongoing cortical dynamics by sensory input during natural vision. *Nature*. 2004; 431:573–578. [PubMed: 15457262]
- 28. Singh G, Memoli F, Ishkhanov T, Sapiro G, Carlsson G, Ringach DL. Topological analysis of population activity in visual cortex. *J Vis*. 2008; 8:11, 11–18. This study presents the first rigorous topological analysis of spontaneous and driven population activity. The authors show how tools from the emerging field of computational topology to estimate basic topological signatures of the population activity. [PubMed: 18831634]
29. Watson BO, MacLean JN, Yuste R. UP states protect ongoing cortical activity from thalamic inputs. *PLoS ONE*. 2008; 3:e3971. [PubMed: 19092994]
30. Lampl I, Reichova I, Ferster D. Synchronous membrane potential fluctuations in neurons of the cat visual cortex. *Neuron*. 1999; 22:361–374. [PubMed: 10069341]
31. Polat U, Mizobe K, Pettet MW, Kasamatsu T, Norcia AM. Collinear stimuli regulate visual responses depending on cell's contrast threshold. *Nature*. 1998; 391:580–584. [PubMed: 9468134]
32. Douglas RJ, Martin KA. Recurrent neuronal circuits in the neocortex. *Curr Biol*. 2007; 17:R496–500. [PubMed: 17610826]
33. Chirimuuta M, Clatworthy PL, Tolhurst DJ. Coding of the contrasts in natural images by visual cortex (V1) neurons: a Bayesian approach. *J Opt Soc Am A Opt Image Sci Vis*. 2003; 20:1253–1260. [PubMed: 12868631]
34. Miller KD, Erwin E, Kayser A. Is the development of orientation selectivity instructed by activity? *Journal of Neurobiology*. 1999; 41:44–57. [PubMed: 10504191]
35. Benyishai R, Baror RL, Sompolsky H. Theory of Orientation Tuning in Visual-Cortex. *Proceedings of the National Academy of Sciences of the United States of America*. 1995; 92:3844–3848. [PubMed: 7731993]
- 36. Peyre G. Manifold models for signals and images. *Computer Vision and Image Understanding*. 2009; 113:249–260. A theoretical paper illustrating the basics of manifold learning and the applications to filtering and de-noising of images and sounds. It shows in more rigorous terms some of the technical concepts that were discussed in the present review.
37. Carlsson G, Ishkhanov T, de Silva V, Zornorodian A. On the local behavior of spaces of natural images. *International Journal of Computer Vision*. 2008; 76:1–12.
38. Srivastava A, Lee AB, Simoncelli EP, Zhu SC. On advances in statistical modeling of natural images. *Journal of Mathematical Imaging and Vision*. 2003; 18:17–33.
39. Yao H, Shi L, Han F, Gao H, Dan Y. Rapid learning in cortical coding of visual scenes. *Nat Neurosci*. 2007; 10:772–778. [PubMed: 17468750]
- 40. Han F, Caporale N, Dan Y. Reverberation of recent visual experience in spontaneous cortical waves. *Neuron*. 2008; 60:321–327. Uses voltage-sensitive dyes to demonstrate that after the repetition of a visual stimulus, activity in the cortex reverberates so that it is more frequently

present in the spontaneous activity of the cortex. Such activity could serve to shape the cortical network by the statistics of natural signals. [PubMed: 18957223]

41. Super H, van der Togt C, Spekreijse H, Lamme VA. Internal state of monkey primary visual cortex (V1) predicts figure-ground perception. *J Neurosci.* 2003; 23:3407–3414. [PubMed: 12716948]
42. Shadlen MN, Newsome WT. Neural basis of a perceptual decision in the parietal cortex (area LIP) of the rhesus monkey. *Journal of Neurophysiology.* 2001; 86:1916–1936. [PubMed: 11600651]
43. Fox MD, Snyder AZ, Vincent JL, Raichle ME. Intrinsic fluctuations within cortical systems account for intertrial variability in human behavior. *Neuron.* 2007; 56:171–184. [PubMed: 17920023]
44. Gilbert CD, Sigman M. Brain states: top-down influences in sensory processing. *Neuron.* 2007; 54:677–696. [PubMed: 17553419]
45. Nienborg H, Cumming BG. Decision-related activity in sensory neurons reflects more than a neuron's causal effect. *Nature.* 2009; 459:89–92. [PubMed: 19270683]
46. Lee TS, Mumford D. Hierarchical Bayesian inference in the visual cortex. *J Opt Soc Am A Opt Image Sci Vis.* 2003; 20:1434–1448. [PubMed: 12868647]
47. Lee TS, Mumford D, Romero R, Lamme VAF. The role of the primary visual cortex in higher level vision. *Vision Research.* 1998; 38:2429–2454. [PubMed: 9798008]
48. Mumford D. On the computational architecture of the neocortex. I. The role of the thalamo-cortical loop. *Biol Cybern.* 1991; 65:135–145. [PubMed: 1912004]
49. Yuille A, Kersten D. Vision as Bayesian inference: analysis by synthesis? *Trends Cogn Sci.* 2006; 10:301–308. [PubMed: 16784882]
50. Rao RP, Ballard DH. Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. *Nat Neurosci.* 1999; 2:79–87. [PubMed: 10195184]

Fig 1. Visual summary of main concepts. (a) Coding of orientation by a population of V1 cells. Different oriented gratings produce a profile of activity centered at a different location. Each panel shows the activity of cells with different preferred orientations. (b) Assuming symmetry, the resulting manifold of population activity is a circle. (c) Conceptual view of how intrinsic dynamics, feedforward and feedback drive, combine to drive the cortical state. In each panel, a “force field” is shown that qualitatively describes how each of the individual mechanisms would act when the cortex is found in different states (in this case represented by the plane). The red dots indicate the equilibrium points for the feedforward, feedback and combined components. (d) A more general view of the framework. At any point in time primary visual cortex integrates feed-forward information (indicated by the variable Y) with top-down contextual information provided by extra-striate areas (denoted

by the variable Z). The state of population activity in primary visual cortex is constrained to the manifold of natural signals, represented by the blue surface. One way of thinking about the integration of information is via Bayesian inference, where the V1 circuitry works to find the state within the manifold that maximizes the conditional probability, $p(X|Y, Z)$. Under some assumptions [46], $p(X|Y, Z) \sim p(X|Y)p(X|Z)$. A similar arrangement can be postulated at different levels in the visual hierarchy.