Published in final edited form as:

Brain Res. 2012 April 23; 1450: 11–23. doi:10.1016/j.brainres.2012.02.035.

Knockout of the γ -aminobutyric acid receptor subunit $\alpha 4$ reduces functional δ -containing extrasynaptic receptors in hippocampal pyramidal cells at the onset of puberty

Nicole Sabaliauskas¹, Hui Shen², Gregg E. Homanics³, Sheryl S. Smith², and Chiye Aoki^{1,*}

¹Center for Neural Science, New York University, New York NY 10003

²Department of Physiology and Pharmacology, SUNY Downstate Medical Center, Brooklyn NY 11203

³Departments of Anesthesiology and Pharmacology and Chemical Biology, University of Pittsburgh, Pittsburgh PA 15261

Abstract

Increased plasmalemmal localization of $\alpha 4\beta\delta$ GABA_A receptors (GABARs) occurs in the hippocampal pyramidal cells of female mice at pubertal onset (Shen et al., 2010). This increase occurs on both dendritic spines and shafts of CA1 pyramidal cells and is in response to hormone fluctuations that occur at pubertal onset. However, little is known about how the $\alpha 4$ and δ subunits individually mediate the formation of functional, plasmalemmal $\alpha 4\beta\delta$ GABARs. To determine whether expression of the $\alpha 4$ subunit is necessary for plasmalemmal δ subunit localization at pubertal onset, electron microscopic-immunocytochemistry (EM-ICC) was employed. CA1 pyramidal cells of female $\alpha 4$ knockout (KO) mice were tested for plasmalemmal levels of the δ subunit within dendritic spine and shaft profiles at the onset of puberty. EM-ICC revealed that the $\alpha 4$ and δ subunits localize on dendritic spines and shafts at sites extrasynaptic to GABAergic input at pubertal onset in tissue of wild-type (WT) mice. At pubertal onset, plasmalemmal localization of the δ subunit is reduced 45.9% on dendritic spines, and 56.7% on dendritic shafts with KO of the $\alpha 4$ subunit, as compared to WT tissue, yet levels of intracellular δ immunoreactivity remain unchanged. The decline in plasmalemmal localization is manifested as decreased responsiveness to the GABA agonist gaboxadol at concentrations that are selective for δ -containing GABARs. Additionally, $\alpha 4$ KO mice have larger dendritic spine and shaft profiles. Our findings demonstrate that $\alpha 4$ subunit expression strongly influences the pubertal increase of δ subunits at the plasma membrane, and that genetic deletion of $\alpha 4$ serves as a functional knock-down of δ -containing GABARs.

© 2012 Elsevier B.V. All rights reserved.

*Corresponding author: Chiye Aoki, 4 Washington Place, Room 1056, New York, NY 10003, Phone: (212) 998-3929, Fax: (212) 995-4011, chiye@cns.nyu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CONTRIBUTIONS:

Nicole Sabaliauskas performed the EM-ICC studies, Hui Shen performed the electrophysiology study, Gregg Homanics created and provided the $\alpha 4$ KO mice, Sheryl Smith contributed the experimental design and oversaw electrophysiology experiments, and Chiye Aoki contributed the experimental design and oversaw EM-ICC experiments. All authors read and approved the final draft of the manuscript.

Keywords

Puberty; GABA type A receptor; THP; tonic inhibition; alpha 4; delta; allopregnanolone

1. INTRODUCTION

γ -aminobutyric acid receptors (GABARs) are ubiquitously expressed throughout the CNS, and are responsible for mediating the majority of fast and tonic inhibition. The variety of unique GABAR subunit combinations results in receptors with diverse biophysical properties. The localization patterns of GABAR subunits reveals that $\alpha 1\beta 2\gamma 2$ GABARs are the most frequently-occurring in the CNS (Wisden et al., 1992). GABARs containing the δ subunit (in lieu of the $\gamma 2$ subunit) localize extrasynaptically, due to the lack of interaction of the δ subunit with the GABAergic postsynaptic density protein gephyrin (Liang et al., 2004; Liang et al., 2006; Studler et al., 2002; Wei et al., 2003). As compared to other GABARs, $\alpha 4\beta \delta$ GABARs exhibit a distinct pattern of localization, with lower total surface expression throughout the CNS and with high concentrations observed in the dentate gyrus of the hippocampus and the thalamus (Pirker et al., 2000; Sperk et al., 1997; Sur et al., 1999; Wisden et al., 1992). In contrast to the brief, saturating levels of GABA that synaptically-localized receptors are exposed to, extrasynaptic $\alpha 4\beta \delta$ GABARs are localized to sites displaced from GABA release (Wei et al., 2003), which results in the exposure to low-level, ambient GABA concentrations. This and their relative lack of desensitization enables them to much more effectively maintain a constant inhibitory tone for the neuron (Bianchi and Macdonald, 2002; Glykys and Mody, 2006; Glykys et al., 2007; Stell and Mody, 2002).

Numerous reports have shown that the surface expression of $\alpha 4\beta \delta$ GABARs increases with fluctuations in female gonadal steroids, observed across the ovarian cycle, throughout pregnancy, at the onset of puberty, and when hormone fluctuations are mimicked through both exogenous administration and withdrawal (Griffiths and Lovick, 2005b; Maguire et al., 2005; Maguire et al., 2009; Shen et al., 2007; 2010; Smith et al., 1998a; 1998b; Gulinello et al., 2001; Hsu and Smith, 2003; Shen et al., 2005; Zhou and Smith, 2007). The importance of $\alpha 4\beta \delta$ GABARs was highlighted in recent studies of female brains at puberty (Shen et al., 2007; 2010). The pubertal increase in $\alpha 4\beta \delta$ GABARs reduced hippocampal excitability, NMDA-generated EPSCs, and synaptic plasticity, which is likely responsible for the decline in hippocampus-dependent spatial learning found at that time (Shen et al., 2010). These reports also showed a heightened level of restraint stress-induced or stress steroid-induced anxiety-like behaviors at the onset of puberty, and that this phenomenon can be ameliorated with knockout (KO) of the δ subunit (Shen et al., 2007), suggesting that increased plasmalemmal localization of $\alpha 4\beta \delta$ GABARs in the pyramidal cells of the CA1 hippocampus has a functional relevance for mood and learning at the onset of puberty.

One question that remains to be answered is whether plasmalemmal localization of the δ subunit at puberty requires $\alpha 4$ expression. In order to test this, we assessed δ subunit immunoreactivity within CA1 hippocampal pyramidal cells of $\alpha 4$ KO mice by analyzing tissue harvested from females at the onset of puberty, which is when $\alpha 4\beta \delta$ GABARs increase to peak levels within wild-type (WT) brains (Shen et al., 2007).

In the principal cells of the thalamus and cerebellum, the δ subunit forms functional pentamers predominantly with the $\alpha 4$ or the homologous $\alpha 6$ subunit, respectively (Jones et al 1997; Sur et al 1999). Since the $\alpha 6$ subunit expresses solely in the cerebellum, it is not a potential partner for δ in the hippocampus (Laurie et al., 1992; Nusser et al., 1996; Nusser et al., 1998). However, the $\alpha 1$ subunit can co-localize with the δ subunit, as seen in dentate gyrus interneurons (Glykys et al., 2007). Previous reports show that both the $\alpha 1$ and δ

subunits express in pyramidal cells of the CA1 hippocampus (Nusser et al., 1996; Wisden et al., 1992). Therefore, it is possible that, for CA1 hippocampal pyramidal cells, the absence of $\alpha 4$ subunit expression could still lead to pentamerization of the δ subunit with the $\alpha 1$ subunit, resulting in trafficking of δ -containing GABARs from intracellular stores to the plasma membrane at pubertal onset. Since the majority of surface-trafficked, stable GABARs capable of leaving the endoplasmic reticulum are complete pentamers, if the δ subunit cannot oligomerize with another α subunit, surface expression of the δ subunit would not be expected to increase at puberty (Bencsits, 1999; Green and Millar, 1995; Kittler et al., 2002).

One working model compatible with all of these observations is that the $\alpha 4$ and $\alpha 1$ subunits compete as binding partners for δ . The hormone fluctuations accompanying pubertal onset promote expression of $\alpha 4$, which in turn promotes the formation of $\alpha 4\beta\delta$ GABARs. Our current study addresses whether the absence of the $\alpha 4$ subunit will impact the surface expression of the δ subunit.

The current study differs in multiple significant ways from previous studies that have analyzed δ subunit expression within $\alpha 4$ KO brains. First, the previous studies that have examined surface expression of the δ subunit in $\alpha 4$ KO mice have done so in the thalamus and dentate gyrus (Chandra et al., 2006; Suryanarayanan et al., 2011). In WT tissue, levels of $\alpha 4\beta\delta$ GABARs remain elevated continuously in these brain regions (Wisden et al., 1992). In contrast, the CA1 hippocampus of females at puberty exhibits hormone fluctuation-dependent transient rises in these GABARs at pubertal onset (Shen et al., 2007; 2010). Secondly, in the dentate gyrus, GABAergic current is inward (Staley and Mody, 1992); this is not the case for the CA1 hippocampus, where GABAergic current is outward (Staley and Mody, 1992). Thus, the effect of this current reversal on δ subunit expression in $\alpha 4$ KO tissue is yet unknown. Finally, previous studies examined changes in δ expression after ethanol administration or epileptic seizure, which may depend on mechanisms different from those evoked by the endogenous changes accompanying pubertal onset (Brooks-Kayal et al., 1998; Goodkin et al., 2005; Liang et al., 2007; Sundstrom-Poromaa et al., 2002; Wei et al., 2004). While our previous light microscopic analysis was limited to showing the expression of $\alpha 4$ and δ GABAR subunits in individual cell types (Shen et al., 2007), electron microscopic-immunocytochemistry (EM-ICC) allows for localization to be distinguished as plasmalemmal or intracellular. This classification elucidates potential functionality of $\alpha 4$ - and δ -containing GABARs. Our findings demonstrate that $\alpha 4$ KO mice show reduced plasmalemmal localization of the δ subunit at pubertal onset, and that $\alpha 4$ KO serves as a functional knock-down of δ -containing GABARs.

2. RESULTS

2.1 CA1 hippocampal $\alpha 4$ subunit immunoreactivity in $\alpha 4$ KO mice, as revealed by light microscopy

Light microscopic (LM) observations showed that expression of the $\alpha 4$ subunit was apparent in the CA1 field of the hippocampus of female WT pubertal mice (Figure 1 A). In contrast, qualitative observations revealed that the corresponding region of female $\alpha 4$ KO brains at puberty (Figure 1 B) showed negligible $\alpha 4$ immunoreactivity. In WT pubertal sections, prominent labeling for the $\alpha 4$ subunit in the CA1 hippocampus allowed for identification of pyramidal cells from the otherwise non-immunoreactive neuropil. Pyramidal cells were characterized as such by their triangular shape, small perikarya, and distinct apical dendrite descending into the stratum radiatum of the hippocampus. Interneurons in WT pubertal sections were also labeled for the $\alpha 4$ subunit, and were differentiated from pyramidal cells by their larger perikarya, rounded shape, and displacement from the pyramidal cell layer. Due to the dramatic reduction in $\alpha 4$ immunoreactivity in $\alpha 4$ KO brains, neither pyramidal

cells nor interneurons were discernable from the surrounding neuropil. For quantitative analysis of subunit expression, we employed EM-ICC.

2.2 $\alpha 4$ KO reduces immunoreactivity for the $\alpha 4$ subunit to levels comparable to primary antibody preadsorption controls

To determine the specificity of the $\alpha 4$ antibody for the target protein, we performed a primary antibody preadsorption control. Upon incubation of the $\alpha 4$ primary antibody with the control peptide corresponding to the N-terminus region, we found a substantial reduction in plasmalemmal immunoreactivity in WT pubertal tissue by EM analysis. With $\alpha 4$ primary antibody preadsorption, plasmalemmal labeling was completely eliminated in WT pubertal dendritic spines ($p < 0.00005$), and was reduced by 96.7% in WT pubertal dendritic shafts ($p < 0.000001$). This finding indicates that the primary antibody we are utilizing is specific for the $\alpha 4$ subunit.

Previous studies indicate that the KO of the $\alpha 4$ subunit in this mouse strain was successful, citing a complete elimination of hippocampal $\alpha 4$ immunoreactivity by both western blot and light-microscopic immunohistochemistry (Chandra et al., 2006; Suryanarayanan et al., 2011). An EM-ICC characterization of this strain of $\alpha 4$ KO mice had not been performed until now. In our analysis of $\alpha 4$ KO tissue, we observed that the plasmalemmal $\alpha 4$ immunoreactivity on dendritic spines was reduced by 69.1% as compared to WT levels ($p < 0.0001$; Figure 2 A–C). Plasmalemmal immunoreactivity for the $\alpha 4$ subunit on dendritic shafts was reduced by 81.3% upon $\alpha 4$ KO ($p < 0.000001$; Figure 2 D–F). For both dendritic spines and dendritic shafts, the decrement of $\alpha 4$ immunoreactivity within the $\alpha 4$ KO was comparable to the low levels of labeling observed following omission of the primary antibody ($p > 0.1$).

2.3 Hippocampal pyramidal cells from WT and $\alpha 4$ KO mice exhibit normal neuronal phenotype

We examined the phenotypes of CA1 hippocampal pyramidal cells from WT and $\alpha 4$ KO tissue by conducting LM analyses of two major neuronal phenotypes – immunoreactivity to α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor subunits GluR2/3, and to the microtubule-associated protein 2 (MAP2). CA1 pyramidal cells from pubertal WT and $\alpha 4$ KO mice exhibit equivalent levels of immunolabeling for both the GluR2/3 subunits and MAP2 (Figure 3 A–D). Between the two genotypes, we also observed proper pyramidal cell morphology, and stratification of the CA1 hippocampus like that of WT tissue. Together, these data indicate that hippocampal neurons are likely to be developing normally and are capable of neurotransmission. As such, we concluded that any observed changes in localization of the δ subunit in the CA1 hippocampus at the onset of puberty in $\alpha 4$ KO mice are solely the result of the absence of $\alpha 4$ subunit expression.

2.4 $\alpha 4$ KO hippocampus exhibits reduced δ subunit immunoreactivity at plasma membrane of dendritic spines and shafts at puberty

EM-ICC data provide direct evidence for changes in plasmalemmal δ localization on CA1 pyramidal neurons upon KO of the $\alpha 4$ subunit. This technique also allows for distinguishing between localization at dendritic spines versus shafts and between synaptic versus extrasynaptic sites. On dendritic spine profiles, $\alpha 4$ KO resulted in a 45.9% reduction of plasmalemmal δ immunoreactivity as compared to spines of WT pubertal mice (Table 1; Figure 4 A–C). EM analysis also revealed a significant reduction in plasmalemmal δ labeling on dendritic shafts of KO tissue, relative to the levels seen in WT mice. Labeling for the δ subunit at the dendritic shaft plasma membrane was reduced by 56.7% in tissue of $\alpha 4$ KO pubertal mice as compared to tissue of WT pubertal mice (Table 1; Figure 4 D–F). These EM-ICC data demonstrate that $\alpha 4$ KO mice have reduced plasmalemmal localization

of δ -containing GABARs on dendritic spines and shafts at pubertal onset as compared to WT mice.

In addition to allowing for differentiation between immunoreactivity on dendritic spines and shafts, EM-ICC also allows for the differentiation of synaptic versus extrasynaptic localization along the plasma membrane. In WT pubertal tissue, of the 93 SIG puncta found localized to the plasma membrane from 100 individual dendritic shaft profiles, only one puncta was localized synaptically, while three puncta were localized perisynaptically (within one axon terminal-width of the site of input). The remaining 90 puncta were outside of this boundary, and were classified as extrasynaptic. In $\alpha 4$ KO pubertal tissue, of the 46 plasmalemmal SIG puncta found on 100 dendrites, there were no synaptic puncta, and only one that was perisynaptic, with the remaining 45 localized extrasynaptically. These findings demonstrate that the localization of the δ -containing GABARs is predominantly extrasynaptic, regardless of expression of the $\alpha 4$ subunit.

2.5 Levels of intracellular δ immunoreactivity are equivalent in $\alpha 4$ KO and WT hippocampal dendritic spines and shafts at puberty

After establishing that $\alpha 4$ KO has an effect upon plasmalemmal localization of the δ subunit, we next sought to determine if $\alpha 4$ KO has an effect upon overall δ expression. We did so by comparing the amount of surface immunolabeling relative to the total labeling (surface + intracellular) between WT and $\alpha 4$ KO pubertal tissue. For both dendritic spines and dendritic shafts, we found a significantly higher ratio of surface-to-surface + intracellular immunolabeling in WT pubertal tissue than seen in $\alpha 4$ KO pubertal tissue (Table 1). This difference in ratio is likely due to the observed decrease in plasmalemmal δ localization upon $\alpha 4$ KO, but could be due to alterations in intracellular immunoreactivity. To test this possibility, we measured levels of intracellular δ immunoreactivity in dendritic spines and shafts of $\alpha 4$ KO tissue and compared these levels to those found in WT pubertal tissue. In dendritic spine profiles, $\alpha 4$ KO pubertal mice showed equivalent levels of intracellular δ immunoreactivity as compared to WT pubertal mice (Table 1). We also found no difference in intracellular δ labeling in dendritic shafts between WT and KO pubertal tissue (Table 1). With the equivalent levels of intracellular immunoreactivity between the two genotypes, these EM-ICC data demonstrate that while plasmalemmal localization of the δ subunit on dendritic spines and shafts is compromised in $\alpha 4$ KO, levels of δ -containing GABARs within dendritic spines and shafts at pubertal onset are equivalent in $\alpha 4$ KO and WT pubertal mice.

2.6 $\alpha 4$ KO hippocampus exhibits larger dendritic spine and shaft profiles

Bigger spines would be expected to have more labeling, simply because they occupy more area and membrane within which to potentially express the δ subunit. Was the observed genotype difference in δ immunoreactivity of spines simply due to differences in the size of spines? In order to answer this, we first measured the area and membrane lengths of dendritic spine profiles, as captured in single (not 3D reconstructed) digitized images from pubertal WT and $\alpha 4$ KO tissue. Between the genotypes, we found that larger spines occurred in $\alpha 4$ KO mice, as indicated by increases in both spine head area and membrane length (Table 2). Therefore, the fact that $\alpha 4$ KO mice have less surface labeling despite having more membrane length strengthens the argument that plasmalemmal localization of the δ subunit is dependent upon $\alpha 4$ subunit expression. As intracellular labeling between WT and $\alpha 4$ KO spines is unchanged, despite the larger spine area in $\alpha 4$ KO tissue, it is likely that the δ subunits that cannot localize to the plasma membrane contribute to the intracellular pool in $\alpha 4$ KO tissue.

As the dendritic shaft area and membrane length vary depending upon the plane of the dendrite within the ultrathin section, we measured the minimal diameter of the dendritic shaft profiles from single digitized images to estimate the relative size of shafts. We found that $\alpha 4$ KO mice also have larger apical dendritic shafts as compared to WT mice (Table 2). This increase in diameter would result in a modest increase in the membrane length and membrane area of each dendrite. However, since our measurements for immunoreactivity in and on dendritic shafts are calculated as densities (SIG per μm for surface, and SIG per μm^2 for intracellular), the difference in size between WT and $\alpha 4$ KO tissue is not a factor for our calculations.

2.7 Hippocampal pyramidal cells of pubertal $\alpha 4$ KO mice show a reduced response to the GABAR agonist, gaboxadol at a concentration specific for the δ -containing GABAR

Our EM-ICC data demonstrate that plasmalemmal localization of the δ GABAR subunit is compromised at the onset of puberty upon elimination of $\alpha 4$ subunit expression. As such, the functionality of δ -containing GABARs would be expected to decrease. In order to test this prediction, hippocampal slices from WT and $\alpha 4$ KO pubertal mice were analyzed for functional δ -containing GABAR expression electrophysiologically. Tonic currents were recorded from pyramidal cells in hippocampal slices using whole cell patch clamp procedures. Administration of 100 nM of the GABAR agonist gaboxadol (GBX), which targets only δ -containing GABARs (Brown et al., 2002; Meera et al., 2011), revealed a robust increase in inhibitory current in slices from WT pubertal mice, but only a minimal response in slices from $\alpha 4$ KO pubertal mice (Figure 5 A and B). Subsequent application of the GABAR antagonist gabazine (GBZ), which blocks all GABARs at a concentration of 120 μM , resulted in a restoration of holding current to pre-GBX levels in both WT and $\alpha 4$ KO pubertal slices, indicating that GBX is targeting only GABARs. Our data are in agreement with previously reported findings, which demonstrated decreased GBX sensitivity in ventrobasal thalamic neurons of $\alpha 4$ KO animals (Chandra et al., 2006). These findings indicate that high expression levels of extrasynaptic $\alpha 4\beta\delta$ GABARs contribute to the tonic current in WT pubertal hippocampus, and are in agreement with EM-ICC data, demonstrating that there still are δ -containing GABARs expressing in $\alpha 4$ KO mice at puberty, but their levels are reduced from those seen in WT mice at puberty.

3. DISCUSSION

By comparing δ immunoreactivity in WT and $\alpha 4$ KO pubertal mice, we found that elimination of $\alpha 4$ subunit expression reduced plasmalemmal localization of δ -containing GABARs in the CA1 hippocampus dendritic shaft and spine profiles at the onset of puberty.

3.1 $\alpha 4$ KO reduces $\alpha 4$ immunoreactivity in the CA1 hippocampus

Previous reports indicate that KO of the $\alpha 4$ subunit resulted in elimination of $\alpha 4$ expression in the dentate gyrus of the hippocampus (Chandra et al., 2006; Suryanarayanan et al., 2011). Our data also reveal that $\alpha 4$ immunoreactivity is extremely low in both $\alpha 4$ KO tissue and upon primary antibody preadsorption of WT tissue. The residual immunoreactivity in $\alpha 4$ KO tissue was not further reduced with primary antibody preadsorption. Therefore, the trace amount of $\alpha 4$ labeling seen in $\alpha 4$ KO tissue is most likely due to non-specific antibody binding, and that $\alpha 4$ protein expression is indeed absent in $\alpha 4$ KO mice.

3.2 $\alpha 4$ knockout impairs plasmalemmal δ localization in the CA1 hippocampus at the onset of puberty

The discovery that $\alpha 4$ KO mice show reduced plasmalemmal localization of δ subunits on dendritic spine and shaft profiles compared to WT mice has two possible interpretations: First, it is possible that in the absence of $\alpha 4$ expression, the δ subunit can co-localize with

the $\alpha 1$ subunit, forming $\alpha 1\beta\delta$ GABARs. This is supported by data from other labs, which show expression of $\alpha 1\beta\delta$ GABARs occurring on interneurons of the dentate gyrus (Glykys et al., 2007). Second, it is possible δ subunits localize to the plasma membrane in the absence of other GABAR subunits. While it has been established that the majority of GABARs released from the endoplasmic reticulum for insertion and stabilization in the plasma membrane are fully formed pentamers, it is possible for other non-typical GABARs to traffic to the plasma membrane (Bencsits et al., 1999; Connolly et al., 1996; Kittler et al., 2002; Olsen and Sieghard 2009).

Our EM-ICC data revealed that there were δ -containing GABARs localized to the plasma membrane in pubertal $\alpha 4$ KO mice, albeit at a lower level as compared to pubertal WT mice. Furthermore, our electrophysiology data show, using a selective pharmacological test (100 nM GBX), that δ -containing GABARs are present, but markedly reduced, in hippocampal slices from $\alpha 4$ KO mice as compared to those from WT mice. Therefore, the possible interpretation we favor is the following: in the absence of $\alpha 4$ subunits, the δ subunit can form a full pentamer with the $\alpha 1$ and β subunits and that these localize to extrasynaptic sites of the plasma membrane. However, because this pentamer is not a favored GABAR composition, levels of plasma membrane insertion of putative $\alpha 1\beta\delta$ GABARs at puberty are much less than those for the $\alpha 4\beta\delta$ GABARs in WT tissue.

3.3 $\alpha 4$ knockout does not impair the rise of intracellular δ in the CA1 hippocampus at the onset of puberty

Electron microscopic analysis revealed that knockout of the $\alpha 4$ subunit does not affect intracellular stores of the δ subunit. In both WT and $\alpha 4$ KO pubertal sections, we found equivalent levels of intracellular stores of the δ subunit, both in dendritic spines and shafts. This equivalent labeling for intracellular δ in dendritic spines and shafts in WT and $\alpha 4$ KO animals suggests that at puberty, synthesis of δ subunits is not dependent upon the presence of the $\alpha 4$ subunit, but that trafficking to the membrane is dependent upon $\alpha 4$ expression. There is a novel explanation for the source of intracellular δ subunits in dendritic spines and shafts. It is known that predominantly fully-formed GABARs are able to overcome ER retention (Connolly et al. 1996; Kittler et al. 2002). Since it has already been proposed that $\alpha 1\beta\delta$ GABARs are possibly expressed in the CA1 hippocampal apical dendrites, intracellular pools of δ may be representative of fully formed $\alpha 1$ -containing GABARs. While protein synthesis is most often thought to occur in the somatic region, studies reveal that the cellular machinery required for mRNA translation has been found in both dendritic spines and shafts of the CA1 hippocampus (Chicurel et al. 1993; Bitran et al. 1999; Ostroff et al. 2002; McCarthy and Milner 2003; Wang and Tiedge 2004; Bourne et al. 2007). Therefore, it is also possible that intracellular δ expression in the distal processes of $\alpha 4$ KO mice is indicative of isolated δ subunits that are synthesized locally and sequestered within the ER.

3.4 $\alpha 4$ KO yields larger dendritic spine and shaft profiles

Our EM-ICC and electrophysiology data reveal decreased functional δ -containing GABARs on CA1 hippocampal pyramidal cells. As such, these pyramidal cells should exhibit an increase in excitability and a decrease in the current necessary for triggering a spike due to the decreased GABAergic inhibition they receive. Upon examining the morphology of CA1 pyramidal cells between the two genotypes, we found that pyramidal cells of $\alpha 4$ KO hippocampi have larger dendritic spines, as measured by spine area and membrane length, and dendritic shafts, as measured by shaft diameter, as compared to those from WT hippocampi. These findings indicate that the expression of the $\alpha 4$ subunit has an effect upon the size of dendritic spine and shaft profiles. Furthermore, this increase in profile size would be expected to decrease the input resistance of the pyramidal cells, thus increasing the

current necessary for triggering the neuron to fire a spike. Therefore, it is possible that neurons are undergoing a compensatory change in profile size in response to the decreased inhibition caused by $\alpha 4$ KO.

3.5 Potential functional outcomes of $\alpha 4$ KO

It has been reported that certain learning deficits are observed at the onset of puberty (Johnson and Newport, 1989; McGivern et al., 2002). Previous research from our laboratory has shown that increased plasmalemmal localization of $\alpha 4\beta\delta$ GABARs on CA1 hippocampal pyramidal cells increases the shunting, tonic inhibition and impairs activation of NMDA receptors (Shen et al, 2010). This leads to an impairment in the induction of long-term potentiation and a decline in spatial memory at the onset of puberty (Shen et al., 2010). Furthermore, when δ KO mice were tested for their spatial learning abilities, we found that the pubertal spatial learning deficit was eliminated in these mice (Shen et al., 2010). The EM-ICC and physiology data from the present study show that, in the absence of the $\alpha 4$ subunit, the δ subunit shows minimal plasmalemmal localization on pyramidal cells, both on CA1 dendritic shaft and spine profiles. Therefore, based on this finding, we would expect the spatial learning decline observed at the onset of puberty in WT female mice to also be eliminated in $\alpha 4$ KO mice.

Additional research from this laboratory has shown that $\alpha 4\beta\delta$ GABARs on pyramidal cells of the CA1 hippocampus mediate an increase in stress steroid-induced anxiety-like behaviors at the onset of puberty (Shen et al., 2007). The release of the progesterone metabolite, THP, is known to occur in response to stress (Girdler and Klatzkin, 2007; Purdy et al., 1991). In adults, neurosteroids such as THP acts to subsequently reduce the anxiety-inducing effects of stress (Bitran et al., 1999) by potentiating most GABARs (Belelli et al., 2002; Bianchi and Macdonald, 2003; Brown et al., 2002; Wohlfarth et al., 2002). However, the effect of THP on $\alpha 4\beta\delta$ GABARs is uniquely polarity-dependent, such that it increases inward current (outward flux of Cl^-), but decreases outward current (inward flux of Cl^-) at this receptor subunit combination (Shen et al., 2007). THP has been shown to rapidly desensitize outward current gated by $\alpha 4\beta\delta$ GABARs expressed in human embryonic kidney cells (HEK-293) (Shen et al., 2007). At puberty, when plasmalemmal localization of $\alpha 4\beta\delta$ GABARs is increased on CA1 hippocampal pyramidal cells, where GABAergic current is outward, THP administration reduces the shunting tonic inhibition produced by these receptors, thereby increasing the excitability of hippocampal neurons and of anxiety-like behaviors (Shen et al., 2007; Shen et al., 2010). Our EM-ICC and physiology data report limited localization of the δ subunit to the plasma membrane upon elimination of $\alpha 4$ subunit expression. Therefore, due to the absence of $\alpha 4\beta\delta$ GABARs, we would expect THP to diminish stress-induced excitability of CA1 pyramidal neurons and of anxiety-like behaviors, much like what is observed in adult mice (Bitran et al., 1999). In addition to mediating stress steroid anxiety responses, $\alpha 4\beta\delta$ GABARs are also increased by neuronal activity and traumatic brain injury (Mtchedlishvili et al., 2010; Payne et al., 2008; Santhakumar et al., 2010), which suggests they may also serve a neuroprotective role (Santhakumar et al., 2010).

The data presented here suggest that $\alpha 4$ subunit expression strongly affects the increases in the plasmalemmal δ subunit localization at the onset of puberty. While the mechanisms which underlie $\alpha 4\beta\delta$ GABAR expression at puberty have not yet been elucidated, recent work by other labs has described some of the molecular mechanisms of $\alpha 4$ promoter regulation, which can involve brain derived neurotrophic factor (BDNF) and heat shock proteins (Roberts et al., 2006; Pignataro et al., 2007). These findings may have relevance for alterations in mood and learning reported across fluctuations in ovarian hormones not only at pubertal onset, but also across the estrous and menstrual cycles, during and after pregnancy, and at menopause.

4. EXPERIMENTAL PROCEDURES

4.1 Experimental subjects

Twenty pubertal (36 to 43 days old) female mice were housed on a 12-hour reverse light-dark cycle. $\alpha 4$ KO mice were generated as described previously (Chandra et al., 2006). Heterozygous $\alpha 4$ KO mice on a C57BL/6J $\times$ Strain 129X1/S1 genetic background of the F6 generation were imported from the University of Pittsburgh. Heterozygous mice were bred and offspring were genotyped using Southern blot analysis, as described previously (Chandra et al., 2006). Homozygous offspring were subsequently bred with each other to produce the $\alpha 4$ KO mice used for EM-ICC and slice electrophysiology in this study. C57BL/6J WT mice used for EM-ICC and slice electrophysiology were purchased from Jackson Laboratories (Bar Harbor, Maine). For EM-ICC analysis, mice from single shipments from Jackson Laboratories were perfused for pubertal time-points.

Female mice were chosen for use in this study because our previous findings showing increased $\alpha 4\beta\delta$ GABAR expression were noted in females. Vaginal opening was used to determine the onset of puberty because this physical sign is directly correlated with the hormonal changes that trigger $\alpha 4\beta\delta$ GABAR expression (Shen et al., 2007). Pubertal mice were used after the emergence of the vaginal opening. At these ages, the stage of the estrous cycle does not affect expression of the $\alpha 4$ and δ subunits (Shen et al., 2010). Five WT pubertal, and five $\alpha 4$ KO pubertal mice were utilized in this study for EM-ICC. An additional 5 $\alpha 4$ KO pubertal and 5 WT pubertal mice were used for slice electrophysiology.

All procedures involving live animals were in accordance with NIH guidelines and the Institutional Animal Care and Use Committees of SUNY Downstate Medical Center, University of Pittsburgh, and New York University Washington Square Campus.

4.2 Characterization of antibodies

The $\alpha 4$ antibody utilized in our experiments was purchased from Santa Cruz Biotechnology (catalog #sc-7355). This antibody was an affinity-purified goat polyclonal antibody directed against the extracellular N-terminus (amino acids 32–50) of the human origin peptide sequence. The $\alpha 4$ antibody used has been characterized previously (Griffiths and Lovick, 2005a; Griffiths and Lovick, 2005b; Sanna et al., 2003). On western blots, this antibody has been shown to recognize a single protein of the appropriate molecular weight (Sanna et al 2003; Griffiths and Lovick 2005a). Ultimate proof of antibody specificity using tissue from $\alpha 4$ KO mice had not yet been performed, and the results of such an experiment are described in this manuscript.

The δ primary antibody utilized in our experiments was obtained as a generous gift from W. Sieghart (Medical University Vienna). This antibody was an affinity purified rabbit polyclonal antibody directed against the extracellular N-terminus (amino acids 1–44) of the rat origin peptide sequence. The δ antibody used has been characterized previously by western blot analysis, in which it recognized a single band of the appropriate molecular weight (Fritschy and Mohler, 1995; Jechlinger et al., 1998). In brains from δ KO mice, no immunoreactivity was observed using the same antibody batch as was utilized in these experiments, indicating antibody specificity for the target antigen (Peng et al., 2002).

The GluR2/3 (Millipore catalog #AB1506) and MAP2 (Boehringer Mannheim Biochemicals catalog #1284 959) primary antibodies utilized in our experiments were affinity-purified. Each antibody has been extensively characterized, and recognizes a single protein of the appropriate molecular weight.

4.3 Preparation of brain tissue for EM analysis

All mice were anesthetized with 20% urethane (0.05–0.2 mL intraperitoneally) from 30 minutes to 3 hours before the onset of the dark stage of the circadian cycle and perfused transcardially using a peristaltic pump (flow rate 30 mL/min), first with saline containing heparin (1U/mL) followed by 360 mL of 4% paraformaldehyde buffered to pH 7.4 with 0.1M phosphate buffer (PB) over a 12 minute period. The brains were removed from their skulls and stored in the same fixative until sectioning, which occurred within one week after perfusion. Urethane was chosen as an anesthetic, rather than Nembutal, because Nembutal predominantly targets GABA receptors, the subject of this study, while urethane targets multiple neurotransmitter systems in a more balanced fashion (Hara and Harris, 2002).

All procedures involving transcardial perfusion of animals were carried out in accordance with the NIH guidelines and were approved by the New York University and SUNY Downstate Animal Care and Use Committees.

4.4 Preparation of sections for immunocytochemistry

The tissue processing protocols described previously were followed (Aoki et al., 2000). Coronal sections were cut on a vibratome (Leica VT 100M) at a thickness of 40 μ m, then stored at 4° C in a buffer that contained 0.01M PB with 0.9% sodium chloride (PBS), supplemented with 0.05% sodium azide to prevent bacterial growth. Sections containing dorsal hippocampus were selected for immunocytochemical analysis. Prior to the immunocytochemical procedure, sections were freeze-thaw permeabilized, then treated with 1% hydrogen peroxide (H₂O₂) at room temperature for 30 minutes to retrieve antigenicity, and blocked in 0.01M PBS supplemented with 1% bovine serum albumin and 0.05% sodium azide (PBS/BSA/Azide) for 30 minutes.

4.5 Use of light microscopy

In order to gain an initial insight into the α 4 immunoreactivity in α 4 KO tissue, sections were analyzed qualitatively by light microscopy (LM) using silver-intensified gold (SIG) as the immunolabel. LM allowed for assessment of global expression level changes in α 4 immunoreactivity. For LM analysis, the ability to identify individual cell types (pyramidal cells, interneurons, and glia) due to contrast generated by the presence of immunoreactivity was used as evidence of expression of the α 4 subunit.

LM was also utilized to assess CA1 hippocampal neuronal morphology. Immunoreactivity for the neuronal markers GluR2/3 and MAP2 allowed for characterization of neuronal phenotype and proper stratification of the CA1 hippocampus.

4.6 Choice of immunolabels

Labeling for the α 4 and δ subunits is readily detectable in both dendritic spine profiles and apical dendritic shafts of CA1 hippocampal pyramidal cells at the onset of puberty using SIG as an immunolabel for electron microscopy (Shen et al., 2010). Therefore, we utilized this immunolabel for our electron microscopic analysis. Because SIG is a particulate label, levels of immunoreactivity are clearly quantifiable and resolvable as plasmalemmal versus intracellular with respect to its location. Furthermore, due to the lack of diffusion of SIG particles, labeling can be characterized as occurring synaptically or extrasynaptically. However, due to the lack of enzymatic amplification of the marker, this results in a very high threshold for detection.

Horseradish peroxidase-diaminobenzidine (HRP-DAB) was chosen as an immunolabel for the neuronal markers MAP2 and GluR2/3. Due to the enzymatic amplification step of HRP-DAB labeling, immunoreactivity could be readily detected at the LM level. Furthermore, the

diffuse nature of HRP-DAB provides a more complete saturation of cell labeling, allowing for better structural resolution.

4.7 Procedure using horseradish peroxidase-diaminobenzidine (HRP-DAB) as the immunolabel for light microscopy

Sections were incubated overnight at room temperature in a primary antibody directed against the GluR2/3 subunit of AMPA receptors (1.5 µg/ml) or MAP2 (5 µg/ml), using PBS/BSA/Azide as the diluent. Sections were then rinsed, and incubated in 2° antibody for 3 hours at room temperature (biotinylated anti-rabbit IgG, 1:200 for GluR2/3 1° antibody; biotinylated anti-mouse IgG, 1:200 for MAP2 1° antibody, Vector), followed by Vector's ABC mix ('Elite' kit) for 30 minutes. The HRP reaction used 0.3% DAB with 0.003% H₂O₂ as a substrate, stopped at 10 minutes with PBS rinses. Sections were dehydrated serially with ethanol, defatted with xylene, mounted on slides, and coverslipped for light microscopic analysis. All steps in the immunocytochemical procedure were conducted in parallel for WT and α4 KO tissue.

4.8 Procedure using silver-intensified gold (SIG) as the immunolabel for electron microscopy

Sections were incubated overnight at room temperature in a primary antibody directed against the α4 subunit (1:100 dilution from 0.2 µg/ml) or the δ subunit (1:400 dilution from 0.23 µg/ml), using PBS/BSA/Azide as the diluent. Sections were then incubated in 0.8 nm colloidal gold-conjugated 2° antibody for a minimum of 3 hours at room temperature (horse-anti-goat IgG, 1:100 for α4 immunocytochemistry; goat-anti-rabbit IgG, 1:100 for δ immunocytochemistry, EM Sciences), followed by post-fixation in 2% glutaraldehyde in PBS for 10 minutes. Prior to the silver intensification step, sections were rinsed for 1 minute in a 0.1M citric acid buffer (trisodium citrate salt adjusted to pH 7.4 with monohydrate citric acid salt) to remove phosphate and Cl⁻ ions. Silver IntensEM kit (KPL, Kirkegaard & Perry Laboratories, Inc.) was applied for 10–12 minutes. The silver intensification reaction was terminated by a brief citrate buffer rinse, followed by PBS rinses. Ultrastructure and SIG particles were preserved using several post-fixation steps that avoid the use of osmium tetroxide: 1% tannic acid for 40 min, 1% uranyl acetate for 40 min, and 0.5% iridium tetrabromide for 20 min, all dissolved in 0.1M maleate buffer (maleic acid monosodium salt adjusted to pH 6.0 with maleic acid disodium salt). Vibratome sections were further processed with incubation in 1% *p*-phenylenediamine for 15 min and 1% uranyl acetate overnight, both dissolved in 70% ethanol (Phend et al., 1995). Sections were then dehydrated serially with ethanol and acetone, and infiltrated with EPON-812 (EMScience), first in a 1:1 concentration with acetone, then in full EPON-812. Sections were flat embedded between sheets of Aclar plastic, and ultrathin sectioned at a thickness of 70±1.5 nm, as measured using the minimum folds method, and collected on formvar-coated, 200-mesh EM grids. Grids were counterstained with 2% lead citrate for 30 seconds to increase contrast prior to imaging. All steps in the immunocytochemical procedure were conducted in parallel and with the experimenter blind to the genotype of the tissue source.

4.9 Selection of the CA1 hippocampal region for the electron microscopic study

In order to obtain data from a more uniform population of dendritic spine profiles, EM analysis was restricted to the dorsal CA1 hippocampal pyramidal cell apical dendrites in stratum radiatum. Schaffer collateral input from the CA3 terminates on dendritic spine profiles of the stratum radiatum (Megas et al., 2001; Smith et al., 2003). This region was chosen due to the role that hippocampal GABAR expression plays in the alterations in mood, spatial learning, and neuronal excitability that occur at the onset of puberty (Shen et al., 2007; 2010).

4.10 Dendritic spine analysis

For dendritic spine analysis, spine profiles were identified by the presence of an excitatory synaptic junction. These asymmetric synapses were composed of an electron dense post-synaptic density (PSD) on one side of the synapse, which was immediately opposed by a presynaptic terminal, containing clusters of small, round, clear synaptic vesicles. It has been demonstrated extensively that dendritic spines are biological entities capable of acting autonomously of one another regardless of arising from the same parent dendrite (Carter et al., 2007; Matsuzaki et al., 2004; Yuste and Denk, 1995). Therefore, measurements of $\alpha 4$ and δ immunoreactivity in spines were pooled from hippocampi from multiple mice. A spine was considered labeled for the GABAR subunit when it contained even a single SIG particle. Plasmalemmal labeling was distinguished from intracellular labeling when the SIG was in contact with the plasma membrane of the spine head. A minimum of $380 \mu\text{m}^2$ of neuropil containing at least 110 independent spine profiles was sampled from each mouse for analysis of $\alpha 4$ and δ immunoreactivity within dendritic spines.

4.11 Dendritic shaft analysis

CA1 hippocampal apical dendrites were imaged for the dendritic shaft analysis. Apical dendrites were identified by their parallel protrusion from pyramidal cells into the stratum radiatum. Dendritic shaft membranes were considered labeled when SIG particles were found in contact with the plasma membrane, as this labeling is indicative of potentially functional receptors. Labeling was characterized as synaptic when it occurred directly opposed to the presynaptic terminal, and perisynaptic when it occurred within one axon terminal's width of the presynaptic terminal. All labeling outside of these margins was characterized as extrasynaptic. The total membrane lengths of dendritic shafts were measured using the segmented line tool of ImageJ software (NIH version 10.2), and labeling was quantified as the number of immunolabeled clusters occurring per μm of dendritic plasma membrane, also referred to as linear density. A minimum of $1597 \mu\text{m}^2$ of hippocampal neuropil containing at least 20 dendrites was sampled for each mouse for $\alpha 4$ and δ dendritic shaft analysis.

4.12 Image acquisition

The surface-most portion of vibratome sections from each mouse was examined for labeling on a JEOL transmission electron microscope (1200 EX II), as determined by proximity to the tissue-EPON interface, to ensure adequate antibody exposure. Images were digitally acquired (AMT) under identical conditions across genotypes, using a CCD camera (Hamamatsu) at magnifications of $20,000\times$ (for dendritic shaft analysis) and $40,000\times$ (for dendritic spine analysis). For dendritic spine analysis, non-overlapping images from the CA1 stratum radiatum were acquired. For dendritic shaft analysis, images of at least 20 pyramidal cell apical dendrites per mouse were acquired. The experimenter remained blind to the genotype of the tissue source throughout image acquisition and analysis.

4.13 Capturing and reproduction of figures

Electron microscopic images were captured and stored using the CCD camera attached to the JEOL 1200XL via software developed by AMT. Light microscopic images were captured at a magnification of $20\times$ using a CCD camera attached to a Zeiss Axiophot microscope. For reproduction, a subset of these images was adjusted in contrast, brightness, and image size using Adobe Photoshop (v. CS3).

4.14 Hippocampal slice electrophysiology

$250 \mu\text{m}$ -thick hippocampal slices were generated on a Leica oscillating microtome as described previously (Shen et al., 2007). Pyramidal cells in the CA1 hippocampal slices

were visualized using a Leica DIC-infrared upright microscope, and recorded using whole cell patch clamp procedures in both voltage and current clamp modes. Patch pipets were fabricated from borosilicate glass using a Flaming-Brown puller (Sutter Instruments) to yield open tip release of 2–4 M Ω . Bath solution contained (in mM): NaCl 124, KCl 5, CaCl₂ 2, KH₂PO₄ 1.25, MgSO₄ 2, NaHCO₃ 26 and glucose 10, saturated with 95% O₂, 5% CO₂ and buffered to pH 7.4. Recordings were carried out at room temperature (22–24°C) at a 10kHz sampling frequency (2kHz 4-pole Bessel filter) using an Axopatch 200B amplifier (Axon Instruments) and pClamp 9.2 software. Electrode capacitance and series resistance were monitored and compensated. Access resistance was monitored throughout the experiment, and cells discarded if the access resistance was greater than 10 M Ω .

The pipet solution consisted of (in mM): CsCl 140, MgCl₂ 2, HEPES 10, BAPTA 10, Mg-ATP 2, CaCl₂-H₂O 0.5, Li-GTP 0.5, pH 7.2, 290 mOsm. Voltage clamp recordings were carried out at a –60 mV holding potential. In order to determine the effect of 100 nM GBX (Sigma) and 120 μ M GBZ (Sigma) on the tonic inhibitory current, drugs were bath-applied after 15–20 minutes of recording baseline current in the presence of 2 mM kynurenic acid (Sigma). Responses were assessed by the shift in the holding current.

4.15 Statistics

All statistical analysis for EM-ICC and physiological experiments was performed using the Statistica program (Statsoft). The Shapiro-Wilks W-test was used to determine if the data were normally distributed. When the data were found not to be normally distributed, the Mann-Whitney U-test was subsequently employed to determine significance. When data were normally distributed, Student's t-test was employed to determine significance. For all statistical analyses, significance was accepted for $p < 0.05$.

Highlights

Hippocampal pyramidal cell dendritic spines and shafts express $\alpha 4\beta\delta$ GABA receptors
 Plasmalemmal $\alpha 4\beta\delta$ GABA receptors occur on dendritic spines, near excitatory synapses
 $\alpha 4$ subunit knockout reduces plasmalemmal, but not intracellular, δ localization in pubertal CA1
 $\alpha 4$ subunit knockout yields larger CA1 hippocampal dendritic spine and shaft profiles

Acknowledgments

We thank Dr. Werner Sieghart for providing the δ antibody, Tunazzina Ahmed and Gauri Wable for technical assistance, and Tara Chowdhury for reading the manuscript.

ROLE OF THE FUNDING SOURCE:

This research was supported by NEI Core grant EY13079 and Klarman Family Grants Program in Eating Disorders Research and 1R21MH091445-01 to C.A., NIH Grant AA10422 to G.H., and NIH Grants DA09618 and AA12958 to S.S.S.

REFERENCES

- Aoki C, Rodrigues S, Kurose H. Use of electron microscopy in the detection of adrenergic receptors. *Methods in Molecular Biology*. 2000; 126:535–563. [PubMed: 10685434]
 Barnard EA, Skolnick P, Olsen RW, Mohler H, Sieghart W, Biggio G, Braestrup C, Bateson AN, Langer SZ. International Union of Pharmacology. XV. Subtypes of GABA_A receptors:

- classification on the basis of subunit structure and receptor function. *Pharmacological Reviews*. 1998; 50:291–313. [PubMed: 9647870]
- Bencsits E, Ebert V, Tretter V, Sieghart W. A significant part of native GABA_A receptors containing $\alpha 4$ subunits do not contain γ or δ subunits. *The Journal of Biological Chemistry*. 1999; 274:19613–19616. [PubMed: 10391897]
- Belelli D, Casula A, Ling A, Lambert J. The influence of subunit composition on the interaction of neurosteroids with GABA_A receptors. *Neuropharmacology*. 2002; 43:651–661. [PubMed: 12367610]
- Bianchi M, Macdonald R. Neurosteroids shift partial agonist activation of GABA_A receptor channels from low- or high-efficacy gating patterns. *The Journal of Neuroscience*. 2003; 23:10934–10943. [PubMed: 14645489]
- Bitran D, Dugan M, Renda P, Ellis R, Foley M. Anxiolytic effects of the neuroactive steroid pregnanolone (3 α -hydroxy-5 β -pregnan-20-one) after microinjection in the dorsal hippocampus and lateral septum. *Brain Research*. 1999; 850:217–224. [PubMed: 10629767]
- Brooks-Kayal A, Shumate M, Jin H, Rikhter T, Coulter D. Selective changes in single cell GABA_A receptor subunit expression and function in temporal lobe epilepsy. *Nature Medicine*. 1998; 4:1166–1172.
- Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA. Pharmacological characterization of a novel cell line expressing human $\alpha 4\beta 3\delta$ GABA_A receptors. *British Journal of Pharmacology*. 2002; 136:965–974. [PubMed: 12145096]
- Buchanan C, Eccles J, Becker J. Are adolescents the victims of raging hormones: evidence for activational effects of hormones on moods and behavior at adolescence. *Psychological Bulletin*. 1992; 111:62–107.
- Carter AG, Soler-Llavina GJ, Sabatini BL. Timing and location of synaptic inputs determine modes of subthreshold integration in striatal medium spiny neurons. *The Journal of Neuroscience*. 2007; 27:8967–8977. [PubMed: 17699678]
- Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, Spigelman I, Houser CR, Olsen RW, Harrison NL, Homanics GE. GABA_A receptor $\alpha 4$ subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and the action of gaboxadol. *Proceedings of the National Academy of Sciences of the United States of America*. 2006; 103:15230–15235. [PubMed: 17005728]
- Connolly CN, Krishek BJ, McDonald BJ, Smart TG, Moss SJ. Assembly and cell surface expression of heteromeric and homomeric GABA_A receptors. *Journal of Biological Chemistry*. 1996; 271:89–96. [PubMed: 8550630]
- Fritschy JM, Mohler H. GABA_A receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *Journal of Comparative Neurology*. 1995; 359:154–194. [PubMed: 8557845]
- Girdler S, Klatzkin R. Neurosteroids in the context of stress: implications for depressive disorders. *Pharmacology and Therapeutics*. 2007; 116:125–139. [PubMed: 17597217]
- Glykys J, Mody I. Hippocampal network hyperactivity after selective reduction of tonic inhibition in GABA_A receptor $\alpha 5$ subunit-deficient mice. *Journal of Neurophysiology*. 2006; 95:2796–2807. [PubMed: 16452257]
- Glykys J, Peng Z, Chandra D, Homanics G, Houser C, Mody I. A new naturally occurring GABA_A receptor subunit partnership with high sensitivity to ethanol. *Nature Neuroscience*. 2007; 10:40–48.
- Goodkin H, Joshi S, Mtchedlishvili Z, Brar J, Kapur J. Subunit-specific trafficking of GABA_A receptors during status epilepticus. *The Journal of Neuroscience*. 2008; 28:2527–2538. [PubMed: 18322097]
- Green WN, Millar NS. Ion-channel assembly. *Trends in Neurosciences*. 1995; 18:280–287. [PubMed: 7571003]
- Griffiths J, Lovick T. Withdrawal from progesterone increases expression of $\alpha 4$, $\beta 1$, and δ GABA_A receptor subunits in neurons in the periaqueductal gray matter in female Wistar rats. *Journal of Comparative Neurology*. 2005a; 486:89–97. [PubMed: 15834956]

- Griffiths JL, Lovick TA. GABAergic neurones in the rat periaqueductal grey matter express $\alpha 4$, $\beta 1$, and δ GABA_A receptor subunits: plasticity of expression during the estrous cycle. *Neuroscience*. 2005b; 136:457–466. [PubMed: 16226387]
- Grobin AC, Morrow AL. 3α -hydroxy- 5α -pregnan-20-one exposure reduces GABA_A receptor $\alpha 4$ subunit mRNA levels. *European Journal of Pharmacology*. 2000; 409:R1–R2. [PubMed: 11104837]
- Gulinello M, Gong QH, Li X, Smith SS. Short-term exposure to a neuroactive steroid increases $\alpha 4$ GABA_A receptor subunit levels in association with increased anxiety in the female rat. *Brain Research*. 2001; 910:55–66. [PubMed: 11489254]
- Hara K, Harris RA. The anesthetic mechanism of urethane: the effects on neurotransmitter-gated ion channels. *Anesthesia and Analgesia*. 2002; 94:313–318. table of contents. [PubMed: 11812690]
- Hsu F-C, Smith S. Progesterone withdrawal reduces paired-pulse inhibition in rat hippocampus: dependence on GABA_A receptor $\alpha 4$ subunit upregulation. *Journal of Neurophysiology*. 2003; 89:186–198. [PubMed: 12522171]
- Hsu F-C, Waldeck R, Faber D, Smith S. Neurosteroid effects on GABAergic synaptic plasticity in hippocampus. *Journal of Neurophysiology*. 2003; 89:1929–1940. [PubMed: 12612040]
- Jechlinger M, Pelz R, Tretter V, Klausberger T, Sieghart W. Subunit composition and quantitative importance of hetero-oligomeric receptors: GABA_A receptors containing $\alpha 6$ subunits. *The Journal of Neuroscience*. 1998; 18:2449–2457. [PubMed: 9502805]
- Johnson JS, Newport EL. Critical period effects in second language learning: the influence of maturational state on the acquisition of English as a second language. *Cognitive Psychology*. 1989; 21:60–99. [PubMed: 2920538]
- Jones A, Korpi E, McKernan R, Pelz R, Nusser Z, Mäkelä R, Mellor J, Pollard S, Bahn S, Stephenson F, Randall A, Sieghart W, Somogyi P, Smith A, Wisden W. Ligand-gated ion channel subunit partnerships: GABA_A receptor $\alpha 6$ subunit gene inactivation inhibits δ subunit expression. *The Journal of Neuroscience*. 1997; 15:1350–1362. [PubMed: 9006978]
- Kittler J, McAinsh K, Moss S. Mechanisms of GABA_A receptor assembly and trafficking: implications for the modulation of inhibitory neurotransmission. *Molecular Neurobiology*. 2002; 26:251–268. [PubMed: 12428759]
- Laurie DJ, Seeburg PH, Wisden W. The distribution of 13 GABA_A receptor subunit mRNAs in the rat brain. II. Olfactory bulb and cerebellum. *The Journal of Neuroscience*. 1992; 12:1063–1076. [PubMed: 1312132]
- Liang J, Cagetti E, Olsen R, Spigelman I. Altered pharmacology of synaptic and extrasynaptic GABA_A receptors on CA1 hippocampal neurons is consistent with subunit changes in a model of alcohol withdrawal and dependence. *The Journal of Pharmacology and Experimental Therapeutics*. 2004; 310:1234–1245. [PubMed: 15126642]
- Liang J, Suryanarayanan A, Chandra D, Homanics G, Olsen R, Spigelman I. Functional consequences of GABA_A receptor $\alpha 4$ subunit deletion on synaptic and extrasynaptic currents in the mouse dentate granule cells. *Alcohol Clinical and Experimental Research*. 2008; 32:19–26.
- Liang J, Zhang N, Cagetti E, Houser C, Olsen R, Spigelman I. Chronic intermittent ethanol-induced switch of ethanol actions from extrasynaptic to synaptic hippocampal GABA_A receptors. *The Journal of Neuroscience*. 2006; 26:1749–1758. [PubMed: 16467523]
- Maguire J, Stell B, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA_A receptors mediating tonic inhibition alter seizure susceptibility and anxiety. *Nature Neuroscience*. 2005; 8:797–804.
- Maguire J, Ferando I, Simonsen C, Mody I. Excitability changes related to GABA_A receptor plasticity during pregnancy. *The Journal of Neuroscience*. 2009; 29:9592–9601. [PubMed: 19641122]
- Matsuzaki M, Honkura N, Ellis-Davies GC, Kasai H. Structural basis of long-term potentiation in single dendritic spines. *Nature*. 2004; 429:761–766. [PubMed: 15190253]
- McGivern R, Andersen J, Byrd D, Mutter K, Reilly J. Cognitive efficiency on a match to sample task decreases at the onset of puberty in children. *Brain and Cognition*. 2002; 50:73–89. [PubMed: 12372353]
- Meera P, Wallner M, Otis T. Molecular basis for the high THIP/gaboxadol sensitivity of extrasynaptic GABA_A receptors. *The Journal of Neurophysiology*. 2001 July 27.

- Megas M, Emri Z, Freund TF, Gulys AI. Total number and distribution of inhibitory and excitatory synapses on hippocampal CA1 pyramidal cells. *Neuroscience*. 2001; 102:527–540. [PubMed: 11226691]
- Mtchedlishvili Z, Lepsveridze E, Xu H, Kharlamov E, Lu B, Kelly K. Increase of GABA_A receptor-mediated tonic inhibition in dentate granule cells after traumatic brain injury. *Neurobiology of Disease*. 2010; 38:464–475. [PubMed: 20304069]
- Nusser Z, Sieghart W, Stephenson FA, Somogyi P. The $\alpha 6$ subunit of the GABA_A receptor is concentrated in both inhibitory and excitatory synapses on cerebellar granule cells. *The Journal of Neuroscience*. 1996; 16:103–114. [PubMed: 8613776]
- Nusser Z, Sieghart W, Somogyi P. Segregation of different GABA_A receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *The Journal of Neuroscience*. 1998; 18:1693–1703. [PubMed: 9464994]
- Olsen R, Sieghart W. GABA_A receptors: subtypes provide diversity of function and pharmacology. *Neuropharmacology*. 2009; 56:141–148. [PubMed: 18760291]
- Payne H, Ives J, Sieghart W, Thompson C. AMPA and kainate receptors mediate mutually exclusive effects on GABA_A receptor expression in cultured mouse cerebellar granule neurones. *Journal of Neurochemistry*. 2008; 104:173–186. [PubMed: 17986225]
- Peng Z, Hauer B, Mihalek R, Homanics G, Sieghart W, Olsen R, Houser C. GABA_A receptor changes in delta subunit-deficient mice: altered expression of $\alpha 4$ and $\gamma 2$ subunits in the forebrain. *Journal of Comparative Neurology*. 2002; 446:179–197. [PubMed: 11932935]
- Phend KD, Rustioni A, Weinberg RJ. An osmium-free method of epon embedment that preserves both ultrastructure and antigenicity for post-embedding immunocytochemistry. *The Journal of Histochemistry and Cytochemistry*. 1995; 43:283–292. [PubMed: 7532656]
- Pignataro L, Miller A, Ma L, Midha S, Protiva P, Herrera D, Harrison N. Alcohol regulates gene expression in neurons via activation of heat shock factor 1. *The Journal of Neuroscience*. 2007; 27:12957–12966. [PubMed: 18032669]
- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G. GABA_A receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience*. 2000; 101:815–850. [PubMed: 11113332]
- Purdy RH, Morrow AL, Moore PH, Paul SM. Stress-induced elevations of GABA_A receptor-active steroids in the rat brain. *Proceedings of the National Academy of Sciences of the United States of America*. 1991; 88:4553–4557. [PubMed: 1852011]
- Roberts D, Hu Y, Lund I, Brooks-Kayal A, Russerk S. Brain-derived neurotrophic factor (BDNF)-induced synthesis of early growth response factor (Egr3) controls the level of type A GABA receptor $\alpha 4$ subunits in hippocampal neurons. *Journal of Biological Chemistry*. 2006; 281:29431–29435. [PubMed: 16901909]
- Sanna E, Mostallino M, Busonero F, Talani G, Tranquilli S, Mameli M, Spiga S, Follesa P, Biggio G. Changes in GABA_A receptor gene expression associated with selective alterations in receptor function and pharmacology after ethanol withdrawal. *The Journal of Neuroscience*. 2003; 23:11711–11724. [PubMed: 14684873]
- Santhakumar V, Jones R, Mody I. Developmental regulation and neuroprotective effects of striatal tonic GABA_A currents. *Neuroscience*. 2010; 167:644–655. [PubMed: 20206233]
- Shen H, Gong Q, Yuan M, Smith S. Short-term steroid treatment increases δ GABA_A receptor subunit expression in rat CA1 hippocampus: pharmacological and behavioral effects. *Neuropharmacology*. 2005; 49:573–586. [PubMed: 15950994]
- Shen H, Gong Q, Aoki C, Yuan M, Ruderman Y, Dattilo M, Williams K, Smith S. Reversal of neurosteroid effects at $\alpha 4\beta 2\delta$ GABA_A receptors triggers anxiety at puberty. *Nature Neuroscience*. 2007; 10:469–477.
- Shen H, Sabaliauskas N, Sherpa A, Fenton A, Stelzer A, Aoki C, Smith S. A critical role for $\alpha 4\beta \delta$ GABA_A receptors in shaping learning deficits at puberty in mice. *Science*. 2010; 327:1515–1518. [PubMed: 20299596]
- Smith M, Ellis-Davies GCR, Magee J. Mechanism of the distance-dependent scaling of Schaffer collateral synapses in rat CA1 pyramidal neurons. *Journal of Physiology*. 2003; 548:245–258. [PubMed: 12598591]

- Smith SS, Gong QH. Neurosteroid administration and withdrawal alter GABA_A receptor kinetics in CA1 hippocampus of female rats. *Journal of Physiology*. 2005; 564.2:421–436. [PubMed: 15705652]
- Smith SS, Gong QH, Hsu FC, Markowitz RS, French Mullen JM, Li X. GABA_A receptor $\alpha 4$ subunit suppression prevents withdrawal properties of an endogenous steroid. *Nature*. 1998a; 392:926–930. [PubMed: 9582073]
- Smith SS, Gong QH, Li X, Moran M, Bitran D, Frye C, Hsu F-C. Withdrawal from 3 α -OH-5 α -pregnan-20-one using a pseudopregnancy model alters the kinetics of hippocampal GABA_A-gated current and increases the GABA_A receptor $\alpha 4$ subunit in association with increased anxiety. *The Journal of Neuroscience*. 1998b; 18:5275–5284. [PubMed: 9651210]
- Sperk G, Schwarzer C, Tsunashima K, Fuchs K, Sieghart W. GABA_A receptor subunits in the rat hippocampus I: immunocytochemical distribution of 13 subunits. *Neuroscience*. 1997; 80:987–1000. [PubMed: 9284055]
- Staley K, Mody I. Shunting of excitatory input to dentate gyrus granule cells by a depolarizing GABA_A receptor-mediated postsynaptic conductance. *Journal of Neurophysiology*. 1992; 68:197–212. [PubMed: 1381418]
- Staley K, Proctor W. Modulation of mammalian dendritic GABA_A receptor function by the kinetics of Cl[−] and HCO₃[−] transport. *Journal of Physiology*. 1999; 515:693–712. [PubMed: 10457084]
- Stell B, Mody I. Receptors with different affinities mediate phasic and tonic GABA_A conductances in hippocampal neurons. *The Journal of Neuroscience*. 2002; 22:RC223. [PubMed: 12006605]
- Studler B, Fritschy JM, Brnig I. GABAergic and glutamatergic terminals differentially influence the organization of GABAergic synapses in rat cerebellar granule cells *in vitro*. *Neuroscience*. 2002; 114:123–133. [PubMed: 12207960]
- Sundstrom-Poromaa I, Smith D, Gong Q, Sabado T, Li X, Light A, Wiedmann M, Williams K, Smith S. Hormonally regulated $\alpha 4\beta 2\delta$ GABA_A receptors are a target for alcohol. *Nature Neuroscience*. 2002; 5:721–722.
- Sur C, Farrar SJ, Kerby J, Whiting PJ, Atack JR, McKernan RM. Preferential coassembly of $\alpha 4$ and δ subunits of the GABA_A receptor in rat thalamus. *Molecular Pharmacology*. 1999; 56:110–115. [PubMed: 10385690]
- Suryanarayanan A, Liang J, Meyer E, Lindermeyer A, Chandra D, Homanics G, Sieghart, Olsen R, Spigelman I. Subunit compensation and plasticity of synaptic GABA_A receptors induced by ethanol in $\alpha 4$ subunit knockout mice. *Frontiers in Neuroscience*. 2011; Volume 5 Article 110.
- Wei W, Faria L, Mody I. Low ethanol concentrations selectively augment the tonic inhibition mediated by δ subunit-containing GABA_A receptors in hippocampal neurons. *The Journal of Neuroscience*. 2004; 24:8379–8382. [PubMed: 15385620]
- Wei W, Zhang N, Peng Z, Houser C, Mody I. Perisynaptic localization of δ subunit-containing GABA_A receptors and their activation by GABA spillover in the mouse dentate gyrus. *The Journal of Neuroscience*. 2003; 23:10650–10661. [PubMed: 14627650]
- Wisden W, Laurie D, Monyer H, Seeburg P. The distribution of 13 GABA_A receptor subunit mRNAs in the rat brain. I. telencephalon, diencephalon, mesencephalon. *The Journal of Neuroscience*. 1992; 12:1040–1062. [PubMed: 1312131]
- Wohlfarth K, Bianchi M, Macdonald R. Enhanced neurosteroid potentiation of ternary GABA_A receptors containing the δ subunit. *The Journal of Neuroscience*. 2002; 22:1541–1549. [PubMed: 11880484]
- Yuste R, Denk W. Dendritic spines as basic functional units of neuronal integration. *Nature*. 1995; 375:682–684. [PubMed: 7791901]
- Zhou X, Smith S. Steroid requirements for regulation of the $\alpha 4$ subunit of the GABA_A receptor in an *in vitro* model. *Neuroscience Letters*. 2007; 411:61–66. [PubMed: 17081691]

Figure 1. Light microscopic immunohistochemistry of $\alpha 4$ labeling in WT and $\alpha 4$ KO hippocampus

Light micrographs of hippocampal sections from WT and $\alpha 4$ KO pubertal female mice were captured following immunocytochemical labeling for the $\alpha 4$ subunit, using SIG as the label. Expression of the $\alpha 4$ subunit was apparent both in the pyramidal cell layer (PCL; black arrowhead) as well as in hippocampal interneurons (white arrowhead) of the stratum oriens (SO) and stratum radiatum (SR) of WT tissue (panel A). Qualitative analysis revealed marked reduction of the $\alpha 4$ subunit in $\alpha 4$ KO mice at puberty (panel B) when compared to the tissue of WT pubertal mice. Scale bar=500 μ m.

Figure 2. Plasmalemmal $\alpha 4$ immunoreactivity on pubertal CA1 hippocampal pyramidal cells' dendritic spines and shafts is reduced by $\alpha 4$ KO

Representative EM images of dendritic spine profiles from WT pubertal (panel A) and $\alpha 4$ KO pubertal (panel B) CA1 regions following SIG-immunocytochemistry to label $\alpha 4$ subunits. The black arrow denotes plasmalemmal SIG. 'U' denotes unlabeled dendritic spine profiles, while 'L' denotes a labeled dendritic spine profile. Scale bar for panels A and B=500 nm. Mann-Whitney U-test of plasmalemmal $\alpha 4$ labeling revealed a significant effect of genotype, with significantly greater $\alpha 4$ immunoreactivity within WT pubertal tissue, compared to $\alpha 4$ KO pubertal tissue (panel C; * $p < 0.0001$). $n = 1400$ spines, sampled equally from 5 WT pubertal and 5 KO pubertal mice. Representative EM images of dendritic shafts from WT pubertal (panel D) and $\alpha 4$ KO pubertal (panel E) mice. The black arrow denotes plasmalemmal SIG. 'Sh' denotes dendritic shafts. Scale bar for panels C and D=500 nm. Mann-Whitney U-test plasmalemmal $\alpha 4$ labeling revealed a significant effect of genotype, with significantly greater $\alpha 4$ immunoreactivity within WT pubertal tissue, compared to $\alpha 4$ KO pubertal tissue (panel F; ** $p < 0.000001$). $n = 200$ dendrites sampled equally from 5 WT pubertal and 5 KO pubertal mice. Values indicate means $\pm$ SEM.

Figure 3. CA1 pyramidal cells exhibit a normal neuronal phenotype: A comparison of WT and $\alpha 4$ KO pubertal mice

Light microscopy reveals normal expression levels of two neuronal markers. Expression levels of the AMPA receptor subunit GluR2/3 are indistinguishable between WT pubertal (panel A) and $\alpha 4$ KO pubertal (panel B) tissue. Pubertal tissue of WT (panel C) and $\alpha 4$ KO (panel D) mice also showed similar immunoreactivity to the neuronal marker MAP2. The CA1 hippocampus from both genotypes also exhibits proper pyramidal cell morphology and stratification. Scale bar for panels A–D=500 μ m.

Figure 4. Plasmalemmal δ immunoreactivity on pubertal CA1 hippocampal pyramidal cells' dendritic spines and shafts is reduced by $\alpha 4$ KO

Representative EM images of dendritic spine profiles from WT pubertal (panel A) and $\alpha 4$ KO pubertal (panel B) CA1 regions following SIG-immunocytochemistry to label δ subunits. The black arrow denotes plasmalemmal SIG. 'U' denotes unlabeled dendritic spine profiles, while 'L' denotes a labeled dendritic spine profile. Scale bar for panels A and B=500 nm. Mann-Whitney U-test of plasmalemmal δ labeling revealed a significant effect of genotype, with significantly greater δ immunoreactivity within WT pubertal tissue, compared to $\alpha 4$ KO pubertal tissue (panel C; * $p < 0.0001$). $n = 1400$ spines, sampled equally from 5 WT pubertal and 5 KO pubertal mice. Representative EM images of dendritic shafts from WT pubertal (panel D) and $\alpha 4$ KO pubertal (panel E) mice. The black arrow denotes plasmalemmal SIG. 'Sh' denotes dendritic shafts. Scale bar for panels C and D=500 nm. Mann-Whitney U-test of plasmalemmal δ labeling revealed a significant effect of genotype, with significantly greater δ immunoreactivity within WT pubertal tissue, compared to $\alpha 4$ KO pubertal tissue (panel F; * $p < 0.0001$). $n = 200$ dendrites sampled equally from 5 WT pubertal and 5 KO pubertal mice. Values indicate means $\pm$ SEM.

Figure 5. Reduced gaboxadol responsiveness of $\alpha 4$ KO pubertal mouse hippocampus as compared to WT pubertal mouse

The GABAR agonist gaboxadol (GBX) is selective for δ -containing GABAR at a concentration of 100 nM, while gabazine (GBZ) is an antagonist specific to all GABARs at a concentration of 120 μ M. Representative currents (panel A) and averaged population data (panel B) reveal that the response to GBX was greater at puberty in WT hippocampus than in $\alpha 4$ KO hippocampus. The response to 120 μ M GBZ is the current deflection after treatment with GBX. (* $p < 0.05$), $n = 5$ cells per group. Scale bar = 50 pA, 50 s.

Table 1

<i>δ IMMUNOREACTIVITY</i>		WT Mean ± SEM (n)	KO Mean ± SEM (n)	p-value* (Z)
SPINES	<i>Surface (percent labeled)</i>	5.16 ± 0.83 (700)	2.79 ± 0.70 (700)	0.00008 (3.955)
	<i>Intracellular (percent labeled)</i>	4.92 ± 0.86 (700)	4.22 ± 0.91 (700)	0.5 (−0.629)
	<i>Surface proportion of total labeling (percent)</i>	59.9 ± 6.5 (54)	36.3 ± 7.0 (45)	0.03 (2.091)
SHAFTS	<i>Surface (label per μm)</i>	0.097 ± 0.011(100)	0.042 ± 0.009(100)	0.00006 (−4.402)
	<i>Intracellular (label per μm²)</i>	1.151 ± 0.085(100)	1.021 ± 0.070(100)	0.5 (−0.715)
	<i>Surface proportion of total labeling (percent)</i>	0.147 ± 0.016(100)	0.047 ± 0.011(100)	0.00001 (−4.921)

* Bold numbers indicate significant p-values

Table 2

<i>PYRAMIDAL CELL PARAMETERS</i>		WT Mean ± SEM (n)	KO Mean ± SEM (n)	p-value* (Z)
SPINES	<i>Spine area (nm²)</i>	64319 ± 1743 (500)	72308 ± 1796 (500)	0.00005 (4.080)
	<i>Spine membrane (nm)</i>	1012 ± 15 (500)	1070 ± 15 (500)	0.001 (3.287)
SHAFTS	<i>Diameter (nm)</i>	1504 ± 35 (100)	1609 ± 42 (100)	0.05 (−1.985)

*
Bold numbers indicate significant p-values