Published in final edited form as:

Public Health Nurs. 2011 ; 28(6): 494–502. doi:10.1111/j.1525-1446.2011.00945.x.

Intention to Smoke Tobacco Using Waterpipe among Students in a Southeastern U.S. College

Devon Noonan, PhD, MPH, FNP-BC[Research Fellow],

Health Promotion/ Risk Reduction Interventions with Vulnerable Populations, University of Michigan School of Nursing, 400 North Ingalls, Ann Arbor, MI, 48109-5482

Pamela Kulbok, RN, DNSc, PHCNS-BC[Professor of Nursing], and

Coordinator of Community & Public Health Leadership, University of Virginia, 225 Jeanette Lancaster Way, Charlottesville, VA 22908

Guofen Yan, PhD[Assistant Professor]

Biostatistics & Epidemiology, University of Virginia, P.O. Box 800717, Charlottesville, VA 22908-0717

Devon Noonan: dnoonan@umich.edu

Abstract

Objective—Guided by the Theory of Reasoned Action this study examined the association of behavioral beliefs, attitudes, normative beliefs and subjective norms with waterpipe tobacco smoking intention in college students.

Design and Sample—A cross-sectional design was used. A web-based survey was sent to a random sample of 1,000 undergraduate students from a public institution in the Southeast to recruit participants.

Measures—The Theory of Reasoned Action Waterpipe Questionnaire, a modified version of the Fishbein-Ajzen-Hanson Questionnaire, was used to capture modal constructs of the Theory of Reasoned Action related to waterpipe use. Cronbach's alpha coefficients for the scales of the Theory of Reasoned Action Waterpipe Questionnaire ranged from .76 to .95.

Results—Of the sample ($n=223$), 13.5% currently smoked a waterpipe and 61% had ever done so. Using multiple regression, attitudes, behavioral beliefs, and subjective norms were associated with intention to smoke a waterpipe in the next three months and collectively explained 35% of the variance in intention. The full model, which included all the constructs of the Theory of Reasoned Action, demographic variables, and tobacco use variables, explained 83% of the variance in intention to smoke a waterpipe in the next three months.

Conclusions—This study provides valuable information that may be used to target students at risk for waterpipe smoking and serves as a starting point in developing theoretically-driven interventions to prevent waterpipe smoking.

Keywords

health promotion; health risk behaviors; tobacco; waterpipe

Tobacco use is the leading preventable cause of death in the United States (U.S.) and is responsible for an estimated 443,000 deaths, or 1 out of 5 deaths annually (Centers for Disease Control and Prevention [CDC], 2008). There are many forms of tobacco use, all of which contain nicotine and are highly addictive. Waterpipe smoking has gained attention as a new trend in tobacco use among U.S. college students and young adults, and the American Lung Association (ALA) has labeled it the first new tobacco trend of the 21st century

(2007). In addition, the worldwide increase in smoking tobacco using a waterpipe among young people during the past decade may represent the second global tobacco use epidemic since cigarettes (Maziak, 2010). The purpose of this study, guided by the Theory of Reasoned Action (Fishbein & Ajzen, 1975), was to examine predictors of waterpipe tobacco smoking intention in U.S. college students. Theory and evidence-based knowledge related to waterpipe smoking attitudes, beliefs, and behavior may be a useful starting point for developing and testing future preventive interventions.

Background and Literature

Waterpipe smoking is generally defined as a tobacco use method in which smoke passes through water before it is inhaled (Maziak, Ward, Afifi, Soweid, & Eissenberg, 2004). A typical waterpipe contains five parts: a bowl, head, base, hose, and mouthpiece. The tobacco used to smoke a waterpipe does not burn in a self-sustaining manner, so charcoal is placed in the bowl on top of the tobacco, although the coal is usually separated from the tobacco by a piece of perforated aluminum foil (World Health Organization [WHO], 2005). Many people believe that waterpipe smoking is a safe alternative to the use of conventional tobacco products. However, in a recent study by Eissenberg and Shihadeh (2009) comparing waterpipe smoke toxicant exposure to cigarette smoke toxicant exposure, waterpipe smoke was found to contain comparable peak nicotine levels and three times greater peak Carboxyhemoglobin (COHb) levels (a measure of carbon monoxide exposure). Furthermore, because of the long duration of most waterpipe sessions (45-minutes or more) waterpipe smoking results in much greater nicotine exposure, 1.7 times that of smoking a cigarette. Waterpipe smoke contains high levels of other toxic compounds including heavy metals and cancer causing chemicals, and therefore may be associated with multiple health risks including nicotine addiction, cancer of the lungs, and heart disease (Jabbuor, El-Roueiheb, & Sibai, 2003; Eissenberg & Shihadeh, 2009; Sepetdjian, Shihadeh & Saliba, 2008; Shihadeh & Saleh, 2005). Recent research highlighted the danger of secondhand waterpipe smoke, and yet waterpipe smoking is able to circumvent most smoke-free indoor air laws (Daher et al., 2010; Maziak, Ibrahim, Rastam, Ward & Eissenberg, 2008; Noonan, 2010a).

Current research has highlighted growing waterpipe use on college campuses. In a cross-sectional study of U.S. college students, Grekin and Ayna (2008) reported “current” waterpipe use, defined as having smoked a waterpipe in the last 30 days, to be approximately 15 % ($n=602$). Other studies support the 15–20% estimate of waterpipe use among college students and young adults in the U.S. (Eissenberg, Ward, Smith-Simone, & Maziak, 2008; Smith-Simone, Maziak, Ward, & Eissenberg, 2008; Ward, Eissenberg, Gray, Srinivas, Wilson, & Maziak 2008). Waterpipe smoking in college student populations has been strongly linked to cigarette use (Dugas et al., 2010; Primack et al., 2008; Smith-Simone et al., 2008). Smith-Simone and colleagues (2008) found that 54% of students that used waterpipe currently also currently smoked cigarettes. Similarly, Dugas and colleagues (2010) found that young adults who had used waterpipe in the past year were more likely to have smoked cigarettes in the past year. Jensen and colleagues (2010) reported that waterpipe smoking among male adolescents was predictive of cigarette smoking. Furthermore, students who previously experimented with waterpipe smoking are also more likely to have intentions to smoke waterpipe in the future (Primack et al., 2008).

Waterpipe smoking has been associated with demographic variables including age, gender, and race. In a study of 744 U.S. college students, Eissenberg and colleagues (2008) reported past year waterpipe smoking to be associated with being male, being younger (18–19 vs. 20–23), and being Caucasian American. Dugas and colleagues (2010) found similar results with younger age and male gender also predicting waterpipe smoking. Researchers have also examined, though to a lesser degree, psychosocial factors associated with waterpipe

smoking in college populations. Smith-Simone and colleagues (2008) assessed attitudes towards waterpipe smoking in a convenience sample of 201 users and reported common positive attitudes to include pleasant taste, pleasant smell, relaxing effects, and the opportunity to socialize with friends. Similarly, Primack and colleagues (2008) examined attitudes, norms and intentions towards waterpipe smoking among 647 undergraduate and graduate college students and found that one-third of the sample considered waterpipe smoking socially acceptable and over half viewed waterpipe smoking as less dangerous than cigarette smoking. Approximately 87% of waterpipe users and 20% of non-users had intentions to smoke a waterpipe in the future. Notably, very few of these studies used a theoretical framework to examine psychosocial factors associated with waterpipe use in college students, although this information is necessary to design effective interventions for this population.

The Theory of Reasoned Action (Fishbein & Ajzen, 1975) was used to guide this study and has been used successfully in predicting smoking intentions in past tobacco use studies (Hanson 1997, 2006; Primack et al., 2008). According to the Theory of Reasoned Action, human action is determined by behavioral intention. Attitudes and subjective norms related to a behavior, determine behavioral intention (Fishbein & Ajzen, 1975). Behavioral beliefs and normative beliefs surrounding a behavior determine attitudes and subjective norms, respectively. Behavioral beliefs and the outcome evaluation of these beliefs produce a favorable or unfavorable attitude towards a particular behavior and in turn determine attitudes surrounding a behavior. Normative beliefs and the motivation to comply with those beliefs produce perceived subjective norms and in turn determine subjective norms surrounding a behavior. Behavioral and normative beliefs are often referred to as indirect measures, therefore indirectly affecting behavioral intentions.

The purpose of this study was to determine if the constructs of the Theory of Reasoned Action (behavioral beliefs, attitudes, normative beliefs and subjective norms) were associated with waterpipe smoking intention in the next three months. For this study each of the aforementioned constructs of the Theory of Reasoned Action were examined individually for their association with waterpipe smoking intention. The study also examined the relationship of waterpipe smoking intention with demographics and tobacco use. The interaction of gender and school year with the four constructs of the Theory of Reasoned Action was also explored to identify target groups for prevention efforts.

Methods

Design, Setting, and Sample

This study used a cross-sectional design. Participants were recruited from a public institution in a Southeastern state with approximately 13,000 undergraduate students: 55% female 45% male; 64% Caucasian American, 11% Asian American, 8% African American, 4% Hispanic, 0.2% Native American, and 12% other/unclassified (Office of Institutional Assessment, 2006). The Institutional Review Board (IRB) at the university approved the study. This study was conducted in the Spring Semester of 2009. The Office of Student Services provided a computer-generated simple random sample of 1,000 undergraduate students as a pool from which to recruit participants.

To improve response rates, the random sample of 1,000 undergraduates received a pre-notice email informing them that they would receive an email invitation in the next few days to participate in a survey about waterpipe smoking (Dillman, 2000). Three days later, the sample received an email inviting them to participate in the waterpipe survey via a link provided. It was made clear in the invitation that only tobacco waterpipe smoking was of interest. SurveyGizmo, online-computerized survey software, was used to recruit

participants and manage the survey. By agreeing to participate in the study based on the narrative provided, students gave implicit consent for participation. Participants received privacy information for SurveyGizmo.com with the consent information. The survey was open for 2 weeks with reminders sent every 3 days to encourage participation. In an effort to encourage participation, respondents who completed the survey were eligible to enter a lottery drawing of a gift certificate for \$250 to a well known bookstore.

Measures

A 45-item online survey was used to gather information on demographic variables, tobacco use behavior, and variables measuring the four constructs of the Theory of Reasoned Action. Demographic information included age, gender, racial and/or ethnic identity, and year in school. Current (“During the past thirty days have you tried smoking tobacco in a waterpipe even one or two puffs?” and “In the past 30 days how many days did you smoke tobacco using a waterpipe”) and ever waterpipe smoking (“Have you ever tried smoking tobacco in a waterpipe, even one or two puffs?”) was collected. “Smoking tobacco” in these items was bolded to avoid confusion with other substances. Current and ever cigarette data was also collected. The cigarette use items were from the 2007 Youth Risk Behavior Survey (CDC, 2007). The waterpipe measures were from Smith’s College Freshman Nicotine Study (Smith, 2006).

A 37-item investigator-developed Theory of Reasoned Action Waterpipe Questionnaire assessed psychosocial factors related to waterpipe smoking intention in college students. The questionnaire was a modified version of the Fishbein-Ajzen-Hanson Questionnaire (FAHQ) (Hanson, 1997). The original Fishbein-Ajzen-Hanson Questionnaire is a 50-item questionnaire based on the Theory of Reasoned Action/Theory of Planned Behavior (TPB). This questionnaire included scales to measure the psychological determinants of cigarette smoking. Seven direct measures including intention, attitude and subjective norm were adapted from the Fishbein-Ajzen-Hanson Questionnaire, with waterpipe use substituted for cigarette use. All direct measures were assessed using 7-point semantic differential evaluative scales. Intentions were captured by three items, “I intend to smoke tobacco using a waterpipe three months from now...” (true/false, likely/unlikely, and probably/probably not). Attitudes were captured by three items, “For me to smoke tobacco using a waterpipe in the next three months would be...” (pleasant/not-pleasant, nice/awful, and fun/not fun). Subjective norm was captured by one item “If I smoke tobacco using a waterpipe most people that are important to me would...” (approve/disapprove).

The indirect measures of the Fishbein-Ajzen-Hanson Questionnaire assessed beliefs and referents salient to cigarette smoking and were not useful for this study. Therefore, a pilot study was conducted to gather necessary information to create belief-based, indirect measures of attitudes and subjective norms related to waterpipe use among college students. The pilot study used free response questions suggested by Ajzen (2002) to obtain salient beliefs and referents of waterpipe smoking in 58 college students. Content analysis was performed and salient behavioral beliefs that emerged from the analysis included, “waterpipe smoking is... safer than regular cigarettes, causes lung cancer, may harm my health, is less irritating than cigarette smoke, gives me a good buzz, costs a lot of money, allows me to have a good time with friends, is relaxing, tastes good, and smells good.” These beliefs were used to create the behavioral belief and outcome evaluation scales in relation to waterpipe use. A total of twenty, 7- point semantic differential scales were used, ten for modal belief items ranging from “likely to unlikely” and ten outcome evaluation items ranging from desirable to undesirable” were captured. Referents that emerged (family, friends, significant others, parents and siblings) were used to create the normative belief and motivation to comply scales in relation to waterpipe use. A total of ten 7-point semantic differential scales were used, five perceptions of referents items ranging from “approve to

disapprove” and five motivation to comply items ranging from “agree to disagree” were captured. These indirect measures of attitude and subjective norms were modeled after the belief-based measures of the FAHQ. The scoring of the Theory of Reasoned Action Waterpipe Questionnaire followed that of the Fishbein-Ajzen-Hanson-Questionnaire (Hanson, 1997, 2006).

To test the feasibility of the questionnaire and the reliability of the scales, a second pilot study was conducted with a convenience sample of 100 undergraduate college students to examine internal consistency using Cronbach’s alpha reliability coefficients. Cronbach’s alpha for the entire 37-item Theory of Reasoned Action Waterpipe Questionnaire was 0.86. Cronbach’s alpha coefficients for the scales of the Theory of Reasoned Action Waterpipe Questionnaire ranged from .76 to .95. The intention scale had the highest reliability (.95) followed by attitude (.94), behavioral beliefs (.76), and normative beliefs (.76) respectively.

Data Analysis

Multiple linear regression was used to examine the relationship between intention to smoke waterpipe in the next three months and the independent variables of behavioral beliefs, attitude, normative beliefs, and subjective norms. The relationships of age, gender, year in school, race, current waterpipe and cigarette use, with intention to smoke waterpipe, were also examined based on associations reported in the current literature (Dugas et al., 2010; Eissenberg et al., 2008; Smith-Simone et al., 2007; Ward et al., 2008). First, the regression model examined the relationship between intention to smoke and each of the four individual constructs (behavioral beliefs, attitude, normative beliefs, and subjective norms controlling for age, gender, year in school, race, current cigarette use, and current waterpipe use. Second, the regression model was used to examine the joint relationship between intention to smoke a waterpipe and the four Theory of Reasoned Action constructs (entered into the model simultaneously), controlling for age, gender, year in school, race, current cigarette use, and current waterpipe use. The regression models that included the interaction terms of gender and each individual construct of the Theory of Reasoned Action as well as the interaction terms of school year and each individual construct of the Theory of Reasoned Action were then examined. Model assumptions were checked including normality, linearity, multicollinearity, and homoscedasticity with no major violations noted. Statistically significant results were interpreted at a p value of 0.05.

Results

Of the 1000 emails sent representing the target population, only seven were undeliverable. Of the 993 college students from the target population who received the survey invitation, 261 (26%) completed the survey. Of the 261 students who completed the survey, two cases had a calculated Mahalanobis distance above the critical threshold indicating they contained multivariate outliers and were deleted to reduce the chances of error (Meyers, Gamst, & Guarino, 2006). Another 36 participants were eliminated from the analysis because of incomplete questionnaires resulting in a final sample of 223 participants. Two sample t -tests and chi-square tests were used to compare the demographic and tobacco use variables of the 36 cases that were eliminated and the 223 cases that were included in the study. The results of this analysis revealed no significant differences between non-respondents and the final study sample.

Demographics and Tobacco Use Behavior

The mean age (SD) of the sample was 19.9 years (1.3); more females (54%) than males completed the survey. The majority were Caucasian American. This is consistent with the demographic characteristics of the university students sampled. Sixty-one percent reported

ever waterpipe use and 13.5% reported current waterpipe use. Fifty percent of the sample reported ever cigarette use and 18% reported current cigarette use (see Table 1).

Correlates of Waterpipe Smoking Intention

Each construct of the Theory of Reasoned Action was found to be significantly associated with intention to smoke waterpipe in the next three months, after adjusting for demographic variables and tobacco behaviors (attitude: $B = .250$, $p < .001$, behavioral beliefs: $B = .037$, $p < .001$, subjective norm: $B = .657$, $p < .001$, and normative beliefs: $B = .031$, $p < .001$). The amount of variance explained for each construct ranged from 22% to 75% with attitude accounting for the most variance (75%), followed by behavioral beliefs (43%), normative beliefs (34%) and subjective norms (21%). As expected, the relationships were positive; that is, smoking intention increased as favorable attitudes, perceived subjective norms, and behavioral beliefs about waterpipe smoking increased.

When all four constructs of the Theory Reasoned Action, demographics, (age, gender, race, year in school) and tobacco use (current waterpipe, and current cigarette use) were simultaneously examined for their joint relationship with waterpipe smoking intention, these variables collectively accounted for 83% of the variance in intention. Attitude, behavioral beliefs, and subjective norm were all significantly associated with intention and explained 35% of the variance in waterpipe smoking intention. Table 2 presents the regression coefficients, significance levels and R^2 for all of the variables entered into the model. Current cigarette use, current waterpipe use, and school year were also significantly associated with intention, with second and third year students having stronger intentions to smoke waterpipe in the next 3 months than fourth year students.

The interactions of gender and school year with the Theory of Reasoned Action constructs were not significant, indicating that the effects of attitudes, norms, behavioral beliefs, and normative beliefs on intention did not differ by gender or year in school.

Discussion

This is one of the first theoretically-driven studies to examine waterpipe smoking intentions among college students. Furthermore, in the available literature examining smoking tobacco using a waterpipe, very few studies use measures based on the Theory of Reasoned Action to guide their analysis. Theoretically-driven studies are necessary to design interventions and tailor health education messages geared toward waterpipe tobacco use (Kasprzyk, Montano, & Fishbein, 1998). In this study, when examined individually, each of the Theory of Reasoned Action constructs were significantly associated with waterpipe smoking intention, while controlling for demographic and tobacco use variables. Attitudes towards waterpipe smoking explained the most variance (75%) in intention to smoke tobacco using a waterpipe in the next three months. In the full model, the Theory of Reasoned action constructs of attitude, behavioral beliefs, and subjective norms remained significantly associated with intention to smoke in this study. In the full model, attitudes towards waterpipe smoking accounted for the most variance (33%) in intention to smoke at three months. This finding suggests that future research developing and testing prevention messages that focus on decreasing favorable attitudes towards waterpipe smoking is warranted. These findings are similar to those reported in other studies conducted in the U.S. where positive attitudes towards smoking were associated with use in the college population (Primack et al., 2008; Smith-Simone et al., 2008). However, the majority of these studies were not theoretically based. Therefore, this study serves as a strong foundation for future theory-based studies examining attitudes towards waterpipe smoking and their effect on intentions and behavior.

Although behavior beliefs and subjective norms were significant, they accounted for very little variance in intention to smoke tobacco using a waterpipe in the full model. It is noteworthy that subjective norms did not account for a large proportion of variance in the model. Many other U.S. based studies have reported subjective norms (perceived social acceptability of waterpipe smoking) as a strong indicator of waterpipe use in this population (Eissenberg et al., 2008; Primak et al., 2008; Smith-Simone et al., 2008). This relationship should be re-examined in future studies with larger more diverse samples.

The demographic variables of age, gender, and race have been shown to be associated with waterpipe smoking in the literature (Dugas et al., 2010; Eissenberg et al., 2008; Smith-Simone et al., 2007; Ward et al., 2008), although they were not associated with waterpipe smoking intention after adjusting for the Theory of Reasoned Action constructs. The demographic variables (age, gender, and race) were examined individually for their association with intention and only sex was found to be a significant predictor; male sex was associated with stronger intentions to smoke. Furthermore, the interaction of the constructs of the Theory of Reasoned Action with gender and school year were not significant, suggesting that broad, campus-wide antismoking messages that target attitudes, subjective norms, and behavioral beliefs surrounding waterpipe use may be beneficial to all students, regardless of gender or year in school. This interpretation of the results needs to be tested in larger, more diverse populations.

This study highlights the problem of waterpipe smoking among college students and young adults. In this study, current waterpipe use (defined as use in the last 30 days) was 13.5%. This is similar to current rates reported by Smith (2006) and by Grekin and Anya (2008), and is consistent with the data from two previous pilot studies examining waterpipe smoking in a similar university setting (Noonan, 2010b). Sixty-one percent of this sample had ever smoked a waterpipe. This percentage was higher than seen in similar studies, which reported ever waterpipe use between 40 and 50% (Eissenberg, Ward, Smith-Simone, & Maziak, 2008; Primack et al., 2008). This finding may reflect the increasing popularity of waterpipe use over the past two years, and is a cause for concern. Moreover, more students had ever smoked a waterpipe (61%) than had ever smoked cigarettes (49.8%), suggesting that waterpipe smoking may be more appealing to tobacco naïve persons. However, because current cigarette and waterpipe use were strongly associated with smoking intentions, explaining 29% of the variance in intention to smoke in the next three months, those experienced with other forms of tobacco use, i.e., smoking cigarettes, should also be targeted for waterpipe prevention interventions. This supports similar studies where traditional forms of tobacco use were also a stronger predictor of waterpipe smoking (Dugas et al., 2010; Smith-Simone et al., 2008). Further investigation examining the relationship between waterpipe smoking intentions and multiple different tobacco use methods are necessary to target appropriately those at risk for smoking waterpipe in the future.

Study Limitations

While this study provides important information about factors predicting waterpipe smoking intention in college students, there were some notable limitations. The response rate was low (26%), although this is common in college student populations and for email surveys. Response rates for college students average between 17% and 70% (Porter & Umbach, 2006). However, exclusion of subjects who did not respond could introduce bias. In addition, the sample was from a single university in the Southeast and the majority of students (71%) were Caucasian American. Although the racial distribution in the study was similar to that of the university, future studies should focus on minority recruitment by over sampling to diversify the study population. In addition, future studies may include a more geographically diverse samples obtained from multiple colleges and universities. It should also be noted, since this was a cross-sectional observational study, that causality cannot be

established between the independent and dependent variables. Therefore, the results of this study should be interpreted with caution. Future theory-based research of this nature should employ a longitudinal approach. Finally, the measures used for this study are new and although they were developed from preliminary work in accordance with Ajzen's (2002) recommendations, they may not fully capture the theoretical constructs. Further testing and refinement of the Theory of Reasoned Action Waterpipe Questionnaire is recommended.

Implications for Practice

Healthcare practitioners often play a large role in designing college campus prevention strategies and can use the results of this study to inform current tobacco and waterpipe prevention messages for students in similar university settings. Notably, the results of this study coupled with the available current literature indicate that waterpipe smoking and use of other tobacco products are present on many college campuses. Healthcare providers working with college students need to recognize this and screen patients accordingly. Individually, providers can capitalize on opportunities to promote the dangers of waterpipe smoking during patient visits and help to dispel myths about using a waterpipe to smoke tobacco. For example, the myth that smoking tobacco using a waterpipe is safer than using other tobacco products. Providers should highlight the unpleasant and serious aspects of waterpipe smoking, including nicotine addiction, various chronic health effects and communicable diseases. Providers can suggest alternate, healthier ways to socialize with friends, which do not involve addictive substances (like substance-free campus sponsored events) and refer patients for cessation services as necessary. Providers can also empower students to disapprove of their peer's waterpipe smoking to challenge campus norms surrounding the acceptability of waterpipe smoking, as well as teach students peer pressure resistance techniques. Finally, providers should advocate for smoke-free environments for all students by promoting a tobacco-free campus.

Conclusions

Waterpipe smoking has recently become a new trend among college students in the U.S. and is associated with multiple health problems, including addiction (ALA, 2007). Moreover, waterpipe smoking is recognized as a public health threat in the U.S. and worldwide, due in part to its growing popularity among young people (Cobb, Ward, Maziak, Shihedeh, & Eissenberg, 2010; Maziak, 2010). College students ages 18 to 24 represent a growing vulnerable cohort of individuals that are in need of targeted interventions, intensifying the need for new, theoretically-driven prevention initiatives to prevent tobacco use in this population (IOM, 2007). Developing and testing theory- and evidence-based interventions and programs to prevent waterpipe smoking, which are lacking in the literature, are critical from a public health perspective to curtail this new potential pathway for nicotine addiction before it evolves further.

References

- Ajzen, I. Constructing a TpB questionnaire: conceptual and methodological considerations. 2002. Retrieved May 1, 2007 from <http://www.people.umass.edu/ajzen/pdf/tpb.measurement.pdf>
- American Lung Association. An emerging deadly trend: waterpipe tobacco use. Tobacco Policy Trend Alert. 2007. Retrieved April 1, 2007 from http://www.lungusa2.org/embargo/slati/Trendalert_Waterpipes.pdf
- Bonnie, R.; Stratton, K.; Wallace, R., editors. Institute of Medicine Ending the tobacco problem: a blueprint for the nation. Washington, DC: National Academy Press; 2007.
- Centers for Disease Control and Prevention. Cigarette smoking among adults-United States, 2004. Morbidity and Mortality Weekly Report. 2005; 54(44):1121-1124. [PubMed: 16280969]

- Centers for Disease Control and Prevention. Tobacco use among adults: United States, 2005. Morbidity and Mortality Weekly Report. 2006; 55(42):1145–1148. [PubMed: 17065979]
- Centers for Disease Control and Prevention. Youth Risk Behavior Survey. 2007. Retrieved from <http://cdc.gov/yrbss>
- Centers for Disease Control and Prevention. Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses—United States, 2000–2004. Morbidity and Mortality Weekly Report. 2008; 57(45):1226–8. [PubMed: 19008791]
- Cobb C, Ward KD, Maziak W, Shihadeh AL, Eissenberg T. Waterpipe tobacco smoking: an emerging health crisis in the United States. American Journal of Health Behavior. 2010; 34(3):275–85. [PubMed: 20001185]
- Daher A, Saleh R, Jaroudi E, sheheiti Badr, Sepetdjian E, Al Rashidi M, Saliba N, Shihadeh A. Comparison of carcinogen, carbon monoxide, and ultrafine particle Emissions from narghile waterpipe and cigarette smoking: sidestream smoke Measurements and assessment of second-hand smoke emission factors. Atmospheric Environment. 2010; 44:8–14.10.1016/j.atmosenv.2009.10.004 [PubMed: 20161525]
- Dillman, D. Mail and Internet Surveys. 2. New Jersey: John Wiley and Sons; 2007.
- Dugas E, Tremblay M, Low NCP, Cournoyer D, O'Loughlin J. Water-pipe smoking among North American youths. Pediatrics. 2010; 125:1184–1189.10.1542/peds.2009-2335 [PubMed: 20457679]
- Eissenberg T, Shihadeh A. Waterpipe tobacco and cigarette smoking direct comparison of toxicant exposure. American Journal of Preventive Medicine. 2009; 37(6):518–523.10.1016/j.amepre.2009.07.014 [PubMed: 19944918]
- Eissenberg T, Ward KD, Smith-Simone S, Maziak W. Waterpipe tobacco smoking on a U.S. college campus: prevalence and correlates. Journal of Adolescent Health. 2008; 42(5):526–529.10.1016/j.jadohealth.2007.10.004 [PubMed: 18407049]
- Fishbein, M.; Ajzen, I. Belief, attitude, intention and behavior: An introduction to theory and research. Reading, Massachusetts: Addison-Wesley Publishing; 1975.
- Grekin ER, Ayna D. Argileh use among college students in the United States: an emerging trend. Journal of Studies on Alcohol and Drugs. 2008; 69(3):472–475. [PubMed: 18432392]
- Hanson M. The theory of planned behavior applied to cigarette smoking in African-American, Puerto Rican, and non-Hispanic white teenage females. Nursing Research. 1997; 46(3):155–162. [PubMed: 9176505]
- Hanson M. Predicting smoking behavior in college students. The American Journal for Nurse Practitioners. 2006; 10(6):26–30.
- Jabbour S, El-Roueiheb Z, Sibai AM. Narghile (waterpipe) smoking and incident coronary heart disease: a case control study. {abstract}. Annals of Epidemiology. 2003; 13:570.
- Jensen PD, Cortes R, Engholm G, Kremers S, Gislum M. Waterpipe use predicts progression to regular cigarette smoking among Danish youth. Substance Use and Misuse. 2010; 45:1245–1261.10.3109/10826081003682909 [PubMed: 20441461]
- Kasprzyk D, Montano DE, Fishbein M. Application of an intergrated behavioral model to predict condom use: a prospective study among high HIV risk groups. Journal of Applied Sociology. 1998; 28(17):1557–1583.10.1111/j.1559-1816.1998.tb01690.x
- Maziak W. The global epidemic of waterpipe smoking. Addictive Behaviors. 2010a; 6:1–5.
- Maziak W, Ward KD, Afifi Soweid RA, Eissenberg T. Tobacco smoking using a waterpipe: A re-emerging strain in a global epidemic. Tobacco Control. 2004a; 13(4):327–333.10.1136/tc.2004.008169 [PubMed: 15564614]
- Maziak W, Ibrahim I, Rastam S, Ward K, Eissenberg TE. Waterpipe-associated particulate matter emissions. Nicotine and Tobacco Research. 2008; 10(3):519–523.10.1080/15622200801901989 [PubMed: 18324571]
- Meyers, LS.; Gamst, G.; Guarino, AJ. Applied Multivariate Research Design and Interpretation. Thousand Oaks, Calif: Sage; 2006.
- Noonan D. Exemptions for hookah bars in clean indoor air laws: a public health concern. Public Health Nursing. 2010a; 27(1):49–53.10.1111/j.1525-1446.2009.00826.x [PubMed: 20055968]
- Noonan, D. Dissertation Thesis. University of Virginia; Charlottesville, VA: 2010b. The new trend in tobacco: predicting waterpipe use among college students.

- Office of Institutional Assessment. Undergraduate Student Enrollment, 2006. 2006. Cited from University of Virginia Health Behavior Survey Retrieved December 1, 2008 from <http://www.virginia.edu/case/about/documents/HBS2007forWebsite.doc>
- Porter SR, Umbach PD. Student survey response rates across institutions: why do they vary? *Research in Higher Education*. 2006; 47(2):229–247.10.1007/s11162-005-8887-1
- Primack B, Sidani J, Agarwal A, Shandel WG, Donny E, Eissenberg TE. Prevalence of and associations with waterpipe smoking among US university students. *Annals of Behavioral Medicine*. 2008; 36(1):81–86.10.1007/s12160-008-9047-6
- Shihadeh A, Saleh R. Polycyclic aromatic hydrocarbons, carbon monoxide, “tar”, and nicotine in the mainstream smoke aerosol of narghile water pipe. *Food and Chemical Toxicology*. 2005; 43:655–66.10.1016/j.fct.2004.12.013 [PubMed: 15778004]
- Sepetdjian E, Shihadeh A, Saliba NA. Measurement of 16 polycyclic aromatic hydrocarbons in narghile smoke waterpipe tobacco smoke. *Food and Chemical Toxicology*. 2008; 46(5):1582–1590.10.1016/j.fct.2007/12/008 [PubMed: 18308445]
- Smith, S. Dissertation Thesis. John Hopkins University; Baltimore, MD: 2006. Nicotine product harm perception and psychosocial correlates of use in college freshman.
- Smith-Simone S, Maziak W, Ward KD, Eissenberg T. Waterpipe tobacco smoking: knowledge, attitudes, beliefs, and behavior in two U.S. samples. *Nicotine and Tobacco Research*. 2008; 10(2): 393–398.10.1080/14622200701825023 [PubMed: 18236304]
- Ward KD, Eissenberg T, Gray JN, Srinivas V, Wilson N, Maziak W. Characteristics of U.S. waterpipe users: a preliminary report. *Nicotine and Tobacco Research*. 2008; 9(12):1339–1346.10.1080/14622200701705019 [PubMed: 18058352]
- World Health Organization. Tobacco regulation advisory note: waterpipe tobacco smoking: health effects, research needs, and recommended action by regulators. 2005. Retrieved October 18, 2006 from http://www.who.int/tobacco/global_interaction/tobreg/Waterpipe%20recommendation_Final.pdf

Table 1

Demographics and Tobacco Use Behavior in the Study Sample (n=223)

Characteristics	Value
Age: <i>M (SD)</i>	19.85 (1.3)
Male (#, %)	102 (45.7)
Race: (#, %)	
Asian	29(13)
Black	7 (3.1)
Caucasian	160 (71.7)
Hispanic	8 (3.6)
Other	19 (8.5)
Year in School: (#, %)	
1 st year	67 (30)
2 nd year	58 (26)
3 rd year	40 (17.9)
4 th year	58 (26)
Ever Waterpipe Smoking: (#, %)	
Yes	136 (61)
Current Waterpipe Smoking: (#, %)	
Yes	30 (13.5)
Ever Cigarette Use: (#, %)	
Yes	111 (49.8)
Current Cigarette Use: (#, %)	
Yes	42 (18.8)

Table 2

Relationship of Intention to Smoke Waterpipe with Predictors (n=223)

Construct/Variable	Regression Coefficient	Standard Error	P Value	R ²
Attitude	.226	.017	<.001*	.334
Subjective Norm	.154	.076	.045*	.009
Behavioral Beliefs	.013	.004	<.001*	.001
Normative Beliefs	-.007	.004	.099	.002
Age	.026	.093	.780	.002
Gender -.084	.112	.493	.116	
School year: 1 st year	.441	.329	.182	.048
School year: 2 nd year	.538	.246	.030*	
School year: 3 rd year (4 th year reference group)	.488	.203	.017*	
Race: Asian	.152	.174	.386	.022
Race: Black and Hispanic	.005	.233	.982	
Race: Other (Caucasian reference group)	.236	.207	.205	
Current Cigarette Use (Yes vs. No)	.613	.162	<.001*	.295
Current Waterpipe Use (Yes vs. No)	1.194	.185	<.001*	

Note:

* Significant values

Cumulative R² of Theory of Reasoned Action constructs only = 34.6%Cumulative R² of entire model = 82.9%