Two RFLPs at the glutathione S-transferase 3 gene

E.L.Wong, G.Kandpal and A.E.Bale*

Department of Human Genetics, Yale University School of Medicine, New Haven, CT 06510, USA

Source/Description: GSTPi-1, a 0.73 kb EcoRI cDNA fragment containing the entire coding region of the glutathione S-transferase 3 gene (Moscow, J.A. *et al.*, 1988).

Polymorphism: PstI identifies constant bands at 2.8 kb and 2.05 kb and simple two allele polymorphism with a band at either 2.25 kb (A1) or 2.20 kb (A2). EcoRI identifies constant bands at 16 kb and 4.8 kb and a simple two allele polymorphism with a band at either 7.8 kb (B1) or 7.4 kb (B2).

Frequency:

PstI: 2.25 kb (A1) 0.03 (38 unrelated Caucasians)
2.20 kb (A2) 0.97
EcoRI: 7.8 kb (B1) 0.05 (29 unrelated Caucasians)
7.4 kb (B2) 0.95

Not Polymorphic For: BamHI, BglII, HindIII, MspI, RsaI, SstI, TaqI, or XbaI.

Chromosomal Location: Mapped to 11q13 by in situ hybridization (Moscow, J.A. *et al.*, 1988).

Mendelian Inheritance: Codominant inheritance demonstrated for the EcoRI polymorphism in 12 members of one family.

Probe Availability: ATCC.

Other Comments: The EcoRI and PstI RFLPs show no evidence for linkage disequilibrium.

Reference: Moscow, J.A. *et al.* (1988) *Proc. Natl. Acad. Sci. USA* **85**, 6518–6522.

A StuI RFLP in the human COL11A2 gene

K.S.E.Cheah, V.C.H.Lui, E.C.L.Yu¹ and L.C.S.Hsu²

Departments of Biochemistry, ¹Paediatrics and ²Orthopaedic Surgery, Hong Kong University, Sassoon Road, Hong Kong

Description: pVL39 is a 4.4 kb EcoRI subclone of the cosmid clone cosHcol.11 which contains the human $\alpha 2(XI)$ collagen (COL11A2) gene (1).

Polymorphism: Using pVL39 as a probe, a 2 allele StuI polymorphism is identified (A1: 9.1 kb; A2: 6.1 kb). A third fragment of 3.0 kb fragment derived from cleavage of allele A1 is also detected.

Frequency: In 41 unrelated Chinese, allele A1: 0.53; allele A2: 0.47.

Not Polymorphic For: TaqI, RsaI (100 chromosomes tested).

Chromosomal Localisation: COL11A2 has been mapped to chromosome 6p21.3 and is 45 kb centromeric to the HLA-DP β 2 locus (1).

Mendelian Inheritance: Autosomal codominant (15 meioses, 4 families).

Probe Availability: Contact Dr.Cheah.

Other Comments: Probe does not contain repetitive sequences. The COL11A2 gene also contains an EcoRI RFLP (2). The StuI polymorphic site is 11.2 kb 3' to the EcoRI RFLP.

Acknowledgements: This work was supported by a Hong Kong University Strategic Research Grant and the Arthritis & Rheumatism Council, UK. We thank M.S.Pui for technical assistance.

References: 1) Hanson, L.M., Gorman, P., Lui, V.C.H., Cheah, K.S.E., Solomon, E. and Trowsdale, J. (1989) *Genomics* **5**, 925. 2) Cheah, K.S.E., Pui, M.S., Lui, V.C.H., Wong, L., Hsu, L.C.S., Pun, K.K., Priestley, L.M. and Sykes, B.C. (1990) *Nucl. Acids Res.* **18**, 387.

* To whom correspondence should be addressed