Published in final edited form as:

Science. 2011 February 18; 331(6019): 912–916. doi:10.1126/science.1197454.

The antiproliferative action of progesterone in uterine epithelium is mediated by Hand2

Quanxi Li¹, Athilakshmi Kannan¹, Francesco J. DeMayo³, John P. Lydon³, Paul S. Cooke¹, Hiroyuki Yamagishi⁴, Deepak Srivastava⁵, Milan K. Bagchi^{2,*}, and Indrani C. Bagchi^{1,*}

¹Department of Comparative Biosciences, University of Illinois Urbana/Champaign, Urbana, IL

²Department of Molecular and Integrative Physiology, University of Illinois Urbana/Champaign, Urbana, IL

³Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX

⁴Keio University School of Medicine, Tokyo, Japan

⁵Gladstone Institute of Cardiovascular Disease, University of California, San Francisco, CA

Abstract

During pregnancy, progesterone inhibits the growth-promoting actions of estrogen in the uterus. However, the mechanism for this is not clear. The attenuation of estrogen-mediated proliferation of the uterine epithelium by progesterone is a prerequisite for successful implantation. Our study reveals that progesterone-induced expression of the basic helix-loop-helix transcription factor Hand2 in the uterine stroma suppresses the production of several fibroblast growth factors (FGFs), which act as paracrine mediators of mitogenic effects of estrogen on the epithelium. In mouse uteri lacking Hand2, continued induction of these FGFs in the stroma maintains epithelial proliferation and stimulates estrogen-induced pathways, resulting in impaired implantation. Thus, Hand2 is a critical regulator of the uterine stromal-epithelial communication that directs proper steroid regulation conducive for the establishment of pregnancy.

A sequential and timely interplay of the steroid hormones 17 β -estradiol (E) and progesterone (P) regulates critical uterine functions during the reproductive cycle and pregnancy (1–3). Whereas E drives uterine epithelial proliferation in cycling females, P counteracts E-induced endometrial hyperplasia. In mice, preovulatory ovarian E stimulates uterine epithelial growth and proliferation on days 1 and 2 of pregnancy (1). However, starting on day3, P produced by the corpora lutea terminates E-mediated epithelial proliferation. In response to P, epithelial cells exit from the cell cycle and enter a differentiation pathway to acquire the receptive state that supports embryo implantation on day4 of pregnancy (4–6). To identify the P-regulated pathways that underlie the implantation process, we had previously examined alterations in mouse uterine mRNA expression profiles in the peri-implantation period in response to RU486, a well-characterized progesterone receptor (PR) antagonist (7). Our results identified *Hand2*, a critical regulator of morphogenesis in a variety of tissues (8, 9), as a potential PR-regulated gene. Real-time PCR confirmed that the expression of Hand2 mRNA was greatly reduced in the uteri of RU486-treated mice (Fig. S1A). The expression of Hand2 protein, localized exclusively in the uterine stroma, was also abolished upon RU486 treatment (Fig. S1B), indicating that PR controls Hand2 expression in mouse uterus during early pregnancy.

*To whom correspondence should be addressed: Indrani C. Bagchi, ibagchi@illinois.edu, Milan K. Bagchi, mbagchi@life.illinois.edu.

To further confirm P-regulation of *Hand2*, ovaries were removed from non-pregnant mice and then these animals were injected with either vehicle or P. We observed intense nuclear expression of Hand2 protein in uterine stromal cells upon P treatment. Similar treatment of PR-null females showed no induction of Hand2 protein (Fig. 1A). These results established that P induces Hand2 expression in the uterine stroma. Consistent with its regulation by P, Hand2 expression was observed in the stromal cells underlying the luminal epithelium on days 3 and 4 of pregnancy (Fig. 1B).

To investigate the function of *Hand2* in the uterus, we created a conditional knockout of this gene in the adult uterine tissue. Crossing of mice harboring the “floxed” *Hand2* gene (*Hand2^{ff}*) with PR-Cre mice generated *Hand2^{d/d}* mice in which the *Hand2* gene is deleted selectively in cells expressing PR. As shown in Fig. S2, *Hand2* expression was successfully abrogated in uteri of *Hand2^{d/d}* mice. A breeding study demonstrated that *Hand2^{d/d}* females are infertile (Table S1). An analysis of the ovulation and fertilization in *Hand2^{ff}* and *Hand2^{d/d}* females revealed no significant difference in either the number or the morphology of the embryos recovered from their uteri (Figs. S3A and 3B). The serum levels of P and E were comparable in *Hand2^{ff}* and *Hand2^{d/d}* females on day4 of pregnancy, indicating normal ovarian function (Figs. S3C and S3D).

We next examined embryo attachment to the uterine epithelium by employing the blue dye assay, which assesses increased vascular permeability at implantation sites. *Hand2^{ff}* mice displayed distinct blue bands, indicative of implantation sites on day5 of pregnancy (Fig. S4). In contrast, none of the *Hand2^{d/d}* females showed any sign of implantation. Implanted embryos with decidual swellings were also absent in *Hand2^{d/d}* uteri on days 6 and 7 of pregnancy. Histological analysis of *Hand2^{ff}* females on day5 of pregnancy showed, as expected, a close contact of embryonic trophoblast with uterine luminal epithelium (Fig. 1C, a and b). In contrast, in *Hand2^{d/d}* uteri, blastocysts remained unattached in the lumen (c and d). These results suggested that in the absence of Hand2 expression in the stroma, the luminal epithelium fails to acquire competency for embryo implantation.

In mice, the window of uterine receptivity coincides with the P-mediated down-regulation of ER activity in uterine luminal epithelium (5, 6). As shown in Fig. S5, the levels of PR and ER α proteins in the luminal epithelium or stroma of *Hand2^{d/d}* uteri were comparable to those of *Hand2^{ff}* controls. An examination of the phosphorylation of ER α at serine 118, indicative of its transcriptionally active state (10), revealed a sharp reduction of this modification in the luminal epithelial cells of *Hand2^{ff}* uteri on days 3 and 4 of pregnancy (Fig. S6, a–d). In contrast, an increase in ER α phosphorylation was evident on these days in luminal epithelium of *Hand2^{d/d}* uteri (Fig. S6, e–h).

Consistent with this increase in ER’s transcriptional activity, expression of mRNAs corresponding to mucin 1 (*Muc-1*) and lactoferrin (*Lf*), well-characterized E-responsive genes in uterine epithelium (11), was significantly elevated in the *Hand2*-null uterus on day4 of pregnancy (Fig. 2A). In contrast, the expression of *Ihh* (12), *Alox15* (7), and *Irg1* (7), known P-responsive genes in uterine epithelium, remained unaltered in *Hand2^{d/d}* uteri (Fig. S7). Additionally, the mRNA levels of *Hoxa10*, a P-regulated stromal factor (3), and *Nr2f2*, a downstream target of *Ihh* in the uterine stroma (12), were unaffected in the uteri of *Hand2^{d/d}* mice. However the expression of leukemia inhibitory factor (*Lif*), a glandular factor that regulates uterine receptivity (2), was significantly reduced in *Hand2^{d/d}* uteri (Fig. S8).

Down-regulation of Muc-1 in the luminal epithelium is indicative of a receptive uterus (11). In contrast, persistent Muc-1 expression impairs acquisition of uterine receptivity and embryo implantation. On days 4 and 5 of gestation, a marked reduction in Muc-1 level was

seen in uterine epithelia of *Hand2^{ff}* mice, consistent with the attainment of receptive status (Fig. S9). However, Muc-1 expression persisted in uteri of *Hand2^{d/d}* mice at the time of implantation. Thus, elevated epithelial ER signaling led to increased expression of Muc-1 and corresponded to disrupted uterine receptivity and implantation failure in *Hand2^{d/d}* mice.

In normal pregnant uteri, the receptive state is also marked by a cessation in epithelial cell proliferation prior to implantation (1, 2). As expected, in *Hand2^{ff}* mice, Ki-67, a cell proliferation marker, was undetectable in the uterine epithelium as it attains receptive status on day4 of pregnancy (Fig. 2B, a). *Hand2^{d/d}* uteri, on the other hand, exhibited robust Ki67 expression in the luminal epithelium, which indicated sustained epithelial cell proliferation in the absence of *Hand2* (Fig. 2B, b).

The persistent proliferative state of uterine epithelium in the *Hand2^{d/d}* mice raised the possibility that stromal expression of *Hand2* mediates P action that opposes E-mediated epithelial proliferation. As shown in Fig. 2C, administration of E to ovariectomized *Hand2^{ff}* and *Hand2^{d/d}* mice led to robust uterine epithelial proliferation (a and b). Treatment with P alone induced proliferation exclusively in the uterine stromal cells of both genotypes (c and d). In *Hand2^{ff}* mice pretreated with P, administration of E showed no proliferative activity in the epithelium, suggesting a complete blockade of E-dependent proliferation by P. Under similar treatment conditions, *Hand2^{d/d}* uteri exhibited a marked E-induced epithelial proliferation, indicating an absence of antiproliferative effects of P on the uterine epithelium (e and f). These results established *Hand2* as a critical mediator of the actions of P in the stroma that inhibit E-dependent epithelial proliferation.

To identify downstream target(s) of *Hand2* in the uterus, we performed gene expression profiling of uterine stromal cells isolated from *Hand2^{ff}* and *Hand2^{d/d}* mice on day 4 of pregnancy. This study revealed elevated expression of mRNAs corresponding to several members of the fibroblast growth factor family (FGFs), namely *Fgf1*, *Fgf2*, *Fgf9*, and *Fgf18*, in uterine stroma of *Hand2^{d/d}* mice. Real-time PCR confirmed the induction of *Fgf1*, *Fgf9*, *Fgf2* and *Fgf18* mRNAs in uterine stromal cells of *Hand2^{d/d}* mice (Fig. 3A). The expression of *Hbegf* mRNA, encoding the heparin-binding epidermal growth factor (HB-EGF), was also increased in the mutant uteri (Fig. S10). In contrast, the expression of mRNAs of several other growth factor genes was either unaffected or slightly reduced in the *Hand2*-null uteri (Fig. S10). We also observed that the uterine expression of *Fgf2*, *Fgf9*, and *Fgf18* progressively declined with the rise of *Hand2* expression as pregnancy advanced from day 1 to day 4 (Fig. S11).

FGFs exert their paracrine responses through the cell surface FGF receptors (FGFRs) and a docking protein complex (13). FGF-stimulation of FGFRs leads to phosphorylation of specific tyrosine residues in a critical docking protein, FGFR substrate2 (FRS2), which guides the assembly of distinct multi-protein complexes, leading to activation of either ERK1/2 or PI3K/AKT signaling cascades (13). We determined the expression of FGFRs 1 and 2 in uteri on days 1 and 4 of pregnancy and found these receptor localized to the epithelium (Fig. S12A). The levels of mRNAs corresponding to *Fgfr* 1–3 were comparable in uteri of *Hand2^{ff}* and *Hand2^{d/d}* mice on day4 of pregnancy (Fig. S12B).

The activation of the FGFR signaling pathway in uteri of *Hand2^{ff}* and *Hand2^{d/d}* mice was monitored by examining the tyrosine phosphorylation status of FRS2. Whereas only low levels of phospho-FRS2 were seen in the uterine epithelium of *Hand2^{ff}* mice on day4 of pregnancy (Fig. 3B, a), a marked increase in its level was observed in the epithelium of *Hand2*-null uteri, indicating that FGF signaling is activated in the absence of *Hand2* (b). Phospho-FRS2 was undetectable in uterine epithelium of ovariectomized *Hand2^{ff}* mice in which P effectively blocks E-mediated proliferation (c). However, the epithelial cells of

Hand2-null uteri, which showed proliferative activity under similar hormone treatment conditions, exhibited a marked elevation in the level of phospho-FRS2, reflecting the activation of FGFR signaling in these cells (d). We also monitored the activated phosphorylated states of ErbB1 and ErbB4, the primary receptors mediating the actions of *Hbegf* (14). The activated forms of these receptors were undetectable in uterine epithelia of *Hand2^{fl/fl}* and *Hand2^{d/d}* mice on day4 of gestation (Fig. S13), indicating that the increased HB-EGF produced by the stroma of *Hand2*-null uteri did not act directly on the epithelial cells.

We next investigated whether the ERK1/2 and/or PI3K/AKT signaling pathways were activated downstream of FGFR in the epithelia of *Hand2*-ablated uteri. As shown in Fig. 3C, an increase in the level of phospho-ERK1/2 was seen in uterine epithelium of *Hand2*-null mice on day4 of pregnancy (b) and also in response to E treatment in the presence of P (d). In contrast, the phospho-AKT levels were undetectable or low and remained unaltered in the uterine epithelia of these mice (Fig. S14), suggesting that the ERK1/2 pathway, but not the PI3K/AKT pathway, is the key downstream mediator of enhanced FGF signaling in *Hand2*-null uteri.

To examine whether the elevated mitogenic activity in the luminal epithelium of *Hand2^{d/d}* uteri is induced by FGF and ERK1/2 signaling, we administered PD173074, a FGFR-specific inhibitor (15), or vehicle into uterine horns of *Hand2^{d/d}* mice at the time of implantation. The epithelia of vehicle-treated horn showed prominent expression of p-FRS2 and p-ERK1/2 (Fig. 4A, a and c). However, the levels of both p-FRS2 and p-ERK1/2 were reduced in the epithelia of PD173074-treated horn on day4 of pregnancy (Fig. 4A, b and d). We also observed a marked decline in the proliferative activity of *Hand2*-null uterine epithelia as indicated by decreased Ki67 staining (Fig. 4A, e and f). In parallel experiments, administration of PD184352, an ERK1/2-specific inhibitor (16), to uterine horns of *Hand2^{d/d}* mice suppressed the level of pERK1/2 (Fig. 4B, a and b) as well as luminal epithelial proliferation (Fig. 4B, c and d). Collectively these results are consistent with the hypothesis that increased FGF production by the *Hand2*-null uterine stroma stimulates epithelial proliferation by activating the FGFR-ERK1/2 pathway.

The ERK1/2-dependent phosphorylation of epithelial ER α at serine-118 is critical for the transcriptional activation of ER α (10). Interestingly, administration of either PD173074 (Fig. 4C, a–d) or PD184352 (Fig. 4C, e–h) to *Hand2*-null uterine horns blocked the phosphorylation of epithelial ER α at Ser 118 as well as Muc-1 expression. This result supported our view that elevated signaling by FGFR-ERK1/2 pathway in *Hand2^{d/d}* uteri is responsible for phosphorylation and activation of ER α in epithelial cells, thereby promoting persistent expression of Muc-1, which creates a barrier, preventing embryo attachment.

Earlier studies, employing tissue recombinants prepared with uterine epithelium and stroma isolated from neonatal wild-type and PR-null mice, indicated that the stromal PR plays an obligatory role in mediating the inhibitory actions of P on E-induced epithelial cell proliferation (17). However, the mechanism of this stromal-epithelial communication remained unknown. Our study has delineated a unique pathway in which *Hand2* operates downstream of P to regulate the production of FGFs, mitogenic paracrine signals that originate in the stroma and act on the FGFR(s) in epithelium to control its E-responsiveness (Fig. S15). The antiproliferative action of P in uterine epithelium is of clinical significance since breakdown of this action underpins E-dependent endometrial cancer (18). *Hand2*, therefore, is an important factor to be considered for the hormone therapy to block the proliferative actions of E in the endometrium.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank M. Laws for genotyping and Y. Li for IHC. This work was supported by the Eunice Kennedy Shriver NICHD/NIH through U54HD055787 as part of the Specialized Cooperative Centers Program in Reproduction and Infertility Research. The GEO accession number for microarray is GSE25881.

References and Notes

1. Finn CA, Martin L. J Reprod Fert. 1974; 39:195.
2. Carson DD, et al. Dev Biol. 2000; 223:217. [PubMed: 10882512]
3. Ramathal CY, et al. Semin Reprod Med. 2010; 28:17. [PubMed: 20104425]
4. Martin L, Das RM. J Endocrinol. 1973; 57:549. [PubMed: 4715727]
5. Bagchi IC, et al. Front Biosci. 2003; 8:s852. [PubMed: 12957875]
6. Pan H, Deng Y, Pollard JW. Proc Natl Acad Sci. 2006; 103:14021. [PubMed: 16966611]
7. Bagchi IC, et al. Semin Reprod Med. 2005; 23:38. [PubMed: 15714388]
8. Srivastava D, et al. Nat Genet. 1997; 16:154. [PubMed: 9171826]
9. Firulli AB. Gene. 2003; 312:27. [PubMed: 12909338]
10. Kato S, et al. Science. 1995; 270:1491. [PubMed: 7491495]
11. Surveyor GA, et al. Endocrinology. 1995; 136:3639. [PubMed: 7628404]
12. Lee KY, et al. Nat Genet. 2006; 38:1204. [PubMed: 16951680]
13. Eswarakumar VP, Lax I. Cytokine Growth Factor Rev. 2005; 16:139. [PubMed: 15863030]
14. Iwamoto R, Mekada E. Cytokine Growth Factor Rev. 2000; 11:335. [PubMed: 10959080]
15. Koziczak M, Holbro T. Oncogene. 2004; 23:3501. [PubMed: 15116089]
16. Solit DB, et al. Nature. 2006; 439:358. [PubMed: 16273091]
17. Kurita T, et al. Endocrinology. 1998; 139:4708. [PubMed: 9794483]
18. Kim JJ, Chapman-Davis E. Semin Reprod Med. 2010; 28:81. [PubMed: 20104432]

Fig. 1A

Fig. 1B

Fig. 1C

Figure 1.

P-regulated expression of Hand2 in the uterus is critical for implantation. **(A)** IHC of Hand2 protein in the uterine sections of ovariectomized WT mice treated with vehicle (a) or P (b) and PR-null mice treated with P (c). Panel d represents sections treated with non-immune IgG. **(B)** Uterine sections obtained from mice on days 2 to 4 of pregnancy were subjected to IHC using Hand2 antibody. Magnification 20X. **(C)** H and E staining of uterine sections obtained from *Hand2^{ff}* (a, b) and *Hand2^{dd}* (c, d) mice on day 5 (n=6) of pregnancy. b and d represent magnified images of a and c, respectively. Solid and dotted arrows point to embryo and luminal epithelium. L and S represent luminal epithelium and stroma, respectively.

Fig. 2A

Fig. 2B

Fig. 2C

Figure 2.

Enhanced ER α activity and proliferation in the luminal epithelium of *Hand2^{d/d}* uteri. **(A)** Real-time PCR was performed to monitor the expression of *Muc1* and *Ltf* in the uteri of day 4 pregnant *Hand2^{ff}* and *Hand2^{d/d}* mice, *P<0.001. **(B)** IHC of Ki67 in *Hand2^{ff}* (a) and *Hand2^{d/d}* (b) uteri on day 4 of pregnancy, 20X. Panel c shows uterine sections from *Hand2^{d/d}* mice treated with non-immune IgG, 40X. **(C)** IHC of Ki67 in the uterine sections of ovariectomized *Hand2^{ff}* and *Hand2^{d/d}* mice treated with E for one day (a and b), P for three days (c and d) or two days of P treatment followed by P and E (e and f).

Fig. 3A

Fig. 3B

Fig. 3C

Figure 3.

Enhanced FGFR signaling in the luminal epithelium of *Hand2*^{d/d} uteri. (A) Relative level of expression of *Fgf* family of growth factors in the uterine stroma of *Hand2*^{f/f} and *Hand2*^{d/d} mice on day 4 of pregnancy, *P<0.001. The levels of p-FRS2 (B) and p-ERK1/2 (C) were examined in the uterine sections (n=3) of both genotypes by IHC. a & b represent uterine sections obtained from day 4 pregnant mice. c & d indicate uterine sections from ovariectomized mice treated with two days of P followed by P and E.

Fig. 4A

Hand2^{d/d}

Fig. 4B

Hand2^{d/d}

Fig. 4C

Figure 4.

The inhibitor PD173074 (**A**) or PD184352 (**B**) was administered to one uterine horn of *Hand2^{d/d}* mice on day 3 of pregnancy (n=5). The other horn served as vehicle-treated control. Uterine horns were collected on day 4 morning and sections were subjected to IHC to detect p-FRS2, p-ERK1/2, and Ki67 (**C**) IHC of pERα and Muc-1 in uterine sections of *Hand2^{d/d}* mice treated with PD173074 or PD184352.