Published in final edited form as:

Org Lett. 2012 April 6; 14(7): 1880–1883. doi:10.1021/ol300476f.

Enantioselective Synthesis of (+)-Crocacin C. An Example of A Highly Challenging Mismatched Double Asymmetric δ -Stannylcrotylboration Reaction

Ming Chen and William R. Roush*

Department of Chemistry, Scripps Florida, Jupiter, Florida 33458

Abstract

A concise, enantioselective synthesis of (+)-crocacin C is described, featuring a highly diastereoselective mismatched double asymmetric δ -stannylcrotylboration of the stereochemically demanding chiral aldehyde **9** with the bifunctional crotylboration reagent (*S*)-**E-10**. The total synthesis of (+)-crocacin C was accomplished in seven steps (longest linear sequence) starting from commercially available precursors.

The crocacin A–D are a family of natural products isolated from *Chondromyces crocatus* and *Chondromyces pediulatus* (Figure 1).¹ Crocacin A, B, and D are dipeptides of glycine and a 6-aminohexenoic or a 6-aminohexadienoic acid with a polyketide-derived acyl residue connected to the nitrogen atom, while crocacin C is a primary amide of the acyl polyketide fragment. Initial biological studies revealed that the crocacin A–C display antifungal and cytotoxic activities. Compared to crocacin A–C, only crocacin D exhibited potent activity against *Saccharomyces cerevisiae* with a MIC of 1.4 ng/mL, which indicates that the dipeptide moiety of the crocacin A–C is crucial for their biological properties.¹ Recent crystallographic data suggest that the crocacin A–C are a new class of inhibitors of the cytochrome *bc*₁ complex.²

A characteristic structural feature of crocacin C is the *anti,anti*-dipropionate stereotriad (highlighted in yellow in Figure 2). It is evident that a mismatched double asymmetric crotylation reaction of an aldehyde substrate, such as **2**, with a chiral crotylmethyl reagent would be a direct, logical approach to this structural motif.^{3, 4}

However, an attempted^{5f} mismatched double asymmetric crotylboration of aldehyde **2** with crotylborationate **5** provided a 1:4 diastereomeric mixture in 60% yield, favoring the undesired 3,4-*anti*-4,5-*syn*-stereotriad **4** (Figure 2). Attempted crotylboration of aldehyde **2** using

roush@scripps.edu.

Supporting Information Available: Experimental procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Brown's crotylborane reagent **6** gave a 1:3 mixture of diastereomers, again favoring the undesired diastereomer **4**. Owing to the inability to directly access this requisite *anti,anti*-stereotriad (e.g., **3**), the central theme of multiple approaches developed for the synthesis of crocacin C^{5, 6} utilize indirect methods⁷ to prepare the *anti,anti*-stereotriad with high diastereoselectivity. Strategies involving aldol reactions,^{5a, 5e–g} epoxide ring-opening reactions,^{5b–d} or the desymmetrization of *meso* cyclic precursors^{5i, 6d} have been adopted to access the *anti,anti*-stereotriad units of crocacin C precursors.

We recently described⁸ highly diastereoselective syntheses of *anti,anti*-stereotriads using mismatched double asymmetric δ -stannylcrotylboration reactions of chiral aldehydes with crotylborane reagent (*S*)-**E-10**⁹ (Figure 3). Because it has been reported that reagents such as **5** and **6** are incapable of overriding the intrinsic diastereofacial preference of aldehyde **2** (Figure 2), we were intrigued whether our new reagent (*S*)-**E-10** could be adopted for synthesis of the *anti,anti*-stereotriad unit in **7**. Furthermore, the vinylstannane unit in **7** can be used in subsequent C–C bond forming reactions, for example, Stille¹⁰ coupling with vinyl iodide **8**.^{5a} We chose crocacin C as the target molecule for this study because it can be converted into other members of the crocacin family using a Cu-catalyzed coupling reaction as demonstrated by Dias and coworkers.¹¹

Starting from acyl oxazolidinone **11**, aldehyde **9** was obtained in four steps according to known procedures (Scheme 1).¹² Addition of aldehyde **9** to the crotylborane reagent (*S*)-**E-10**, generated from the enantioselective and enantioconvergent hydroboration of racemic allenylstannane ($\pm$)-**16**¹³ with (*d*Ipc)₂BH, at –78 °C followed by warming the reaction mixture to ambient temperature for a 24 h reaction period provided the targeted *anti,anti*-stereotriad **15** in 61% yield and with >15:1 diastereoselectivity.

Methylation of the secondary alcohol of **15** with Me₃O·BF₄ and Proton Sponge provided methyl ether **7**^{5a} in 88% yield. A Pd(0)-catalyzed Stille coupling^{5a, 10} of vinylstannane **7** with vinyl iodide **8**^{5a} gave (+)-crocacin C (**1**) in seven steps (longest linear sequence) and in 21% overall yield from **11** without any protecting group manipulations. The spectroscopic data (¹H NMR, ¹³C NMR, [α]_D) of synthetic (+)-crocacin C were in excellent agreement with the data previously reported for the natural product.^{1, 5}

The intrinsic diastereofacial preference of aldehyde **9** was assessed by using an anti-crotylboration reaction with the achiral pinacol (*E*)-crotylboronate **17** (Scheme 2). This reaction provided an 18:1 mixture of 3,4-*anti*-4,5-*syn*-stereotriad **18** and *anti, anti*-stereotriad **19** in 77% yield, with **18** as the major product (as expected^{3, 14}). In contrast, the mismatched double asymmetric δ -stannylcrotylboration of aldehyde **9** with (*S*)-**E-10** provided the *anti,anti*-stereotriad **15** with >15:1 diastereoselectivity. No other crotylation diastereomers were observed in the reaction mixture. Protodestannylation of **15** under acidic conditions (TsOH·H₂O) provided alcohol **19** in 87% yield, which matched the minor isomer obtained from crotylboration of **9** with achiral crotylboronate **17**.

The mismatched double asymmetric δ -stannylcrotylboration of **9** with (*S*)-**E-10** thus represents yet another case⁸ where a significant intrinsic diastereofacial barrier, as presented by chiral aldehyde **9**, is overridden by the chiral reagent (*S*)-**E-10**. The free energy contribution of reagent (*S*)-**E-10** (i.e., the enantioselectivity of the reagent expressed in energetic terms) necessary to override the 18:1 intrinsic diastereofacial preference of **9** and to generate homoallylic alcohol **15** with >15:1 mismatched diastereoselectivity is ≥ 3.3 kcal/mol (reaction at 23 °C). The exceptional enantioselectivity of (*S*)-**E-10** defines a new standard of excellence that all future methodological studies on enantioselective crotylboration or crotylmatal-carbonyl addition reactions should be judged against.

In conclusion, the total synthesis of (+)-crocacin C (**1**) was completed in seven steps (longest linear sequence), which represents the shortest synthesis of **1** reported to date. Most importantly, the mismatched double asymmetric δ -stannylcrotylboration of aldehyde **9**, with a significant 18:1 intrinsic diastereofacial preference, was achieved with exceptional selectivity (>15:1) by using the crotylborane reagent (*S*)-**E-10**. The vinylstannane unit in the derived *anti*, *anti*-stereotriad **15** facilitates the subsequent Stille reaction that was used to complete this short synthesis of crocacin C. Other applications of reagent (*S*)-**E-10** in the synthesis of biologically active natural products will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support provided by the National Institutes of Health (GM038436) and Eli Lilly (for a predoctoral fellowship to M.C.) is gratefully acknowledged.

References

1. (a) Kunze B, Jansen R, Hofle G, Reichenbach H. J Antibiot. 1994; 47:881. [PubMed: 7928674] (b) Jansen R, Washausen P, Kunze B, Reichenbach H, Hofle G. Eur J Org Chem. 1999:1085.
2. Crowley PJ, Berry EA, Cromarti T, Daldal F, Godfrey CRA, Lee DW, Phillips JE, Taylor A, Viner R. Bioorg Med Chem. 2008; 16:10345. [PubMed: 18996700]
3. Reviews of reactions of carbonyl compounds with crotylmetal reagents: (a)Roush WR. Trost BM. Comprehensive Organic Synthesis. Pergamon PressOxford1991; 2:1.(b) Yamamoto Y, Asao N. Chem Rev. 1993; 93:2207.(c)Denmark SE, Almstead NG. Otera J. Modern Carbonyl Chemistry. Wiley-VCHWeinheim2000:299.(d) Denmark SE, Fu J. Chem Rev. 2003; 103:2763. [PubMed: 12914480] (e) Lachance H, Hall DG. Org React. 2008; 73:1.
4. Masamune S, Choy W, Petersen JS, Sita LR. Angew Chem, Int Ed Engl. 1985; 24:1.
5. For total syntheses of crocacin C, see: (a) Feutrill JT, Lilly MJ, Rizzacasa MA. Org Lett. 2000; 2:3365. [PubMed: 11029212] (b) Chakraborty TK, Jayaprakash S. Tetrahedron Lett. 2001; 42:497. (c) Chakraborty TK, Jayaprakash S, Laxman P. Tetrahedron. 2001; 57:9461.(d) Dias LC, de Oliveira LG. Org Lett. 2001; 3:3951. [PubMed: 11720577] (e) Sirasani G, Paul T, Andrade RB. J Org Chem. 2008; 73:6386. [PubMed: 18630880] (f) Sirasani G, Paul T, Andrade RB. Bioorg Med Chem. 2008; 18:3648. [PubMed: 20363632] (g) Gillis EP, Burke MD. J Am Chem Soc. 2008; 130:14084. [PubMed: 18837550] (h) Feutrill JT, Lilly MJ, White JM, Rizzacasa MA. Tetrahedron. 2008; 64:4880.(i) Candy M, Audran G, Bienayme H, Bressy C, Pons JM. J Org Chem. 2010; 75:1354. [PubMed: 20128625]
6. For formal syntheses of crocacin C, see: (a) Gurjar MK, Khaladkar TP, Borhade RG, Murugan A. Tetrahedron Lett. 2003; 44:5183.(b) Raghavan S, Reddy SR. Tetrahedron Lett. 2004; 45:5593.(c) Besev M, Brehm C, Fürstner A. Collect Czech Chem Commun. 2005; 70:1696.(d) Yadav JS, Reddy PV, Chandraiah L. Tetrahedron Lett. 2007; 48:145.(e) Yadav JS, Reddy MS, Rao PP, Prasad AR. Synlett. 2007:2049.
7. For reviews of methods commonly used to synthesize the anti, anti dipropionate stereotriad: (a) Hoffmann RW. Angew Chem, Int Ed Engl. 1987; 26:489.(b) Hoffmann RW, Dahmann G, Andersen MW. Synthesis. 1994:629.
8. Chen M, Roush WR. J Am Chem Soc. 2012; 134:3925. [PubMed: 22332989]
9. (a) Chen M, Roush WR. J Am Chem Soc. 2011; 133:5744. [PubMed: 21449594] For synthetic applications of reagent (*S*)-**E-10**, see: (b) Sun H, Abbott JR, Roush WR. Org Lett. 2011; 13:2734. [PubMed: 21510632] (c) Yin M, Roush WR. Tetrahedron. 2011; 67:10274.(d) Chen M, Roush WR. Org Lett. 2012; 14:426. [PubMed: 22148850]
10. (a) Stille JK. Angew Chem, Int Ed Engl. 1986; 25:508.(b) Farina V, Krishnamurthy V, Scott WJ. Org React. 1997; 50:1.

11. Dias LC, de Oliveira LG, Vilcachagua JD, Nigsch F. J Org Chem. 2005; 70:2225. [PubMed: 15760209]
12. Evans DA, Miller SJ, Ennis MD. J Org Chem. 1993; 58:471.
13. Racemic allenylstannane ($\pm$)-16 is prepared in two steps from commercially available ($\pm$)-3-butyn-2-ol using the route described for (M)-16 and (P)-16. See: (a) Marshall JA, Lu ZH, Johns BA. J Org Chem. 1998; 63:817. [PubMed: 11672079] (b) Marshall JA, Chobanian H. Org Syn. 2005; 82:43.
14. Hoffmann RW, Weidmann U. Chem Ber. 1985; 118:3966.

Figure 1.
Structures of crocacin A–D.

Figure 2. Attempted mismatched double asymmetric crotylboration reactions of aldehyde **2** with reagents **5** and **6** for synthesis of the *anti,anti*-stereotriad of crocacin C.^{4f}

Figure 3.
Crocacin C, retrosynthetic analysis.

Scheme 1.
Total Synthesis of (+)-Crocacin C (1)

Scheme 2.
Crotylboration Studies of Aldehyde **9**