Published in final edited form as:

Bioorg Med Chem Lett. 2012 April 15; 22(8): 2833–2836. doi:10.1016/j.bmcl.2012.02.078.

Deep-red fluorescent imaging probe for bacteria

Alexander G. White^a, Brian D. Gray^b, Koon Yan Pak^b, and Bradley D. Smith^{a,*}

^aDepartment of Chemistry and Biochemistry, 236 Nieuwland Science Hall, University of Notre Dame, Notre Dame, IN 46556, USA

^bMolecular Targeting Technologies Incorporated, 833 Lincoln Ave., Unit 9, West Chester, PA 19380, USA

Abstract

A versatile deep-red fluorescent imaging probe is described that is comprised of a bis(zinc(II)-dipicolylamine) targeting unit covalently attached to a pentamethine carbocyanine fluorophore with Cy5-like spectroscopic properties. A titration assay based on fluorescence resonance energy transfer is used to prove that the probe selectively associates with anionic vesicle membranes whose composition mimics bacterial cell membranes. Whole-body optical imaging experiments show that the probe associates with the surfaces of both Gram-positive and Gram-negative bacteria cells, and it can target the site of bacterial infection in a living mouse. In vivo accumulation at the infection site and subsequent clearance occurs more quickly than a structurally related near-infrared bis(zinc(II)-dipicolylamine) probe. The fact that the same deep-red probe molecule can be used for spectroscopic assays, cell microscopy, and in vivo imaging studies, is an important and attractive technical feature.

Keywords

fluorescent molecular probe; imaging bacterial infection; zinc dipicolylamine; in vivo imaging; optical imaging; FRET

Over the last five years we have developed a series of fluorescent bis(zinc(II)-dipicolylamine) (bis(Zn-DPA)) imaging probes that target anionic membranes and these molecular probes have proven useful for optical imaging of two major classes of cells: dead and dying mammalian cells and bacterial cells.¹ Initially, we synthesized blue and green emitting bis(Zn-DPA) probes that were suitable for bacterial cell studies using flow cytometry and fluorescence microscopy.² We have also developed a near-infrared version that is very effective for in vivo imaging studies of infection models in rodents.³ Imaging probes that incorporate a deep-red fluorophore with excitation/emission wavelengths that match the common Cy5 filter set (approximately 645/665 nm) in microscopes, microarrays, and flow cytometers are particularly useful.⁴ This wavelength region is still visible to the eye of a microscope operator which facilitates microscopic studies, yet it is sufficiently red to allow effective in vivo imaging in living animals.^{4c,d} Performance limitations, due to

© 2012 Elsevier Ltd. All rights reserved.

*Telephone: 1-574-631-8632, Fax: 1-574-631-6652, smith.115@nd.edu.

Supporting Information

Probe synthesis, experimental methods, imaging analyses, and Figs. S1–S4 can be found in the online version at doi:10.1016/j.bmcl.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

undesired absorption and autofluorescence by endogenous biomolecules and scattering of the light by animal skin and tissue, are attenuated at longer wavelengths.⁵ Previously, we have reported deep-red bis(Zn-DPA) probes with structurally unusual squaraine rotaxane fluorophores,⁶ but the need remains for additional deep-red bis(Zn-DPA) probes that are relatively cheap to produce. A low production cost is extremely important for many imaging applications that require large amounts of probe material such as high throughput cell screening assays or temporal studies of large cohorts of animal models.⁷ Here, we describe the new and versatile deep-red probe, **1**, that is comprised of a bis(Zn-DPA) targeting unit covalently attached to a pentamethine carbocyanine fluorophore with Cy5-like spectroscopic properties (Scheme 1). We demonstrate how this probe can be used in a spectrometric Fluorescence Resonance Energy Transfer (FRET) assay that measures probe association with model vesicle membranes of any phospholipid composition. We also show how probe **1** can be used to stain bacteria cells and optically image sites of bacterial infection in a living mouse.

The straightforward synthesis of probe **1** using readily available building blocks is described in the Supporting Information. In short, the bis(Zn-DPA) unit is coupled to the carbocyanine fluorophore by forming an amide-bond.⁸ Probe **1** is highly water soluble and exhibits attractive photophysical properties in aqueous solution ($\lambda_{\text{abs}} = 643 \text{ nm}$, $\epsilon = 42,800 \text{ M}^{-1}\text{cm}^{-1}$, $\lambda_{\text{em}} = 658 \text{ nm}$, $\Phi_{\text{f}} = 0.13$). Based on previous experience, we expected that the bis(Zn-DPA) unit would promote association with membrane surfaces that are rich in anionic phospholipids. One example is the bacterial membrane. The cellular envelope of Gram-positive bacteria is rich in phosphatidylglycerol and other phosphorylated amphiphiles, and the outer membrane of Gram-negative bacteria is known to contain a large fraction of lipopolysaccharides that have anionic Lipid A as the core structure (Scheme 1).² Before starting the bacterial studies we wanted to measure the association of probe **1** with model vesicles that mimic bacterial cell membranes. It is a technical challenge to quantitatively measure the binding of small molecules to bilayer membranes.⁹ We were attracted to the Cy5-like, carbocyanine fluorophore in probe **1** because it can accept energy from a donor Cy3-like chromophore via a FRET pathway.¹⁰ Thus, we constructed a FRET based titration assay for membrane binding (Figure 1A). The lipophilic dye **3** (also known as DiIC₁₈) was incorporated into the membrane of vesicles and then probe **1** was added in incremental amounts. Association of **1** to membranes leads to strong energy transfer, which is observed as a loss in the donor emission band at 570 nm and an increase in the acceptor emission band at 665 nm. The spectral data in Figure 1B–D is plotted in Figure 1E as a set of titration isotherms. The plots show that probe **1** has very weak affinity for zwitterionic membranes composed of 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC), and stronger affinity for membranes that are enriched in Lipid A (mimic of Gram-negative membrane) or 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphoglycerol (POPG, mimic of Gram-positive bacteria). These titration curves fitted well to a binding model of the probe forming a 1:1 complex with the anionic phospholipid that is exposed on the outer surface of the vesicle. The association constant for membranes that are rich in Lipid A ($1.2 \times 10^5 \text{ M}^{-1}$) is higher than for membranes that are rich in POPG ($1.3 \times 10^4 \text{ M}^{-1}$). This is structurally reasonable since the polar head group of the Lipid A structure contains two dianionic phosphate monoesters groups (Scheme 1) that have an inherently higher affinity for the cationic bis(Zn-DPA) unit in probe **1**. The titration curves also show a lower extent of FRET for **1** binding to Lipid A-rich membranes, reflecting the five times lower number of membrane binding sites for the same amount of FRET donor dye **3**. From an operational perspective we find that this FRET-based vesicle binding assay is technically straightforward to conduct and highly reproducible. The fact that it can be used to measure association of **1** with vesicles of any composition makes it highly versatile.

The outer membrane surface of healthy animal cells is composed primarily of zwitterionic phospholipids, and we have previously shown that bis(Zn-DPA) probes can selectively target sites of bacterial infection in living animals.^{2,3} The vesicle titration data in Figure 1 suggested that probe **1** should also exhibit selective and universal affinity for the cell envelope of virtually all genera of bacteria.^{2,3,6} This hypothesis was confirmed first by conducting pelleting experiments that treated representative strains of Gram-positive and Gram-negative bacteria (*Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhimurium*) with probe **1** and showing in each case that the bacterial pellet, produced after centrifugation and washing, exhibited intense deep-red fluorescence. A subsequent animal imaging experiment demonstrated that the deep-red fluorescence could be observed through skin and tissue. Specifically, two separate populations of Gram-positive *Staphylococcus aureus* Xen31 were mixed with either deep-red probe **1** or the known near-infrared probe **2** (excitation/emission wavelengths of 794/810 nm)³ and each cell/probe mixture was converted into a pellet by centrifugation. The two pellets were washed and then injected into the opposite rear legs of an anesthetized mouse, and a whole-body optical image of the mouse was recorded using an in vivo animal imaging station. The images in Figure 2 show that both probes have good affinity for the *S. aureus* and that the deep-red fluorescence signal for **1** could be observed separately from the near-infrared signal of **2**.

Next, two sets of animal imaging experiments demonstrated the effectiveness of probe **1** to selectively target a bacterial infection within a living mouse. In each case, a bolus of $\sim 10^8$ bacterial cells was injected into the rear leg of an anesthetized mouse and then, 3 h later, a dose of probe **1** (10 nmol) was injected intravenously via the tail vein. The whole-body images in Figure 3 were acquired at 3 h after the probe injection. In this case, the infection was a bioluminescent strain of *Salmonella typhimurium* AM3, a Gram-negative bacteria. Shown in Figure 3A is the bioluminescent emission from the *S. typhimurium*, and in Figure 3B is a deep-red fluorescence image. The overlaid image in Figure 3C confirms that probe **1** is highly colocalized at the same anatomical site as the bioluminescent bacteria.

The images in Figure 4 summarize an analogous experiment that used probe **1** to target an infection of Gram-positive *S. aureus* Xen31. The montage shows the accumulation of probe **1** and its subsequent clearance from the animal. A region-of-interest analysis of these in vivo images indicated that the maximum target to background ratio of 4.2 was achieved after 4 h, and that there was very little fluorescent signal remaining at the site of bacterial infection after 24 h. This latter fact was confirmed with an ex vivo biodistribution analysis of the excised tissues after animal sacrifice at 24 h (see Figure S4). A comparison of the in vivo imaging data in Figure 4 with literature shows that probe **1** (10 nmol) can be employed at about a four-fold lower dosage than probe **2** for the same T/NT ratio.³ Furthermore, probe **1** washes out much faster from the infection site than probe **2**. This suggests that probe **1** may be better suited for longitudinal imaging studies that must repeatedly image the same infected animal over time.¹¹ Fluorescent probes like **1** and **2** with a single bis(Zn-DPA) targeting unit can detect a mouse leg infection containing $\sim 10^7$ or more bacterial cells, but this visualization threshold can be lowered by using higher affinity bacterial probes whose chemical structures containing multiple bis(Zn-DPA) units.⁶ While the tissue penetration of near-infrared light associated with **2** is slightly greater than the red light of **1**, neither probe is expected to be very useful for fluorescent planar imaging of deep tissue infection sites in larger animals. However, continued advances in fluorescence tomography may ameliorate this technical limitation.^{7g}

In summary, we find that deep-red bis(Zn-DPA) probe **1** exhibits several useful photophysical and supramolecular properties that make it a very attractive and versatile targeted probe for fluorescence imaging studies of membranes that are rich in anionic phospholipids. The probe can be employed in a technically straightforward FRET-based

assay that measures affinity for model vesicle membranes of any composition. The bis(Zn-DPA) unit in probe **1** selectively targets the abundant phosphorylated amphiphiles within the cell envelope of Gram-positive and Gram-negative bacteria.^{2,3,6} Thus, probe **1** is expected to be a universal fluorescent imaging probe for any bacterial infection in a living mouse. The fact that the same probe molecule can be used for spectroscopic assays, cell microscopy, and in vivo imaging studies, is an important technical feature that simplifies the experimental design and reduces experimental uncertainty since there is no change in probe molecular structure.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH (GM059078 to B.D.S.) and the Notre Dame Integrated Imaging Facility. All animal care and imaging protocols were approved by the Notre Dame Institutional Animal Care and Use Committee. We are grateful to members of the Freimann Life Sciences Center for professional animal care.

References

- (a) O'Neil EJ, Smith BD. *Coord Chem Rev.* 2006; 250:3068–3080.(b) Drewry JA, Gunning PT. *Coord Chem Rev.* 2011; 255:459.
- Leevy WM, Johnson JR, Lakshmi C, Morris J, Marquez M, Smith BD. *Chem Commun.* 2006; 15:1595.
- (a) Leevy WM, Gammon ST, Jiang H, Johnson JR, Maxwell DJ, Marquez M, Piwnica-Worms D, Smith BD. *J Am Chem Soc.* 2006; 128:16476. [PubMed: 17177377] (b) Leevy WM, Gammon ST, Johnson JR, Lampkins AJ, Jiang H, Marquez M, Piwnica-Worms D, Suckow MA, Smith BD. *Bioconjugate Chem.* 2008; 19:686.
- (a) Terai T, Nagano T. *Curr Opin Chem Biol.* 2008; 12:515. [PubMed: 18771748] (b) Trinquet E, Mathis G. *Mol Biosyst.* 2006; 2:380. [PubMed: 16880958] (c) Enniga J, Sansonetti P, Tournibize R. *Trends Microbiol.* 2007; 15:483. [PubMed: 17983749] (d) Hoppe AD, Seveau S, Swanson JA. *Cellular Microbiol.* 2009; 11:540.
- Kobayashi H, Ogawa M, Alford R, Choyke PL, Urano Y. *Chem Rev.* 2010; 110:2620. [PubMed: 20000749]
- (a) Johnson JR, Fu N, Arunkumar E, Leevy WM, Gammon ST, Piwnica-Worms D, Smith BD. *Angew Chem Int Ed.* 2007; 46:5528.(b) White AG, Fu N, Leevy WM, Lee JJ, Blasco MA, Smith BD. *Bioconjugate Chem.* 2010; 21:1297.
- (a) Leevy WM, Serazin N, Smith BD. *Drug Discovery Today: Disease Models.* 2007; 4:91. [PubMed: 20376332] (b) Zhao XJ, Hilliard LR, Mechery SJ, Wang YP, Bagwe RP, Jin SG, Tan WHA. *Proc Nat Acad Sci USA.* 2004; 101:15027. [PubMed: 15477593] (c) Qu LW, Luo PG, Taylor S, Lin Y, Huang WJ, Anyadike N, Tzeng TRJ, Stutzenberger F, Latour RA, Sun YP. *J Nanosci Nanotechnol.* 2005; 5:319. [PubMed: 15853155] (d) Zhu C, Yang Q, Liu L, Wang S. *Angew Chem Int Ed.* 2011; 50:9607.(e) Inglese J, Johnson RL, Simeonov A, Xia M, Zhang W, Austin CP, Auld DS. *Nature Chem Biol.* 2007; 3:466. [PubMed: 17637779] (f) Ning X, Lee S, Wang Z, Kim D, Stubblefield B, Gilbert E, Murthy N. *Nature Materials.* 2011; 10:602.(g) Signore A, Mather SJ, Piaggio G, Malviya G, Dierckx RA. *Chem Rev.* 2010; 110:3112. [PubMed: 20415479] (h) Kong Y, Yao H, Ren H, Subbian S, Cirillo SLG, Saccettini JC, Rao J, Cirillo JD. *Proc Natl Acad Sci USA.* 2010; 107:12239. [PubMed: 20566877] (i) Melican K, Richter-Dahlfors A. *Curr Opin Microbiol.* 2009; 12:31. [PubMed: 19135408]
- (a) Shao F, Yuan H, Josephson L, Weissleder R, Hilderbrand SA. *Dyes Pigments.* 2011; 90:119.(b) Joeri KJ, Buckle T, Yuan H, van den Berg NS, Oishi S, Fujii N, Josephson L, van Leeuwen FWB. *Bioconjugate Chem.* 2011; 22:859.
- (a) Hofmann A, Huber R. *Methods Enzymol.* 2003; 372:186–216. [PubMed: 14610814] (b) Surman AJ, Kenny GD, Kumar DK, Bell JD, Casey DR, Vilar R. *Chem Commun.* 2011; 47:10245.

10. (a) Massey M, Algar WR, Krull UJ. *Analyt Chim.* 2006; 568:181–189.(b) You M, Li E, Wimley WC, Hristova K. *Anal Biochem.* 2005; 340:154. [PubMed: 15802141]
11. While a complete toxicity study has not yet been conducted, it is worth noting that the mice easily tolerate the presence of probe **1** (10 nmol) for at least 24 h, as judged by their continued grooming and nesting activities.

Figure 1.

Association of fluorescent energy acceptor **1** to the surface of vesicles containing 1 mol % of the energy donor **3** induces FRET. [A] Schematic of the FRET process. [B, C, D] Normalized spectra of FRET induced by incremental addition of **1** to vesicles containing **3** and composed of only POPC, POPC:POPG (50:50), or POPC:lipid A (90:10), respectively. [E] Plots of the spectral data showing changes in the ratio of acceptor/donor fluorescence intensities due to association-induced FRET.

Figure 2.

Optical imaging of anesthetized hairless mouse with separate rear leg injections of *S. aureus* Xen31 ($\sim 10^8$ cells) prelabeled with 10 nmol of either **1** (A, cyan color showing deep-red signal via the Cy5.5 filter set) or **2** (B, red color showing infrared signal via the ICG filter set). Panel C shows an overlaid image of the spectrally unmixed emission signals from the two populations of *S. aureus* Xen31 labeled with **1** or **2**. Intensity scale for each image is in arbitrary units. N = 3.

Figure 3. Optical imaging of anesthetized hairless mouse with a rear leg injection of bioluminescent *S. typhimurium* AM3 ($\sim 10^8$ cells) and treated 3 h later with 10 nmol of **1** via the tail vein. [A] Bioluminescent signal from bacteria, [B] fluorescent signal from **1**, [C] overlay of panels A and B. Images were acquired 3 h after probe dosage. Intensity scale for each image is in arbitrary units. N =3.

Figure 4.

Fluorescence imaging montage of anesthetized hairless mouse containing a rear leg injection of *S. aureus* Xen31 ($\sim 10^8$ cells) and treated 3 h later with 10 nmol of **1** via the tail vein. Mouse was imaged at [A] 0 h, [B] 3 h, [C] 6 h, and [D] 24 h after probe dosage. Intensity scale (arbitrary units) applies to all images. N = 4.

Scheme 1.

Chemical structures of the fluorescent probes and phospholipids used in this study.