Polymerase chain reaction (PCR) for detection of PstI polymorphism in the human PF4 gene

Banumathi Ramachandran, Eric F. Rappaport, Saul Surrey, Mortimer Poncz and Elias Schwartz

Division of Hematology, The Children's Hospital of Philadelphia and Departments of Pediatrics and Human Genetics, University of Pennsylvania School of Medicine, Philadelphia, PA 19104, USA

Source and Description: A cDNA clone isolated from a human erythroleukemia (HEL) cell cDNA library was used initially for detection of blotted genomic DNA fragments. Once the polymorphic site was identified, PCR and PstI digestion were employed for detection of the site. A 1.6 kb fragment encompassing the polymorphic site was amplified by PCR using the following two synthetic oligonucleotide primers: 5' GGA-TCATGATCGCAGCTGCTC 3' and 5' TCTGGTCCATGGT-GGTAGGCT 3'. This fragment is located in the PF4var1 gene (1).

Polymorphism: On a genomic Southern blot, PstI generates 2 constant bands (1.4 kb and 0.6 kb) and either a 2.0 or 1.0 kb band. PstI digestion of the PCR amplified fragment yields a 1.3 kb fragment if the site is absent and a 1.0 kb fragment if present (Figure).

Frequency: Studied in 17 unrelated individuals.

1.3 kb (A1): 0.68

1.0 kb (A2): 0.32

Not Polymorphic For: 10 restriction enzymes (see ref. 2).

Chromosomal Location: The human PF4 gene has been localized to the region 4q12-q21 by hybridization in situ (3).

Mendelian Inheritance: Co-dominant inheritance was observed in 3 families (13 members).

Comments: Genomic DNA was first amplified using 2 primers to give a 2.0 kb fragment (5' CAGGTCCCGCCTCACCCGCGC 3' and 5' TAGGTACCTCCCTCCCTTCCA 3'). 30 cycles of PCR were carried out by denaturing at 95°C for 2 minutes, annealing at 60°C for 2 minutes and extending at 72°C for 3 minutes. The final extension was at 72°C for 7 minutes. The DNA was then subjected to a second round of PCR using the nested primers listed above to generate the 1.6 kb fragment used for the PstI digestion. The reaction conditions used were identical to the previous PCR except that annealing was done at 55°C for 2 minutes.

Acknowledgements: This work was supported by grants from the National Institutes of Health (DK16691, HL28157, P01 HL40387 and HL37419).

References: 1) Green *et al.* (1989) *Mol. Cell. Biol.* **9**, 1445–1451. 2) Guzzo *et al.* (1987) *Nucl. Acids Res.* **15**, 380. 3) Griffin *et al.* (1987) *Cytogenet. Cell Genet.* **45**, 67–69.

A three allele TaqI polymorphism at TOP1 gene

Lone Sunde*, Eigil Kjeldsen, Toshiwo Andoh¹, Jack D. Keene² and Lars Bolund

Institute of Human Genetics, University of Aarhus, DK-8000 Aarhus C, Denmark, ¹Department of Hygienic Chemistry, Meiji College of Pharmacy, Tanashi, Tokyo 188, Japan and ²Department of Microbiology and Immunology, Duke University, NC 27710, USA

Source and Description of Clone: cDNA clones encoding human topoisomerase I were isolated from an expression vector library (λ gt11) by screening with autoimmune anti-topoisomerase I serum. A 3.6 kb EcoRI fragment, comprising the carboxyl-terminal fragment of the human topoisomerase I gene, was sub-cloned into pUC19 and used for hybridization. The clone (pSc170) recognizes a 4.1 kb mRNA in Northern blots (1).

Polymorphism: Hybridization to TaqI digested human genomic DNA identifies a three allele polymorphism with bands of 8.0 kb (A1), 7.0 kb (A2), or 5.4 kb (A3); and invariant bands of 6.0, 5.0, 4.0, 3.5 and 2.1 kb.

Frequency: Studied in 68 unrelated European Caucasians:

8.0 kb (A1) 0.32

7.0 kb (A2) 0.60

5.4 kb (A3) 0.07

Not Polymorphic for: EcoRI, HindIII, BglII, PstI, PvuII. Tested by comparing the hybridization pattern of retracted DNA from one universally homozygous hydatidiform mole with that of DNA pooled from 64 unrelated European Caucasians.

Chromosomal Localization: 20q11.2–13.1 (2). (Pseudogenes have been demonstrated on chromosomes 1 and 22 (2). The described polymorphism is unique for the active gene, as it was also detected by a 5' fragment of the cloned sequence, not present in the pseudogenes).

Mendelian Inheritance: Co-dominant segregation observed in 17 unrelated families (53 individuals).

Probe Availability: Contact E. Kjeldsen, T. Andoh or J. D. Keene.

Acknowledgements: Supported by grants from the Danish Biotechnology Programme and the Danish Cancer Society.

References: 1) D'Arpa, P. *et al.* (1988) *Proc. Natl. Acad. Sci. USA* **85**, 2543–2547. 2) Kunze, N. *et al.* (1989) *Hum. Genet.* **84**, 6–10.

* To whom correspondence should be addressed