Published in final edited form as:

J Investig Med. 2010 December ; 58(8): 987–990. doi:10.231/JIM.0b013e3181fa02f7.

Airway Platelet Activation is Associated with Airway Eosinophilic Inflammation in Asthma

Angela S Benton, BAS^{1,2,*}, Nikila Kumar, BS^{2,*}, Jennifer Lerner, BS¹, Andrew Wiles, BS¹, Matthew Foerster, BA¹, Stephen J Teach, MD, MPH^{3,4,5}, and Robert J Freishtat, MD, MPH^{1,4,5,6}

¹Center for Genetic Medicine Research, Children's National Medical Center, Washington, DC, United States

²The George Washington University School of Medicine and Health Sciences, Washington, DC, United States

³Center for Clinical and Community Research, Children's National Medical Center, Washington, DC, United States

⁴Division of Emergency Medicine, Children's National Medical Center, Washington, DC, United States

⁵Department of Pediatrics, The George Washington University School of Medicine and Health Sciences, Washington, DC, United States

⁶Department of Integrative Systems Biology, The George Washington University School of Medicine and Health Sciences, Washington, DC, United States.

Abstract

Background—Allergic asthma is characterized by airway inflammation associated with recruitment and activation of eosinophils. In mice, allergen exposure induces platelet migration to the airways that is necessary for eosinophil recruitment and activation. We therefore hypothesized that in the airways of human subjects with asthma, (1) platelet activation would be positively associated with eosinophil activation and (2) platelet and eosinophil activation would both be associated with clinical asthma characteristics.

Methods—Nasal wash levels of p-selectin (a measure of platelet activation) and eosinophil cationic protein (ECP) (a measure of eosinophil activation) were compared with each other and with clinical asthma characteristics in a cross-sectional study of urban children and adolescents (6–20y) with asthma.

Results—Regression analysis revealed a significantly positive association between log₁₀ p-selectin levels and log₁₀ ECP levels (Beta=0.50 ng/mL [95%CI: 0.05 to 0.94 ng/mL]; P=0.029). Additionally, ECP was significantly and negatively associated with two asthma-related quality of life measurements and p-selectin was associated with one of these.

Conclusions—Our study shows the first significant association between platelet and eosinophil activation in airways of human subjects with asthma. These data provide a first step toward delineating what appears to be an important role for platelets in airway eosinophilia.

Corresponding Author: Robert J. Freishtat, MD, MPH Division of Emergency Medicine Children's National Medical Center 111 Michigan Avenue, NW Washington, DC 20010 (202) 476-2971 Fax (202) 476-6014 rfreishtat@cnmcresearch.org.

*These authors contributed equally to this manuscript.

Keywords

Asthma; Eosinophils; Platelets

INTRODUCTION

The airway inflammation seen in allergic asthma is associated with recruitment and activation of inflammatory cells (e.g. eosinophils, lymphocytes, mast cells) (1). Recent evidence has shown that activated platelets play a critical role in the development of inflammation in humans with allergic asthma as well as a variety of other disease states (i.e. rhinitis, COPD, rheumatoid arthritis, inflammatory bowel syndrome, atherosclerosis) (2).

In a mouse model of allergic asthma, allergen exposure induces platelet activation and migration to the airways (3-5) where they in turn attach to and activate leukocytes. Activated leukocytes show an increase in expression of CD11b and Very Late Antigen-4 (VLA-4), adhesion molecules that are necessary for inflammatory cell attachment to the airway vascular endothelium (3). The platelet-leukocyte complexes have also been shown in mice to be associated with eosinophil and lymphocyte activation and migration to the lungs where they subsequently adhere to the endothelial walls and cause inflammation (6). Additionally, it has also been shown that platelets play a critical role in airway remodeling as a result of chronic allergen exposure (7).

Therefore, we hypothesized that there would be a positive association between platelet activation and eosinophil activation in the airways of children and adolescents with asthma. Furthermore, we hypothesized that increased platelet and eosinophil activation would be associated with more severe clinical asthma characteristics.

METHODS**Study Cohort**

The subjects studied were enrolled as part of the ongoing Asthma Severity Modifying Polymorphisms (AsthMaP) Project, a cross-sectional study at Children's National Medical Center in Washington, DC. It is made up of 6-20 year old urban children and adolescents living in the DC metropolitan area with physician-diagnosed asthma present for at least one year. Details of the AsthMaP Project have been previously described (8, 9). Briefly, subjects were recruited in the emergency department at Children's National Medical Center and then returned to the Clinical Research Center for one study visit at least four weeks after completion of their most recent oral steroid dose. Study visits were conducted at approximately the same time of day. Informed consent and assent were obtained from participants and/or their guardians as appropriate. The study was approved by our Institutional Review Board.

Clinical Characteristics

Multiple historical and physiologic characteristics were assessed for each subject in the asthma cohort. Of note, nasal washes were performed by instilling isotonic sterile saline into each nare, holding for ten seconds, and then collecting the fluid in a specimen container. Eosinophils and neutrophils from these nasal washes were manually counted on slides stained with Wright's stain. Additionally, serum IgE was measured using chemiluminescence with an Immulite 2000 system (Siemens Healthcare Diagnostics, Deerfield, IL). Parental interviews were conducted using the Integrated Therapeutics Group's (ITG) Child Asthma Short Form (10), National Institutes of Health, National

Asthma Education and Prevention Program (NAEPP) 2007 criteria (11, 12), and, depending on age, either the Asthma Control Test™ (ACT) or Childhood Asthma Control Test™ (cACT) (QualityMetric Incorporated, Lincoln, RI).

P-Selectin as a Measurement of Platelet Activation

Soluble p-selectin (i.e. CD62P) is an α -granule membrane protein that migrates to the cell surface of platelets upon release of α -granules and subsequent platelet activation (13). It is a well established marker for platelet activation (14, 15). Levels of soluble p-selection were measured in nasal wash samples by flow cytometry on a FACSCalibur™ System (BD Biosciences, San Jose, CA) using a FlowCytomix Simplex Kit (Bender MedSystems, Burlingame, CA). Results were analyzed with FlowCytomix Pro 2.3 software (Bender MedSystems, Burlingame, CA).

Eosinophil Cationic Protein as a Measurement of Eosinophil Activation

Levels of eosinophil cationic protein (ECP), a cytotoxic protein released by activated eosinophils, were measured in nasal wash samples using a standard enzyme-linked immunosorbent assay (ELISA) kit (Medical & Biological Laboratories Co., Ltd., Nagoya, Japan) with a minimal detectable level of 0.125 ng/mL. ECP is a cytotoxic protein released by eosinophils following activation by an immune stimulus (16), and was thus chosen as the marker for eosinophil activation.

Statistical Analysis

Linear regression analysis was used to identify significant associations. Data was $\log_{10}$ transformed when not normally distributed. All beta coefficients and P values were corrected for age and gender. All statistical tests were performed with SPSS Statistics 17.0 (SPSS Inc., Chicago, IL).

RESULTS

At the time of this study, 109 children and adolescents had been enrolled in AsthMaP. Of those, 61% were male and the mean age (SE) was 11.4 (0.4) years. Of the 109 subjects, 100 (92%) were self-identified African Americans (AA), and 89 (82%) had persistent asthma as defined by NAEPP 2007 criteria (11). The median blood eosinophil level was 3.6% (interquartile range [IQR]: 1.8 to 6.8). A detailed description of select asthma characteristics can be found in Table 1.

Nasal wash samples were available from a subgroup of subjects (N = 75). There were no significant differences between these 75 subjects and the remaining 34 with respect to age, gender, and NAEPP classification. From those samples, p-selectin and ECP levels were measured in order to assess platelet and eosinophil activation, respectively. The median p-selectin level in the nasal washes was 0.83 ng/mL (IQR: 0.45 to 2.41) and the median ECP level in the nasal washes was 11.16 ng/mL (IQR: 1.27 to 46.77). Linear regression analysis showed that for every unit increase in $\log_{10}$ p-selection, there was a corresponding significant 0.5 unit increase in $\log_{10}$ ECP (Beta = 0.50 [95%CI: 0.05 to 0.94]; age and gender adjusted P = 0.029).

In addition, using regression analysis, we tested for associations between ECP and subjects' clinical asthma characteristics. We chose to analyze eight asthma-related quality of life and allergic markers that were relevant to allergic asthma and have previously been shown to be associated with eosinophil activation (17, 18). (Table 2) $\log_{10}$ ECP was found to be significantly and negatively associated with two asthma-related quality of life characteristics: (1) $\log_{10}$ ITG – nighttime score (−0.06 [−0.12 to 0.00]; adjusted P = 0.015)

and (2) $\log_{10}$ ITG – composite score (-0.04 [-0.08 to -0.01]; adjusted $P = 0.018$). In order to assess whether similar associations were present with p-selectin, regression analyses were carried out between $\log_{10}$ p-selectin and the same eight clinical asthma characteristics tested with ECP. P-selectin was found to be significantly and negatively associated with $\log_{10}$ ITG – nighttime score (-0.12 [-0.22 to -0.01]; adjusted $P = 0.035$).

We have previously identified, using cluster analysis, three distinct phenotypic clusters within the AsthMaP cohort (8). One of the clusters was characterized by an allergic asthma phenotype with increased nasal wash eosinophils and worse asthma control. Nasal washes were available for 25 participants within this cluster. As in the overall cohort, we used regression analysis to test for associations within this allergic phenotypic cluster between $\log_{10}$ ECP and the eight relevant asthma characteristics. The two significant associations previously identified in the overall cohort were also present within this cluster: $\log_{10}$ ITG – nighttime score (-0.14 [-0.26 to -0.02]; adjusted $P = 0.021$) and $\log_{10}$ ITG – composite score (-0.09 [-0.17 to -0.02]; adjusted $P = 0.020$). Additionally, within the cluster, both $\log_{10}$ ITG – daytime score (-0.28 [-0.53 to -0.04]; adjusted $P = 0.025$) and $\log_{10}$ ITG – nighttime score (-0.36 [-0.68 to -0.04]; adjusted $P = 0.029$) were found to be significantly associated with $\log_{10}$ p-selectin. Table 2 highlights these significant associations found both in the overall cohort and within the allergic phenotypic cluster.

DISCUSSION

We found a significant positive association between markers of platelet (i.e. p-selectin) and eosinophil (i.e. ECP) activation in the airways of children and adolescents with asthma. Additionally, p-selectin was significantly associated with one ECP-associated asthma-related quality of life measurement in the overall cohort. When looking within a previously identified allergic phenotypic cluster of the cohort, the same significant association was found along with an additional significant association between p-selectin and another asthma-related quality of life measurement.

It is well established that allergic asthma is associated with inflammation and airway epithelial damage (1). Specifically, following an allergic stimulus, inflammatory cell (e.g. eosinophils, lymphocytes, mast cells) activation and migration to asthmatic airways is followed by subsequent obstruction and inflammation (17, 18). Emerging literature has begun to explore the role of platelets in this inflammatory response (19). For example, activated platelets have been shown to be increased in asthmatic airways (20). Additionally, it has been demonstrated *in vitro* that, in the presence of platelets derived from patients with asthma, eosinophil attachment to airway endothelium is enhanced (21, 22). It has also been shown that mitogens and enzymes released by activated platelets play a direct role in the chronic inflammatory events that lead to airway remodeling in asthma (23).

Pitchford, et al. (6) showed in a mouse model of allergic asthma that p-selectin expression on activated platelets is necessary for contact-dependent leukocyte activation. Furthermore, they found an association between p-selectin expression on the surfaces of activated platelets and eosinophil and lymphocyte activation. These findings suggest that platelets play an indirect, albeit crucial, role on inflammatory cell activation and recruitment through the binding and activation of leukocytes. Our data extend these findings into human asthma by providing initial evidence for a positive association between platelet activation and eosinophil activation in human airways. Additionally, we show that eosinophil and platelet activation are both associated with asthma-related quality of life characteristics in a pediatric asthmatic cohort. Although we only uncovered a few significant associations due to the small sample size, the associations that we did find support the current literature showing that inflammation leads to increased symptoms and worse disease control (17, 18).

We had previously utilized cluster analysis to reduce the heterogeneity of the AsthMaP cohort and to identify distinct phenotypic clusters (8). In this current study we now show that the significant associations found in the overall cohort were stronger when analyzed within the allergic phenotypic cluster, despite its small sample size. Furthermore, one additional association between platelet activation and ITG – daytime score that was not present in the overall cohort was found to be significant within the cluster. Thus, looking within these allergic phenotypic cluster to identify new associations and strengthen previous associations supports the concept that distinct phenotypic subgroups will display different responses to stimuli (18).

This study has several limitations. First, this study had a relatively small sample size. However, despite the low power to detect significant associations, the relationships detected are important first steps to understanding the role of platelets in inflammation in human airways. Second, the use of ECP as a measure of eosinophil activity presents several concerns. The sticky nature of this protein, along with the fact that it is highly charged, makes recoverability difficult (24). Furthermore, eosinophils have been shown to release less intracellular ECP compared to other granule proteins (25). However, despite these inherent difficulties in measuring ECP, the significant association that we found between ECP and p-selectin provides initial evidence for the role of platelet activation in airway eosinophil activation and recruitment. Third, AsthMaP is comprised largely of AA children and adolescents. Therefore, it is difficult to extend our findings into other populations. However, it is important to note that our study provides insight into AA children and adolescents with asthma, which is one of the highest-risk asthma populations. Finally, a fourth limitation is our use of nasal washes in lieu of the gold-standard bronchoalveolar lavage. Because it has been shown in cystic fibrosis and respiratory syncytial virus (RSV) that inflammatory cell proportions in nasal samples accurately reflect what is observed in the lower airways (26, 27), we argue that these samples are valuable as a less invasive means to measure markers of airway inflammation.

In summary, this study extends to humans the current literature showing platelet involvement in airway inflammation in mice. Using a cohort of urban children and adolescents with asthma, we were able to identify a positive association between platelet activation and eosinophil activation in nasal secretions that had previously been demonstrated in the lower airways of a mouse model of allergic asthma. This finding provides a first step to better understanding the relationship between platelets and inflammation and should be further explored in order to better delineate the role of platelets in asthma.

Acknowledgments

Funding/Support: Funding support provided to RJF by Sheldon C. Siegel Investigator Award Grant from the Asthma and Allergy Foundation of America and by grants K23RR020069, P20MD000198, and M01RR020359 from the National Institutes of Health, Bethesda, Maryland.

REFERENCES

1. Barnes PJ. New aspects of asthma. *J Intern Med.* 1992; 231:453–461. [PubMed: 1602282]
2. Pitchford SC. Novel uses for anti-platelet agents as anti-inflammatory drugs. *Br J Pharmacol.* 2007; 152:987–1002. [PubMed: 17603547]
3. Pitchford SC, Yano H, Lever R, Riffo-Vasquez Y, Ciferri S, Rose MJ, Giannini S, Momi S, Spina D, O'Connor B, et al. Platelets are essential for leukocyte recruitment in allergic inflammation. *J Allergy Clin Immunol.* 2003; 112:109–118. [PubMed: 12847487]

4. Sullivan PJ, Jafar ZH, Harbinson PL, Restrict LJ, Costello JF, Page CP. Platelet dynamics following allergen challenge in allergic asthmatics. *Respiration*. 2000; 67:514–517. [PubMed: 11070455]
5. Yamamoto H, Nagata M, Tabe K, Kimura I, Kiuchi H, Sakamoto Y, Yamamoto K, Dohi Y. The evidence of platelet activation in bronchial asthma. *J Allergy Clin Immunol*. 1993; 91:79–87. [PubMed: 8423274]
6. Pitchford SC, Momi S, Giannini S, Casali L, Spina D, Page CP, Gresele P. Platelet p-selectin is required for pulmonary eosinophil and lymphocyte recruitment in a murine model of allergic inflammation. *Blood*. 2005; 105:2074–2081. [PubMed: 15528309]
7. Pitchford SC, Roffo-Vasquez Y, Sousa A, Momi S, Gresele P, Spina D, Page CP. Platelets are necessary for airway wall remodeling in a murine model of chronic allergic inflammation. *Blood*. 2004; 103:639–647. [PubMed: 14504080]
8. Benton AS, Wang Z, Lerner J, Foerster M, Teach SJ, Freishtat RJ. Overcoming heterogeneity in pediatric asthma: Tobacco smoke and asthma characteristics within phenotypic clusters in an african american cohort. *Journal of Asthma*. 2010 In Press.
9. Freishtat RJ, Iqbal SF, Pillai DK, Klein CJ, Ryan LM, Benton AS, Teach SJ. High prevalence of vitamin d deficiency among inner-city african american youth with asthma in washington, dc. *J Pediatr*. 156:948–952. [PubMed: 20236657]
10. Bukstein DA, McGrath MM, Buchner DA, Landgraf J, Goss TF. Evaluation of a short form for measuring health-related quality of life among pediatric asthma patients. *J Allergy Clin Immunol*. 2000; 105:245–251. [PubMed: 10669843]
11. Expert panel report 3 (epr-3): Guidelines for the diagnosis and management of asthma-summary report 2007. *J Allergy Clin Immunol*. 2007; 120:S94–138. [PubMed: 17983880]
12. Gorelick MH, Brousseau DC, Stevens MW. Validity and responsiveness of a brief, asthma-specific quality-of-life instrument in children with acute asthma. *Ann Allergy Asthma Immunol*. 2004; 92:47–51. [PubMed: 14756464]
13. Abrams CS, Ellison N, Budzynski AZ, Shattil SJ. Direct detection of activated platelets and platelet-derived microparticles in humans. *Blood*. 1990; 75:128–138. [PubMed: 2294986]
14. Conway DS, Pearce LA, Chin BS, Hart RG, Lip GY. Plasma von willebrand factor and soluble p-selectin as indices of endothelial damage and platelet activation in 1321 patients with nonvalvular atrial fibrillation: Relationship to stroke risk factors. *Circulation*. 2002; 106:1962–1967. [PubMed: 12370220]
15. O'Connor CM, Gurbel PA, Serebruany VL. Usefulness of soluble and surface-bound p-selectin in detecting heightened platelet activity in patients with congestive heart failure. *Am J Cardiol*. 1999; 83:1345–1349. [PubMed: 10235093]
16. Woschnagg C, Rubin J, Venge P. Eosinophil cationic protein (ecp) is processed during secretion. *J Immunol*. 2009; 183:3949–3954. [PubMed: 19692640]
17. Borish L, Culp JA. Asthma: A syndrome composed of heterogeneous diseases. *Ann Allergy Asthma Immunol*. 2008; 101:1–8. quiz 8-11, 50. [PubMed: 18681077]
18. Wenzel SE. Asthma: Defining of the persistent adult phenotypes. *Lancet*. 2006; 368:804–813. [PubMed: 16935691]
19. Pitchford SC. Defining a role for platelets in allergic inflammation. *Biochem Soc Trans*. 2007; 35:1104–1108. [PubMed: 17956289]
20. Moritani C, Ishioka S, Haruta Y, Kambe M, Yamakido M. Activation of platelets in bronchial asthma. *Chest*. 1998; 113:452–458. [PubMed: 9498966]
21. Jawien J, Chlopicki S, Gryglewski RJ. Interactions between human platelets and eosinophils are mediated by selectin-p. *Pol J Pharmacol*. 2002; 54:157–160. [PubMed: 12139113]
22. Ulfman LH, Joosten DP, van Aalst CW, Lammers JW, van de Graaf EA, Koenderman L, Zwaginga JJ. Platelets promote eosinophil adhesion of patients with asthma to endothelium under flow conditions. *Am J Respir Cell Mol Biol*. 2003; 28:512–519. [PubMed: 12654641]
23. Tutluoglu B, Gurel CB, Ozdas SB, Musellim B, Erturan S, Anakkaya AN, Kilinc G, Ulutin T. Platelet function and fibrinolytic activity in patients with bronchial asthma. *Clin Appl Thromb Hemost*. 2005; 11:77–81. [PubMed: 15678276]

24. Gleich GJ, Adolphson CR. The eosinophilic leukocyte: Structure and function. *Adv Immunol.* 1986; 39:177–253. [PubMed: 3538819]
25. Pronk-Admiraal CJ, Bartels PC. Total amount of ecp per eosinophil as indicator for the activity state of eosinophils. *Scand J Clin Lab Invest.* 2001; 61:453–457. [PubMed: 11681535]
26. Pitrez PM, Brennan S, Turner S, Sly PD. Nasal wash as an alternative to bronchoalveolar lavage in detecting early pulmonary inflammation in children with cystic fibrosis. *Respirology.* 2005; 10:177–182. [PubMed: 15823182]
27. Sheeran P, Jafri H, Carubelli C, Saavedra J, Johnson C, Krisher K, Sanchez PJ, Ramilo O. Elevated cytokine concentrations in the nasopharyngeal and tracheal secretions of children with respiratory syncytial virus disease. *Pediatr Infect Dis J.* 1999; 18:115–122. [PubMed: 10048682]

TABLE 1
CLINICAL CHARACTERISTICS OF SUBJECTS

Variable	Cases (n=109)
Gender, % male	60.6
Age, yr (SE)	11.4 (0.4)
Body mass index percentile (SE)	68.2 (2.5)
Age of asthma onset, yr (SE)	2.9 (0.3)
Pre-bronchodilator FEV1, % predicted (SE)	88.1 (1.8)
Post-bronchodilator FEV1, % predicted (SE)	93.1 (1.6)
ACT/cACT [†] (SE)	20.6 (0.3)
NAEPP [*] severity classification, %	
1 - <i>Intermittent</i>	11.0
2 - <i>Mild Persistent</i>	24.8
3 - <i>Moderate Persistent</i>	33.9
4 - <i>Severe Persistent</i>	30.3
ITG [‡] - composite score (SE)	75.3 (2.0)
Nasal wash eosinophils, % (IQR)	52 (2, 90)
Nasal wash neutrophils, % (IQR)	30 (3, 71)
Eosinophil cationic protein from nasal washes, ng/mL (IQR)	11.2 (1.3, 46.8)
Neutrophil elastase from nasal washes, ng/mL (IQR)	1.4 (0.6, 2.3)
P-selection from nasal washes, ng/mL (IQR)	0.8 (0.5, 2.4)
Blood eosinophils, % (IQR)	3.6 (1.8, 6.8)
Total serum IgE, IU/mL (IQR)	205.5 (72.3, 525.5)
One or more positive allergen skin prick tests, %	36.4

^{*} National Asthma Education and Prevention Program

[†] Asthma Control Test/Childhood Asthma Control TestTM

[‡] Integrated Therapeutics Group's Child Asthma Short Form

SE: Standard error of the mean

IQR: Interquartile Range

TABLE 2
ASSOCIATIONS BETWEEN ASTHMA CHARACTERISTICS AND EITHER EOSINOPHIL ACTIVATION OR PLATELET ACTIVATION OVERALL AND WITHIN AN ALLERGIC PHENOTYPIC SUBGROUP

Variables	All (n=75)		Allergic Subgroup (n=25)	
	Beta Coefficient ^{*,†} (95% CI)	P Value [†]	Beta Coefficient ^{*,†} (95% CI)	P Value [†]
ECP	ACT/cACT™	-0.41 (-1.41 to 0.58)	0.414	0.052
	ITG – daytime score [§]	-0.03 (-0.07 to 0.02)	0.195	0.052
	ITG – nighttime score [§]	-0.06 (-0.12 to 0.00)	0.015	0.021
	ITG – function score [§]	-0.04 (-0.07 to 0.00)	0.076	0.094
	ITG – composite score [§]	-0.04 (-0.08 to -0.01)	0.018	0.020
	Total serum IgE	0.05 (-0.12 to 0.22)	0.542	0.851
	Blood eosinophils, %	-0.03 (-0.14 to 0.07)	0.546	0.156
	Nasal wash eosinophils, %	-0.02 (-0.18 to 0.13)	0.774	0.973
	ACT/cACT™	0.10 (-2.13 to 2.32)	0.932	0.406
	ITG – daytime score [§]	0.08 (-0.02 to 1.18)	0.115	0.025
P-selectin	ITG – nighttime score [§]	-0.12 (-0.22 to -0.01)	0.035	0.029
	ITG – function score [§]	0.02 (-0.07 to 0.10)	0.718	0.800
	ITG – composite [§]	0.00 (-0.08 to 0.08)	0.979	0.106
	Total serum IgE	0.04 (-0.33 to 0.42)	0.829	0.903
	Blood eosinophils, %	0.05 (-0.19 to 0.28)	0.692	0.147
	Nasal wash eosinophils, %	0.23 (-0.12 to 0.58)	0.201	0.428

Abbreviations: ACT/cACT = Asthma Control Test/Childhood Asthma Control Test™, ITG = Integrated Therapeutics Group's Child Asthma Short Form.

* Beta coefficient is for each log₁₀ increase in either ECP or p-selectin.

[†] Adjusted for age and gender. P Values <0.05 are listed in **bold**.

[§] Variables were log transformed to approximate a normal distribution.