Hill's small systems nanothermodynamics: a simple macromolecular partition problem with a statistical perspective

Hong Qian

Received: 27 October 2011 / Accepted: 7 December 2011 /

Published online: 6 January 2012

© Springer Science+Business Media B.V. 2012

Abstract Using a simple example of biological macromolecules which are partitioned between bulk solution and membrane, we investigate T.L. Hill's phenomenological nanothermodynamics for small systems. By introducing a *system size-dependent* equilibrium constant for the bulk-membrane partition, we obtain Hill's results on differential and integral chemical potentials μ and $\hat{\mu}$ from computations based on standard Gibbsian equilibrium statistical mechanics. It is shown that their difference can be understood from an equilibrium re-partitioning between bulk and membrane fractions upon a change in the system's size; it is closely related to the system's fluctuations and inhomogeneity. These results provide a better understanding of nanothermodynamics and clarify its logical relation with the theory of statistical mechanics.

Keywords Nanothermodynamics • Ensemble • Fluctuation • Small systems • Statistical mechanics

1 Introduction

With great advances in material preparations at nanoscale in recent years, there is a growing interest in the thermodynamics of small systems [1, 2]. Nanothermodynamics, a phenomenological theory developed by T.L. Hill in the early 1960s for small equilibrium systems, has become one of the major formalisms for quantitative treatment of equilibrium nanoscale materials [3, 4]. There are many experimental and computational works that have verified the theory (see for example [5, 6]).

The greatest strength of thermodynamics, being absolute and abstract, is often its weakness: It usually provides no molecular mechanism or insights into chemical and biochemical processes. The same dichotomy applies to Hill's nanothermodynamics: The

H. Qian (✉)
Department of Applied Mathematics, University of Washington,
Seattle, WA 98195-2420, USA
e-mail: hqian@u.washington.edu

key concept in the theory is a difference between *differential* and *integral* forms of many non-extensive thermodynamic quantities due to the smallness of a system. For example, since the Gibbs free energy $G(N, p, T)$ is not linearly proportional to the system's size N , $\mu = \partial G / \partial N$ and $\hat{\mu} = G / N$ are different. The difference is intuitively understood: It is due to contributions such as surface effects.

In biophysical chemistry, biological macromolecules are often treated as having discrete conformational states, and interaction with ligands as stoichiometric binding. While this approach is only an approximation, it often provides much more powerful intuitive understanding of the physical processes that underly molecular thermodynamics. A case in point is J.A. Schellman's theory of a three-component system with mixed solvents [7]. While a thorough treatment of this problem involves preferential interaction coefficients, a discrete stoichiometric binding model has provided greater intuitions [8].¹

The fundamental premise of a *small system* is that its thermodynamic properties are not strictly proportional to the system's size [4]. A system's size, however, can be represented by several different quantities: The number of molecules N , the volume V , or the total energy E . In chemistry and biochemistry, the concentration $x = N / V$ is a widely used intensive quantity. In the rest of the paper, we shall refer to a system's size in terms of its volume. In terms of V , the Helmholtz free energy F of a three-dimensional small system could have the form

$$F = V f(x, T) + a(x, T) V^{2/3} + b(x, T) \ln V + \cdots, \quad (1)$$

where the term in $V^{2/3}$ is associated with a surface energy.

In the present work, we first consider the following simple problem: N non-interactive macromolecules M are in a system which consists of bulk solution and membrane (one could think of a vesicle with lipid membrane). Each M molecule has three discrete conformational states: 0, 1, in the bulk and 2 on the membrane. We assume the equilibrium constant between states 0 and 1 is K_1 and between states 0 and 2 is K_2 . $K_1 = p_1 / p_0$ and $K_2 = p_2 / p_0$ where p_i is the probability of an M being in state i in equilibrium. K_1 is independent of the amount of membrane in the system, but K_2 is critically dependent on the ratio of membrane surface S to bulk volume V , thus the system's size. Assuming simple geometry with surface-to-volume ratio $S / V = V^{-1/3}$ and homogeneity in both bulk and membrane phases, it is reasonable to assume that $K_2(V) \propto V^{-1/3}$. Thus, the equilibrium partition function for a single M molecule, with the state 0 as the reference state, is thus:

$$Z_1 = 1 + K_1 + K_2(V) \rightarrow 1 + K_1 \quad (2)$$

in the limit of $V \rightarrow \infty$. This simple example can be easily generalized to more general macromolecular partition problems, and a key formula relating μ and $\hat{\mu}$, which first appeared in [4], will be derived.

A few remarks on (2) are in order: (i) Z_1 looks like a binding polynomial, but it is actually independent of the concentration of the species M . This is because we have assumed that the N macromolecules are non-interactive. Hence, Z_1 is the "partition function" of a single M molecule. The partition function for the whole system is $Z_N = Z_1^N$.

(ii) $K_2(V) = p_2 / p_0$ as a function of V is itself a complex problem in general. For our simple model, we shall completely neglect the surface free energy of the membrane.

¹A difference between differential and integral forms of macromolecular interaction also appears in the theory of binding: Thermodynamic binding can go to zero while molecular interaction is maximized [9].

Furthermore, if one treats macromolecule-membrane association as simple Langmuir absorption, with $N(p_0 + p_1)/V$ and Np_2/S being the concentrations of the species M in the bulk and on the membrane, then one obtains $p_2/(p_0 + p_1) \propto S/V$. Since $(p_0 + p_1) = (1 + K_1)p_0$, one has $p_2/p_0 \propto S/V \sim V^{-1/3}$. For the present analysis, the key is that $K_2(V)$ is a function of V . The result in (6) given below does not depend on a specific functional form of $K_2(V)$.

The novel idea of the present work is to introduce the *system size-dependent* equilibrium constant(s), (i.e., more generally, the potential of mean force or conditional free energy). It is shown that if one starts with a mechanistic-based statistical approach, following Gibbs, to small systems, then the small system's nanothermodynamics naturally emerges.

One of the salient features of Hill's nanothermodynamics is its *ensemble dependence*: When a system's boundary becomes significant, how a statistical thermodynamic system is maintained at the boundary matters. In the past, dependence upon details of an ensemble at its boundary leads to the phenomenon of *entropy-enthalpy compensation* [10, 11], which can be understood from relaxing certain internal constraints. In Sect. 3.1 we show that this perspective is also fruitful in understanding the difference between the differential and integral chemical potentials μ and $\hat{\mu}$.

2 Bulk-membrane partition: a simple example

We now consider a simple system in which there are N non-interactive macromolecules M . Each M has three distinct states: states 0 and 1 are in the bulk solution and state 2 is associated with the membrane. It is clear that the ratio between total membrane to total bulk volume is a function of the system's size V : Thus by simple geometric considerations $K_2(V)$ is volume-dependent. The partition function for the small system with constant pressure p and temperature T is:

$$Z(N, p, T) = (1 + K_1 + K_2(V))^N, \quad (3)$$

in which $V = V(N, p, T)$ is a function of N .² Therefore,

$$G(N, p, T) = -k_B T \ln Z(N, p, T), \quad (4)$$

$$\hat{\mu}(N, p, T) = \frac{G(N, p, T)}{N} = -k_B T \ln (1 + K_1 + K_2), \quad (5)$$

$$\mu(N, p, T) = \frac{\partial G(N, p, T)}{\partial N} = \hat{\mu} - \frac{k_B T K_2 \left(\frac{\partial \ln K_2}{\partial \ln N} \right)_{p, T}}{1 + K_1 + K_2}, \quad (6)$$

where $\hat{\mu}$ and μ are the integral and differential chemical potentials introduced by Hill [4]. In the last term of (6), $(\partial \ln K_2 / \partial \ln N)$ is an intensive quantity, $K_2 \propto V^{-1/3}$ and $1 + K_1 + K_2 \rightarrow 1 + K_1$. Hence it is on the order of $K^{-1/3}$. Thus, μ and $\hat{\mu}$ are the same for a macroscopic system when $G(N, p, T)$ asymptotically becomes an "extensive" quantity: $G \propto N$.

²To be mathematically more rigorous, one needs to solve the problem with self-consistency among (3), (4) and: $V(N, p, T) = (\partial G / \partial p)_{N, T}$.

Equation (6) can be easily generalized into

$$\mu(N, p, T) - \hat{\mu}(N, p, T) = \left\langle \left(\frac{\partial \mu_i^o(N, p, T)}{\partial \ln N} \right)_{p,T} \right\rangle = \left(\frac{\partial \hat{\mu}(N, p, T)}{\partial \ln N} \right)_{p,T}, \quad (7)$$

where

$$\mu_i^o(N, p, T) = -k_B T \ln K_i(N, p, T) \quad (8)$$

and the average $\langle \cdots \rangle$ is performed with respect to $i = 0, 1, 2$. Equation (7) clearly shows why and how the small system's thermodynamics depends on the nature of the ensemble; and why the difference of μ and $\hat{\mu}$ will be ensemble-specific. Equation (7) agrees with what was obtained by Hill ([4], Part I, p. 30, Eqs. 2-12):

$$\mu - \hat{\mu} = \left(\frac{\partial \hat{\mu}}{\partial \ln N} \right)_{p,T}. \quad (9)$$

3 Thermodynamic hierarchy and nanothermodynamics

3.1 Perturbation and relaxation in thermodynamic systems

The chemical potential μ is the increment in free energy of a system when its number of molecules increases by 1 [12]. Therefore, it can be considered as a *perturbation* to the thermodynamic system. The idea advanced in [10], originally proposed by A. Ben-Naim [13], is to view the response of a thermodynamic system to a perturbation as two parts:

- (i) Constraining the distribution among all the populations unchanged, what is the increase in free energy of the system due to the perturbation?
- (ii) Inevitably, however, the perturbation will cause a shift (i.e., re-distribution, re-partitioning) in the equilibrium among the populations. There is an additional free energy change associated with this re-organization process.

This perturbation-relaxation view of the thermodynamics of complex systems yielded a possible explanation for the so-called “entropy-enthalpy compensation” phenomenon [10, 11, 13]: Note that the chemical potentials for all the subcomponents μ_j of an equilibrium system have to be the same. Thus, the free energy change associated with (ii) is essentially zero if the perturbation is infinitesimal. In contrast, the corresponding entropy and enthalpy of the subcomponents s_j and h_j (where $h_j - Ts_j = \mu_j$), can be very different. Thus, large contributions to entropy and enthalpy from the process in (ii) are expected.³

Applying the same idea to the thermodynamics of our simple example, a perturbation leads to a shift in equilibrium between the “bulk” and “membrane” fractions. More specifically, introducing an additional M molecule causes a change in the system's volume V under isobaric (i.e., constant p) conditions. This change in V is global: It causes the change of K_2 for each and every molecule M in the system.

³Both steps in this gedankenexperiment are carried out under isothermal conditions. They cannot be realized in a laboratory. They are different from the adiabatic (isoentropic) and isothermal processes in the derivation of the fundamental equation of thermodynamics.

3.2 Spatial inhomogeneity

“Size” is a geometric concept. The unique feature of the nanothermodynamics of small systems is *spatial inhomogeneity*, which partitions the particles in a system into different populations whose proportions scale differently with system size. In Hill's theory, this partition is implicit: cluster growth naturally leads to “surface” and “interior”; in our simple model, this partition is explicitly assumed: states 0 and 1 versus 2.

With this geometry in mind, there will be two answers to the following question: What is the change in system's free energy if one introduces one additional particle into the system? Accordingly, answer (i) assumes that the proportion is unchanged, and answer (ii) assumes there is a shift in equilibrium distributions between different populations.

The answer (i) is precisely the macroscopic result, $\hat{\mu}$. It is intimately related to the free-energy perturbation theory [14, 15]:

$$\Delta G^o(A \rightarrow B) = -k_B T \ln \left\langle \exp \left(-\frac{E_B - E_A}{k_B T} \right) \right\rangle_A, \quad (10)$$

in which the average is carried out in terms of the equilibrium distribution of the unperturbed system. Note that (10) assumes that the phase space, over which the ensemble average $\langle \cdots \rangle_A$ is carried out, is the same before and after the perturbation.

The answer, (ii) however, contains exactly a contribution from the small systems effect: The addition of a particle changes the size of the system, and thus the phase space, on the order of $1/N$. This change causes a shift in the equilibrium from “surface population” to “bulk population”, which affects all N particles in the system. This is given in (7).

Realizing the difference between these two answers led Hill to introduce dN_i and $d\mathcal{N}$. The former corresponds to a change in the total number of particles, and thus gives μ (answer ii); the latter changes the number of particles but keeps the equilibrium population proportion by “adding an independent copy of the small system”, thus giving $\hat{\mu}$ (answer i).

3.3 Rapidly stirred biochemical reaction systems

For a small biochemical reaction system that is rapidly stirred, the theory of Delbrück-Gillespie process [16, 17], with its probability distribution given by the chemical master equation and its trajectories following the Gillespie algorithm, predicts that its stationary probability distribution, in the limit of large size, has the generic form

$$f(x) \propto \exp(-V\phi(x)), \quad (11)$$

where x is the concentration of biochemical species, V is the system's volume, and $\phi(x)$ is a function of only intensive quantities. Applying Laplace's method for integrals [18, 19], this result leads to the conclusion that, for a rapidly stirred system without geometric inhomogeneity, the partition function is

$$\begin{aligned} Q(V, \mu, T) &= V \int dx e^{-V\phi(x)} \\ &\approx V e^{-V\phi(x^*)} \int dx e^{-\frac{V}{2}\phi''(x^*)(x-x^*)^2} \\ &= e^{-V\phi(x^*)} \sqrt{\frac{2\pi V}{\phi''(x^*)}}, \end{aligned} \quad (12)$$

where $x^*(\mu)$ is the macroscopic concentration. Therefore,

$$-k_B T \ln Q(V, \mu, T) = k_B T V \phi(x^*) + O(\ln V), \quad (13)$$

where $O(\dots)$ is the mathematical symbol called Bachmann-Landau notation. It stands for “on the order of $\dots$ ”. Equation (13) should be compared with (3): $-k_B T \ln Z(N, p, T) = -k_B T V x^* \ln(1 + K_1) + O(V^{2/3})$ where $x^* = N/V$. For rapidly stirred nanoscale systems, the correction to the extensive term is on the order of the logarithm of an extensive quantity: $\ln V$ or $\ln N$. When there is a spatial partitioning, e.g., compartmentalization, however, terms on the order of a fractional power of the extensive quantity arise. For a system with a bulk-membrane partition, this term is on the order of $V^{2/3}$.

A biological cell is a small thermodynamic system. Applying the present result to a collection of cells, one realizes that there are two macroscopic limits: (a) classical biochemistry, which studies cell extracts with membrane removed; (b) permeable cells that preserve the ratio between membrane surface area to bulk volume. They correspond precisely the two ensembles of Hill.

3.4 Kinetics of re-partitioning

All of the above discussion has been based exclusively on equilibrium thermodynamics. We suspect there is in fact a kinetic aspect of the thermodynamics of small systems. Nonequilibrium thermodynamics for heterogeneous systems and for nanoscale systems have become natural subjects in recent years [20–22]. If one takes a kinetic perspective of the response of a small system upon perturbation [10], one sees that the conformation re-partitioning between the bulk and surface could indeed be observed kinetically. This connection between thermodynamics and kinetics deserves further investigation [23–25].

Acknowledgements I thank Prof. Dick Bedeaux for a stimulating week of discussions in June 2008 at XXI Sitges Conference (Spain), Profs. Ralph Chamberlin and Signe Kjelstrup for discussions that stimulated and renewed my interest in nanothermodynamics. I would like to acknowledge my friend and teacher (via his writing), Terrell Hill, whose work has had a major influence on mine, not only in nanothermodynamics, but also in nonequilibrium steady-state cycle kinetics and theory of muscle contraction [26–28].

References

1. Bustamante, C., Liphardt, J., Ritort, F.: The nonequilibrium thermodynamics of small systems. *Phys. Today* **58**, 43–48 (2005)
2. Rajagopal, A.K., Pande, C.S., Abe, S.: Nanothermodynamics—a generic approach to material properties at nanoscale. arxiv.org/abs/cond-mat/0403738 (2004)
3. Hill, T.L.: A different approach to nanothermodynamics. *Nano Lett.* **1**, 273–275 (2001)
4. Hill, T.L.: *Thermodynamics of Small Systems*. Dover, New York (1994)
5. Chamberlin, R.V.: Mean-field cluster model for the critical behaviour of ferromagnets. *Nature* **408**, 337–339 (2000)
6. Schnell, S.K., Vlugt, T.J.H., Simon, J.-M., Bedeaux, D., Kjelstrup, S.: Thermodynamics of a small system in a μT reservoir. *Chem. Phys. Lett.* **504**, 199–201 (2011)
7. Timasheff, S.N.: Thermodynamic binding and site occupancy in the light of the Schellman exchange concept. *Biophys. Chem.* **101–102**, 99–111 (2002)
8. Schellman, J.A.: A simple model for salvation in mixed solvents: applications to the stabilization and destabilization of macromolecular structures. *Biophys. Chem.* **37**, 121–140 (1990)
9. Schellman, J.A.: The relation between the free energy of interaction and binding. *Biophys. Chem.* **45**, 273–279 (1993)
10. Qian H., Hopfield, J.J.: Entropy-enthalpy compensation: perturbation and relaxation in thermodynamic systems. *J. Chem. Phys.* **105**, 9292–9298 (1996)

11. Qian, H.: Entropy-enthalpy compensation: conformational fluctuation and induced-fit. *J. Chem. Phys.* **109**, 10015–10017 (1998)
12. Ben-Naim, A.: *Statistical Thermodynamics for Chemists and Biochemists*. Plenum, New York (1992)
13. Ben-Naim, A.: Hydrophobic interaction and structural changes in the solvent. *Biopolymers* **14**, 1337–1355 (1975)
14. Zwanzig, R.W.: High-temperature equation of state by a perturbation method. I. Nonpolar gases. *J. Chem. Phys.* **22**, 1420–1426 (1954)
15. Widom, B.: Some topics in the theory of fluids. *J. Chem. Phys.* **39**, 2808–2812 (1963)
16. Qian, H.: Nonlinear stochastic dynamics of mesoscopic homogeneous biochemical reaction systems: an analytical theory. *Nonlinearity* **24**, R19–R49 (2011)
17. Qian, H.: Cellular biology in terms of stochastic nonlinear biochemical dynamics: emergent properties, isogenetic variations and chemical system inheritability. *J. Stat. Phys.* **141**, 990–1013 (2010)
18. Qian, H.: An asymptotic comparative analysis of the thermodynamics of non-covalent association. *J. Math. Biol.* **52**, 277–289 (2006)
19. Murray, J.D.: *Asymptotic Analysis*. Springer, Berlin (1984)
20. Kjelstrup, S., Bedeaux, D.: *Non-equilibrium Thermodynamics of Heterogeneous Systems*. World Scientific, Singapore (2008)
21. Reguera, D., Rubí, J.M., Vilar, J.M.G.: The mesoscopic dynamics of thermodynamic systems. *J. Phys. Chem. B* **109**, 21502–21515 (2005)
22. Ge, H., Qian, H.: The physical origins of entropy production, free energy dissipation and their mathematical representations. *Phys. Rev. E* **81**, 051133 (2010)
23. Hill, T.L., Chamberlin, R.V.: Extension of the thermodynamics of small systems to open metastable states: an example. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 12779–12782 (1998)
24. Rubí, J.M., Bedeaux, D., Kjelstrup, S.: Thermodynamics for single-molecule stretching experiments. *J. Phys. Chem. B* **110**, 12733–12737 (2006)
25. Rubí, J.M., Bedeaux, D., Kjelstrup, S.: Unifying thermodynamic and kinetic descriptions of single-molecule processes: RNA unfolding under tension. *J. Phys. Chem. B* **111**, 9598–9602 (2007)
26. Qian, H.: A simple theory of motor protein kinetics and energetics. *Biophys. Chem.* **67**, 263–267 (1997)
27. Qian, H.: Cycle kinetics, steady-state thermodynamics and motors—a paradigm for living matter physics. *J. Phys. Condens. Matter* **17**, S3783–S3794 (2005)
28. Qian, H.: Open-system nonequilibrium steady-state: statistical thermodynamics, fluctuations and chemical oscillations. *J. Phys. Chem. B* **110**, 15063–15074 (2006)