Published in final edited form as:

Org Lett. 2012 March 16; 14(6): 1378–1381. doi:10.1021/ol300045c.

An Asymmetric Synthesis of L-Pyrrolysine

Margaret L. Wong[†], Ilia A. Guzei[†], and Laura L. Kiessling^{†,‡,*}

[†]Department of Chemistry, University of Wisconsin—Madison, Madison, Wisconsin 53706

[‡]Department of Biochemistry, University of Wisconsin—Madison, Madison, Wisconsin 53706

Abstract

An efficient asymmetric synthesis of the 22nd amino acid L-pyrrolysine has been accomplished. The key stereogenic centers were installed by an asymmetric conjugate addition reaction. A Staudinger/aza-Wittig cyclization was used to form the acid-sensitive pyrroline ring. Pyrrolysine was synthesized in thirteen steps in 20% overall yield.

Pyrrolysine is the 22nd genetically encoded amino acid.¹ It consists of a (4*R*,5*R*)-4-methyl-5-carboxypyrroline ring linked to the ϵ -nitrogen of L-lysine.^{2,3} Pyrrolysine was identified by x-ray crystallography, when it was first observed in the structure of *Methanosarcina barkeri* monomethylamine methyltransferase.¹ It has been hypothesized that this unique amino acid plays an important role in methane production in some archaeal species.⁴

Pyrrolysine is encoded by an in-frame UAG codon, which is non-terminating, and its incorporation is mediated by a dedicated tRNA and cognate tRNA synthetase.⁵ The mechanism of pyrrolysine incorporation into proteins offers a platform for developing new protein labeling methods.⁶ Small molecules, such as fluorophores,^{6a} biotin,^{5e} ubiquitin^{6b} and oligosaccharides,^{6c} have been used for labeling pyrrolysine and pyrrolysine-surrogate residues.⁷ Although pyrrolysine has been exploited for protein labeling, an understanding of its biosynthesis,^{5e,8} evolutionary purpose, and distribution in the proteome is incomplete. New synthetic methods that provide sufficient quantities of this novel amino acid would facilitate an understanding of pyrrolysine biochemistry.

Pyrrolysine is a synthetic target.^{2,9} One reported chemical syntheses² involves the coupling of (4*R*,5*R*)-4-methyl-pyrroline-5-carboxylic acid to lysine. While this innovative route affords the desired product, the yield is modest (9% overall). Moreover, we and others^{5a,9} have found that the reactions were irreproducible. We therefore sought to devise an alternative synthetic route.

A drawback of the reported pyrrolysine synthesis is that it entails exposure of the sensitive carboxypyrroline ring to strongly acidic conditions, both during imine formation and lysine

kiessling@chem.wisc.edu.

Supporting Information Available. Experimental procedures and full characterization of compounds **9–15** and associated intermediate compounds and X-ray crystallographic data (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

coupling. We focused on generating the pyrroline ring in the penultimate step of the synthesis. In this way, we hoped to avoid reactions that could lead to pyrroline ring degradation. Specifically, we envisioned forming the cyclic imine from azido aldehyde **2** via a tandem Staudinger/aza-Wittig reaction¹⁰ (Scheme 1). Azido aldehyde **2** could be obtained from coupling of **3** with a selectively protected lysine derivative. We considered two different approaches to compound **3**: an asymmetric conjugate addition of a methylcarbanion followed by α -substitution ("Route I") or a Michael addition with a glycine enolate ("Route II").

We pursued these strategies in parallel. For Route I, we anticipated the methyl group of pyrrolysine precursor **3** could be installed by conjugate addition to a compound such as **4**. The Feringa group has developed efficient catalysts for this type of transformation.¹¹ Accordingly, the Josiphos catalyst gave rise to the 1,4-addition of methyl Grignard to α,β -unsaturated thioester **4** (Scheme 2). Attempts to functionalize the α -position with an electrophilic azide source or by an asymmetric α -halogenation¹² and subsequent nucleophilic displacement with an azide¹³ did not afford the desired products. A more significant drawback of this approach was our observation that the most robust α -halogenation methods¹⁴ were incompatible with the thioester required for the Josiphos-catalyzed reaction. As a result, the conjugate addition and the selective halogenation reactions demanded different carbonyl derivatives (thioester versus aldehyde), thereby necessitating multiple changes in oxidation state.

The roadblocks encountered pursuing Route I prompted us to focus on Route II. In this approach, we planned to install the pyrroline stereochemistry by carrying out Michael addition of a glycine enolate in the presence of a chiral catalyst. Calcium ion-promoted conjugate addition of *N*-(*t*-butylphenyl-methylidene)glycine *t*-butyl ester to methyl crotonate produced **9**¹⁵ in 92% yield with 99% enantiomeric excess and greater than 99:1 diastereomeric excess (Scheme 3). This reaction was not only stereoselective and efficient; it also was robust. These favorable attributes were preserved even when the reactions were conducted on a medium scale (50–100 mmol).

While the Schiff base was necessary for the high stereoselectivity attained,¹⁶ the imine was prone to hydrolysis upon exposure to acid or base. Accordingly, the *t*-butyl-phenyl imine **9** was hydrolyzed, and the unmasked amine was immediately exposed to di-*t*-butyl dicarbonate. We anticipated the *t*-butyl carbamate could be removed later concomitantly with *t*-butyl ester hydrolysis. The methyl ester was then saponified to afford **10**. The three-step interconversion of protecting groups was executed in an overall yield of 86%. The series of transformations proceeded with no loss of stereochemical integrity, as revealed by X-ray crystallography (Figure 1). The structure of acid **10** confirmed it had the expected absolute stereochemistry.

At this point in the synthesis, we recognized that differentiation of the two carboxy termini would be crucial for accessing lysine-coupled azido aldehyde **2** (Scheme 1). To this end, the carboxylic acid was transformed to a mixed anhydride, which was reduced with sodium borohydride to afford hydroxy *t*-butyrate **11** in 73% yield. The resulting primary hydroxyl group was converted to a silyl ether to mitigate side reactions with a free hydroxyl group. The silyl ether **12** was generated under mildly basic conditions as these avoided lactone formation. To minimize the formation of a disiloxane byproduct that co-eluted with the product during chromatography, the reaction was quenched under non-aqueous conditions. Using these conditions, compound **12** could be readily obtained.

Each functional group in ester **12** (Scheme 4) is poised for manipulation to access compound **14**, the precursor for the tandem sequence. As anticipated, the *t*-butyl ester and carbamate

protecting groups could both be removed in a single step. The azide group was installed using a Zn^{2+} -mediated diazotransfer from triflyl azide,¹⁷ which afforded coupling partner **13** in 75% yield over 2 steps.¹⁸ The precursor to pyrroline ring, α -azido carboxylic acid **13**, was appended to the epsilon nitrogen of *N*-trifluoroacetamidyl lysine methyl ester.² The silyl protecting group was cleaved, and the resulting primary alcohol was converted to an aldehyde by the Swern oxidation.¹⁹ This sequence provided compound **14** as the substrate for the Staudinger/aza-Wittig reaction.

Pyrroline **15** was generated using an efficient sequence that minimized purification steps. The reduction of the azide moiety was effected using polystyrene-supported triphenylphosphine. The putative iminophosphorane underwent cyclization to afford protected pyrrolysine derivative **15**. Solution-phase triarylphosphine reagents have been used previously to execute tandem Staudinger reduction and intramolecular aza-Wittig reactions.²⁰ The popularity of this strategy is gaining, presumably because of the efficiency and simplicity it affords for generating even highly sensitive imines such as **15**. The use of an immobilized phosphine to effect the transformation offers an additional advantage. It circumvents the need for chromatography, which can be a major benefit in preparing sensitive imines. To generate the final pyrrolysine product, we removed the protecting groups at the *N*- and *C*-termini² of pyrroline **15** to afford the lithium salt of L-pyrrolysine **16** in 20% overall yield.

The sequence described is high yielding and stereoselective. It affords L-pyrrolysine in 13 steps and 20% overall yield. Through our examination of multiple strategies for the installation of the two stereogenic centers of pyrrolysine, we identified the Ca^{2+} -mediated conjugate addition reaction as an efficient process for generating pyrroline precursors. Another feature of our route is the gentle conditions offered by a polymer-supported phosphine to effect the Staudinger/aza-Wittig cyclization. We anticipate that the mild conditions used to form the pyrroline can be used to generate pyrrolysine analogs. Our route will also be valuable for synthesizing isotopically labeled pyrrolysine to explore the prevalence of this intriguing amino acid in microorganisms and to carry out proteomic analyses.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We acknowledge Dr. Jiyoung Chang of the Kiessling group for helpful discussions and for her preliminary synthetic studies. This research was supported by NIGMS (R01 GM055984). The NMR facilities at UW-Madison are funded by the NSF (CHE-9208463) and NIH (RR08389-01).

References

1. (a) Hao B, Gong W, Ferguson TK, James CM, Krzycki JA, Chan MK. Science. 2002; 296:1462. [PubMed: 12029132] (c) Srinivasan G, James CM, Krzycki JA. Science. 2002; 296:1459. [PubMed: 12029131]
2. Hao B, Zhao G, Kang PT, Soares JA, Ferguson TK, Gallucci J, Krzycki JA, Chan MK. Chem Biol. 2004; 11:1317. [PubMed: 15380192]
3. Soares JA, Zhang L, Pitsch RL, Kleinholz NM, Jones RB, Wolff JJ, Amster J, Green-Church KB, Krzycki JA. J Biol Chem. 2005; 280:36962. [PubMed: 16096277]
4. Hand CE, Honek JF. J Nat Prod. 2005; 68:293. [PubMed: 15730267] Rother M, Krzycki JA. Archaea. 2010; 2010

5. (a) Blight SK, Larue RC, Mahapatra A, Longstaff DG, Chang E, Zhao G, Kang PT, Green-Church KB, Chan MK, Krzycki JA. *Nature*. 2004; 431:333. [PubMed: 15329732] (b) Polycarpo C, Ambrogelly A, Berube A, Winbush SM, McCloskey JA, Crain PF, Wood JL, Soll D. *Proc Natl Acad Sci USA*. 2004; 101:12450. [PubMed: 15314242] (c) Kobayashi T, Yanagisawa T, Sakamoto K, Yokoyama S. *J Mol Biol*. 2009; 385:1352. [PubMed: 19100747] (d) Namy O, Zhou Y, Gundllapalli S, Polycarpo CR, Denise A, Rousset JP, Soll D, Ambrogelly A. *FEBS Lett*. 2007; 581:5282. [PubMed: 17967457] (e) Longstaff DG, Larue RC, Faust JE, Mahapatra A, Zhang L, Green-Church KB, Krzycki JA. *Proc Natl Acad Sci USA*. 2007; 104:1021. [PubMed: 17204561]
6. (a) Fekner T, Li X, Lee MM, Chan MK. *Angew Chem Int Ed Engl*. 2009; 48:1633. [PubMed: 19156778] (b) Li X, Fekner T, Ottesen JJ, Chan MK. *Angew Chem Int Ed Engl*. 2009; 48:9184. [PubMed: 19882608] (c) Nguyen DP, Lusic H, Neumann H, Kapadnis PB, Deiters A, Chin JW. *J Am Chem Soc*. 2009; 131:8720. [PubMed: 19514718] (d) Ou W, Uno T, Chiu HP, Grinewald J, Cellitti SE, Crossgrove T, Hao X, Fan Q, Quinn LL, Patterson P, Okach L, Jones DH, Lesley SA, Brock A, Geierstanger BH. *Proc Natl Acad Sci USA*. 2011; 108:10437. [PubMed: 21670250]
7. Nguyen DP, Elliott T, Holt M, Muir TW, Chin JW. *J Am Chem Soc*. 2011; 133:11418. [PubMed: 21736333]
8. Gaston MA, Zhang L, Green-Church KB, Krzycki JA. *Nature*. 2011; 471:647. [PubMed: 21455182]
9. Bérubé, A. Progress Toward the Total Syntheses of the Polycyclic Terpenes Providencin and Bacchopetiolone, and, Study of Pyrrolisine. Yale University; 2006.
10. (a) Mulzer J, Meier A, Buschmann J, Luger P. *Synth*. 1996:123.(b) Molina P, Vilaplana MJ. *Synth*. 1994; 1994:1197.
11. Ruiz BM, Geurts K, Fernández-Ibáñez MÁ, ter Horst B, Minnaard AJ, Feringa BL. *Org Lett*. 2007; 9:5123. [PubMed: 17975924]
12. (a) Wack H, Taggi AE, Hafez AM, Drury WJ, Lectka T. *J Am Chem Soc*. 2001; 123:1531. [PubMed: 11456741] (b) Brochu MP, Brown SP, MacMillan DWC. *J Am Chem Soc*. 2004; 126:4108. [PubMed: 15053591] (c) Halland N, Braunton A, Bachmann S, Marigo M, Jorgensen KA. *J Am Chem Soc*. 2004; 126:4790. [PubMed: 15080678]
13. Papa AJ. *J Org Chem*. 1966; 31:1426.
14. Brochu MP, Brown SP, MacMillan DWC. *J Am Chem Soc*. 2004; 126:4108. [PubMed: 15053591]
15. Kobayashi S, Tsubogo T, Saito S, Yamashita Y. *Org Lett*. 2008; 10:807. [PubMed: 18225909]
16. Tsubogo T, Saito S, Seki K, Yamashita Y, Kobayashi S. *J Am Chem Soc*. 2008; 130:13321. [PubMed: 18783222]
17. (a) Alper PB, Hung SC, Wong CH. *Tetrahedron Lett*. 1996; 37:6029.(b) Nyffeler PT, Liang CH, Koeller KM, Wong CH. *J Am Chem Soc*. 2002; 124:10773. [PubMed: 12207533]
18. As observed previously in literature in ref. 17b, the metal-mediated diazotransfer reaction is prone to irreproducibility in yields.
19. (a) Omura K, Swern D. *Tetrahedron*. 1978; 34:1651.(b) Mancuso AJ, Huang SL, Swern D. *J Org Chem*. 1978; 43:2480.
20. (a) Charette AB, Boezio AA, Janes MK. *Org Lett*. 2000; 2:3777. [PubMed: 11101417] (b) Grieder A, Thomas AW. *Synth*. 2003; 2003:1707.(c) Gil C, Bräse S. *Chem Eur J*. 2005; 11:2680. [PubMed: 15736275] (d) Baxendale IR, Deeley J, Griffiths-Jones CM, Ley SV, Saaby S, Tranmer GK. *Chem Commun*. 2006:2566.(e) Mahdavi H, Amani J. *Tetrahedron Lett*. 2009; 50:5923.(f) Smith CJ, Smith CD, Nikbin N, Ley SV, Baxendale IR. *Org Biomol Chem*. 2011; 9:1927. [PubMed: 21283873]

Figure 1.

The stereochemistry generated in the production of imine **9** was preserved through multiple transformations (Scheme 3) as revealed by X-ray structure crystallographic analysis of acid **10**.

Scheme 1.
Retrosynthesis of Pyrrolysine

Scheme 2.

Attempts to synthesize pyrrollysine precursors via asymmetric methylcarbanion 1,4-addition and α -substitution

Scheme 3.
Pyrroline precursors via Michael Addition

Scheme 4.
Generating the Staudinger/aza-Wittig substrate

Scheme 5.
Tandem Staudinger/aza-Wittig and protecting group removal.