PstI and RsaI RFLPs in complete linkage disequilibrium at the CYP2E gene

J.Watanabe*, S.-I.Hayashi, K.Nakachi¹, K.Imai¹, Y.Suda², T.Sekine² and K.Kawajiri

Departments of Biochemistry and ¹Epidemiology, Saitama Cancer Center Research Institute, and ²Division of Abdominal Surgery, Saitama Cancer Center Hospital, Inamachi, Kitaadachi-gun, Saitama 362, Japan

Source/Description: A 1.5 kb ApaI-ApaI fragment, containing 5' upstream sequences of the human CYP2E gene (1).

Polymorphism: PstI detects a two allele polymorphism with bands at 6.1 kb (C1) or 3.9 and 2.2 kb (C2). RsaI detects a two allele polymorphism with bands at 1.4 and 0.5 kb (C1) or 1.9 kb (C2). These two polymorphisms are in complete linkage disequilibrium with each other.

Frequency: 202 unrelated healthy Japanese were screened. C1: 0.807, C2: 0.193.

Not Polymorphic For: BamHI, HaeIII, KpnI, MspI, PvuII, Sau3A and XbaI.

Chromosomal Localisation: Human chromosome 10 (1).

Mendelian Inheritance: Codominant inheritance observed in 3 families.

Probe Availability: Contact J.Watanabe.

Other Comments: No linkage disequilibrium with TaqI RFLP (2) was observed. We have found no correlation between the PstI polymorphism and cancers of the stomach and large bowel, tested in 102 and 86 patients, respectively.

References: 1) Umeno, M. *et al.* (1988) *Biochemistry* **27**, 9006–9013. 2) McBride, O.W. *et al.* (1987) *Nucl. Acids Res.* **15**, 10071.

An SspI RFLP at the D13S25 locus identified by the anonymous single copy probe H2 – 42

Anne M.Bowcock* and Joan M.Herbert

Department of Genetics, Stanford University, Stanford, CA 94305, USA

Source/Description: H2–42 contains a 2.2 kb fragment subcloned into the HindIII site of pBR322. The original HindIII fragment was subcloned from a Charon 21A bacteriophage derived from a chromosome 13 specific library (1).

Polymorphism: A three-allele polymorphism is detected with SspI (AAT/ATT). A1 = 1.75 kb; A2 = 1.60 kb; A3 = 1.8 kb. A constant fragment of 1.05 kb is also present.

Frequency: Estimated from 89 unrelated Caucasians.

A1 = 0.41

A2 = 0.58

A3 = 0.01

Not Polymorphic For: ApaLI, AluI, BamHI, BanII, BclI, BglI, BglII, BstNI, DdeI, DraI, EcoRI, EcoRV, HaeIII, HincII, HindIII, HphI, KpnI, MboI, MspI, PstI, PvuII, RsaI, Sau96I, ScaI, ScrFI, SstI, StuI, StyI, TaqI, XbaI, XmnI when a panel of at least 6 unrelated Caucasians was screened for each enzyme.

Chromosomal Localization: Chromosome 13q14 (1, 2).

Mendelian Inheritance: Co-dominant segregation was demonstrated in 24 informative families (293 individuals).

Probe Availability: From the ATCC.

Other Comments: The frequency of the A3 allele was estimated from one individual with an A2/A3 genotype.

Acknowledgement: This work was supported by NIH grant GM28428.

References: 1) Lalande *et al.* (1984) *Cancer Genet. Cytogenet.* **13**, 283–285. 2) Lalande *et al.* (1986) *Cancer Genet. Cytogenet.* **23**, 151–157.

* To whom correspondence should be addressed

*Present address: Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, TX 75235–9063, USA