Published in final edited form as:

Science. 2011 October 14; 334(6053): 242–245. doi:10.1126/science.1210460.

NMR Detection of Structures in the HIV-1 5'-Leader RNA that Regulate Genome Packaging

Kun Lu^{1,+,#}, Xiao Heng^{1,+}, Lianko Garyu¹, Sarah Monti¹, Eric L. Garcia², Siarhei Kharytonchyk², Bilguujin Dorjsuren¹, Gowry Kulandaivel¹, Simonne Jones¹, Atheeth Hiremath¹, Sai Sachin Divakaruni¹, Courtney LaCotti¹, Shawn Barton¹, Daniel Tummillo¹, Azra Hoscic¹, Kedy Edme¹, Sara Albrecht¹, Alice Telesnitsky^{2,*}, and Michael F. Summers^{1,*}

¹Howard Hughes Medical Institute and Department of Chemistry and Biochemistry, University of Maryland Baltimore County, 1000 Hilltop Circle, Baltimore, Maryland 21250

²Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor MI 48109-0620

Abstract

The 5'-leader of the HIV-1 genome regulates multiple functions during viral replication by mechanisms that have yet to be established. We developed an NMR approach that enabled direct detection of structural elements within the intact leader (712 nucleotide dimer) that are critical for genome packaging. Residues spanning the *gag* start codon (AUG) form a hairpin in the monomeric leader and base pair with residues of the Unique-5' region (U5) in the dimer. U5:AUG formation promotes dimerization by displacing and exposing a dimer-promoting hairpin, and enhances binding by the nucleocapsid protein (NC), the cognate domain of the viral Gag polyprotein that directs packaging. Our findings support a packaging mechanism in which translation, dimerization, NC binding, and packaging are regulated by a common RNA structural switch.

The 5'-leader is the most conserved region of the HIV-1 genome and is responsible for regulating multiple activities during viral replication, including RNA encapsidation during virus assembly (1, 2). Like all retroviruses, HIV-1 packages two copies of its genome, enabling strand-transfer mediated recombination during reverse transcription and promoting genetic evolution under environmental and chemotherapeutic pressure (3). Dimeric genomes are trafficked from the cytoplasm to plasma membrane assembly sites by a small number of viral Gag proteins, where additional Gag proteins assemble and budding occurs (4–8). Dimerization and packaging are mediated by interactions between the nucleocapsid (NC) domains of the viral Gag polyproteins and RNA elements within the 5'-leader of the genome, and there is evidence that these activities and translational control are mechanistically coupled (7, 9–13).

Understanding of the mechanisms that regulate 5'-leader activities is limited, due in part to incomplete knowledge of the leader structure (14). Recombinant leader-containing RNAs have been probed by nucleotide accessibility mapping, mutagenesis, and biochemical approaches, and although there is general consensus that transcriptional activation, primer binding, dimerization, and splicing activities are promoted by discrete hairpin structures (10, 15–23) (TAR, PBS, DIS and SD hairpins, respectively; Fig. 1), there is less agreement

*Correspondence: ateles@umich.edu; summers@hhmi.umbc.edu.

#Current address: The RNA Institute, State University of New York at Albany, 1400 Washington Avenue, Albany, NY 12222

+These authors contributed significantly to this work.

regarding the structures that regulate packaging (14). In particular, residues overlapping the *gag* start codon (G328-A356, AUG) that are critical for genome packaging (24) and RNA dimer stability (25) have been proposed to form a hairpin (Fig. 1B), to base-pair with residues of the upstream Unique-5' element (U5, Fig. 1C) (10, 20), or to adopt other conformations (14). *In vivo* nucleotide reactivity mapping has supported multiple AUG models, without consensus (14, 21–23). NMR is potentially well suited to probe RNA structure, but signal degeneracy and relaxation problems have thus far limited applications to relatively small oligonucleotides (typically fewer than 50 residues) (26, 27). Here we describe an NMR approach that enabled direct detection of structures formed by AUG and other packaging elements within the intact, 230 kDalton dimeric 5'-leader, as well as the identification of conformational changes that regulate dimerization, NC binding and packaging.

A 356 nucleotide HIV-1_{NL4-3} 5'-leader RNA that includes the entire 5'-Untranslated Region (5'-UTR) and the first 21 residues of *gag* (5'-L) was prepared by enzymatic ligation of non-labeled 5'-RNA (residues 1–327) and ¹³C-enriched AUG fragments (residues 328–356) (Fig. 1) (28). The 5'-leader exists predominantly as a monomer at low ionic strength (Fig. 1D), and under these conditions the AUG residues of 5'-L gave rise to ¹H-¹³C correlated HMQC (28) NMR spectra similar to those observed for the isolated AUG fragment, which is known to form a hairpin (29) (Fig. 1E,F). NMR line widths of AUG in the intact 5'-leader and the isolated AUG RNA were also similar, indicating that the hairpin is structurally mobile and does not form A-minor-like contacts (30).

Significant changes in the ¹H-¹³C HMQC spectra of AUG-labeled 5'-L were observed upon incubation at physiological ionic strength (PI buffer: 10 mM Tris-HCl, 140 mM KCl, 10 mM NaCl, 1 mM MgCl₂, pH 7.0), which correspond to a reversible equilibrium shift from a predominantly monomeric species to a mixture of monomeric and dimeric species (Fig. 1G,H; dimer dissociation constant $K_d = 0.9 \pm 0.1 \mu\text{M}$). Signals of the AUG hairpin in 5'-L exhibited uniformly reduced intensities as a function of incubation time, and new signals were observed for the 3'-residues of AUG (A345-A356) (Fig. 1H). Similar NMR spectral changes were observed for an isolated AUG oligo-RNA upon titration with an oligo-U5 fragment (Fig. 1I). The NMR chemical shifts and narrow line widths observed for A345-A356 indicate that these residues are unstructured and mobile in the dimeric form of the leader ([5'-L]₂). No ¹H-¹³C HMQC signals were detected for G328-G344 of [5'-L]₂ due to severe line broadening (discussed below). These findings are consistent with a phylogenetically-derived structural model (10, 20), in which the 5'-residues of AUG base pair with U5 and the 3'-residues are disordered (Fig. 1C).

To directly probe for U5:AUG base pairing, we developed an NMR approach that involves replacement of a short stretch of adjacent base pairs by A-U base pairs (lr-AID; long-range probing by Adenosine Interaction Detection). The substituting element (ideally [U_iU_jA_k]: [U_iA_mA_n], but other sequences can suffice) (Fig. 2A) affords an upfield-shifted A_m-C2-¹H NMR signal (~6.5 ppm), enabling direct detection of cross-strand A_k-H2 and -H1' ¹H-¹H nuclear Overhauser effects (NOEs) without the need for heteronuclear spectral editing. The HIV-1_{NL4-3} leader naturally contains one [UUA:UAA] element in the TAR hairpin ([U12-A14]:[U45-A47]), and to preclude signal overlap, A46 was mutated to G (a naturally occurring substitution in 7% of the reported HIV-1 TAR sequences (31)) (Fig. 2B–D). NOESY spectra (28) obtained for the lr-AID modified dimeric 5'-leader ([5'-L^{lr-AID-U5:AUG}]₂, C110-G112 and G338-G339 substituted by UUA and AA, respectively; Fig. 2A) exhibited well-resolved A338-H2 signals with frequencies similar to those observed for an isolated lr-AID modified U5:AUG oligoribonucleotide (Fig. 2E,F). Other outlier signals in the NOESY spectrum were unaffected by the lr-AID substitutions. All of the expected A338-H2 inter-adenosine NOEs were observed, including cross-strand NOEs

with A112-H1' and -H2. The A338-H2 T₁ and T₂ ¹H NMR relaxation rates (8.7 sec and 9.4 msec, respectively) are consistent with restricted rotational motion and indicate that U5:AUG resides within a globular region of the 5'-L structure.

The relationship between U5:AUG formation and dimerization was investigated by site-directed mutagenesis. Mutations in AUG designed to stabilize the hairpin and disrupt base pairing with U5 (5'-L^{AUG-HP}) favored the monomer, whereas mutations that promote U5:AUG base pairing and destabilize the hairpin ([5'-L^{U5:AUG}]₂) favored the dimer (Fig. S1). Deletion of AUG (5'-L^{ΔAUG}) also favored the monomer (Fig. 3A), whereas incubation of 5'-L^{ΔAUG} with a 17-nucleotide AUG oligoribonucleotide (G328-G344; AUG-17) promoted dimerization (Fig. 3B). These findings indicate that dimerization is suppressed when AUG exists in a hairpin conformation (AUG^{HP}), that U5:AUG formation promotes dimerization, and that dimerization is induced by intramolecular U5:AUG base pairing rather than intermolecular tethering.

NMR chemical shifts and NOEs associated with A268-H2 in 5'-L, 5'-L^{ΔAUG}, and an isolated DIS oligo-RNA were similar, indicating that the attenuated dimerization activity of the AUG^{HP} form of 5'-L is not due to re-folding of the DIS (32). Sequence complementarity between U5 and the GC-rich loop of the DIS suggested that dimerization may instead be inhibited by U5:DIS base pairing. Consistent with this hypothesis, mutations in 5'-L designed to enhance U5:DIS base pairing promoted monomer formation (Fig. 3C,D), whereas mutations to disrupt U5:DIS base pairing induced dimerization of 5'-L^{ΔAUG} (Fig. S2). In addition, NMR data obtained for an Ir-AID substituted 5'-L^{ΔAUG} RNA with 5'-L^{ΔAUG}-like dimerization and NC binding properties (Fig. S4) exhibited cross-strand NOEs and NMR chemical shifts consistent with the predicted U5:DIS interface (Fig. 3E-G).

To determine if U5:AUG formation influences NC binding, isothermal titration calorimetry (ITC) experiments were performed with 5'-L and mutant 5'-L^{AUG-HP} and 5'-L^{U5:AUG} RNAs. Under conditions of the ITC experiments, 5'-L^{AUG-HP} and 5'-L^{U5:AUG} exist predominantly as monomers and dimers, respectively, and native 5'-L exists as a 70:30 monomer:dimer equilibrium mixture (Fig. 4A). All three RNAs gave rise to two-component NC binding isotherms that included an initial exothermic event associated with high-affinity NC binding, and a subsequent endothermic event attributed to NC-induced RNA unfolding at high NC:RNA ratios (33) (Fig. 4B and Fig. S3). Whereas 5'-L^{AUG-HP} binds 7 ± 1 NC molecules with high affinity (K_d = 87 ± 3 nM), [5'-L^{U5:AUG}]₂ binds 32 ± 2 NC molecules with high affinity (K_d = 71 ± 3 nM) (Fig. 4B). The ITC profile observed for native 5'-L was similar to that obtained for a 70:30 mixture of 5'-L^{AUG-HP} and 5'-L^{U5:AUG} (Fig. 4A,B), validating the use of 5'-L^{AUG-HP} and 5'-L^{U5:AUG} as models for the native monomer and dimer, respectively. Importantly, the NC binding and dimerization properties of 5'-L^{Ir-AID-U5:AUG} and 5'-L^{Ir-AID-U5:DIS} were similar to those of the 5'-L^{U5:AUG} and 5'-L^{AUG-HP} controls, respectively (Fig. S4), further indicating that the Ir-AID substitutions did not alter the structure of the RNA. Substitution of the dimer-promoting GC-rich loop of the DIS hairpin by a GAGA tetraloop prevented dimerization of 5'-L^{U5:AUG} but did not affect its NC binding properties (Fig. S5), indicating that enhanced NC binding by the U5:AUG form of the 5'-leader is due to intramolecular conformational changes associated with U5:AUG base pairing and not to dimerization *per se*.

The above findings suggested that *in vivo* RNA packaging should be dependent on U5:AUG formation, and we therefore measured packaging efficiencies of vector RNAs containing 5'-L^{AUG-HP} and 5'-L^{U5:AUG} mutations in competition experiments. An HIV-1_{NL4-3} helper construct that expressed the native 5'-leader and all viral proteins except Env was co-expressed with test vector RNAs with 5'-L^{AUG-HP} or 5'-L^{U5:AUG} mutations (Fig. 4C). Consistent with the structural and NC binding studies, the 5'-L^{U5:AUG} RNAs were packaged

nearly as avidly as the native construct (Fig. 4C; lane 5) whereas the 5'-L^{AUG-HP} RNAs exhibited severe packaging defects (Fig. 4C; lane 6).

Previously observed packaging defects associated with mutations in AUG (24, 34) can be attributed to defects in U5:AUG dependent exposure of NC binding sites rather than inhibition of a *cis*-packaging mechanism (34). The inability of helper RNAs to rescue packaging of RNAs with AUG mutations (24) is likely due to a defect in U5:AUG dependent exposure of the DIS since rescue requires DIS-mediated heterodimer formation. The relationship between U5:AUG formation and dimerization also explains why mutations in AUG can lead to the production of virions containing genomes that are sensitive to dissociation by mild denaturants (25). Hybrid 5'-leader structures containing both U5:AUG and AUG^{HP} features, predicted from *in vivo* ribose reactivity measurements (22), were not observed by NMR but can be explained by the presence of the AUG^{HP}/U5:AUG equilibrium. Elevated U5 nucleotide reactivities observed by traditional chemical probing, which are incompatible with an exclusive U5:AUG structure (21), may also be explained by the observed AUG^{HP}/U5:AUG equilibrium. The finding that the *gag* start codon is exposed in a mobile hairpin in the monomeric leader and sequestered in the dimer is consistent with observations that dimerization attenuates both the chemical reactivity of the *gag* start codon and the *in vitro* translational activity of the genome (16).

In summary, we have developed an approach that extends the size of RNAs that can be structurally probed by NMR to ~230 kDaltons. The Ir-AID method avoids NMR relaxation problems associated with heteronuclear editing, probes interactions over a narrow distance range (~5 Å), provides both chemical shift and distance information for structural analysis, and is readily implemented using standard 2D NMR experiments and conservative base pair mutagenesis. Our findings support a translation/packaging RNA structural switch mechanism, in which the dimer-promoting GC-rich loop of the DIS hairpin is sequestered by base pairing with U5 in the AUG^{HP} form of the 5'-leader, and is displaced by AUG upon U5:AUG formation (Fig. 4D). Conformational changes associated with U5:AUG base pairing simultaneously sequester the *gag* start codon and expose the DIS and high affinity NC binding sites, thereby attenuating translation and promoting the packaging of a dimeric genome.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by a grant from the National Institute of General Medical Sciences (NIGMS; R01 GM42561). L.G. was supported by an NIGMS grant for maximizing doctoral diversity (R25 MBRS-IMSD GM55036). B.D., K.E., S.J., G.K. and S.B. were supported by an NIGMS grant for enhancing minority access to research careers (MARC U*STAR 2T34 GM008663). B.D., G.K. and S.B. were supported by an HHMI undergraduate education grant. We thank Steven R. King (Michigan) and HHMI staff (UMBC) for technical assistance.

References

1. Coffin, JM.; Hughes, SH.; Varmus, HE. Retroviruses. Plainview, N.Y.: Cold Spring Harbor Laboratory Press; 1997.
2. Lever AM. Adv Pharmacol. 2007; 55
3. Onafuwa-Nuga A, Telesnitsky A. Microbiol Mol Biol Rev. 2009; 73
4. Poole E, Strappe P, Mok HP, Hicks R, Lever AM. Traffic. 2005; 6
5. Moore MD, et al. PLoS Pathog. 2009; 5

6. Kutluay SB, Bieniasz PD. PLoS pathogens. 2010; 6
7. Paillart J-C, Shehu-Xhilaga M, Marquet R, Mak J. Nature Revs. Microbiol. 2004; 2
8. Jouvenet N, Simon SM, Bieniasz PD. Proc. Natl. Acad. Sci. USA. 2009; 106
9. Miele G, Moulard A, Harrison GP, Cohen E, Lever AM. J. Virol. 1996; 70
10. Abbink TEM, Berkhout B. J. Biol. Chem. 2003; 278
11. Russell RS, Liang C, Wainberg MA. Retrovirology. 2004; 1
12. Groatorex J. Retrovirology. 2004; 1
13. D'Souza V, Summers MF. Nature Revs. Microbiol. 2005; 3
14. Lu K, Heng X, Summers MF. J. Mol. Biol. 2011; 410
15. Harrison GP, Lever AML. J. Virol. 1992; 66
16. Baudin F, et al. J. Mol. Biol. 1993; 229
17. Clever J, Sassetti C, Parslow TG. J. Virol. 1995; 69
18. McBride MS, Panganiban AT. J. Virol. 1996; 70
19. Clever JL, Miranda J, D, Parslow TG. J. Virol. 2002; 76
20. Damgaard CK, Andersen ES, Knudsen B, Gorodkin J, Kjems J. J. Mol. Biol. 2004; 336
21. Paillart JC, et al. J. Biol. Chem. 2004; 279
22. Wilkinson KA, et al. PLoS Biology. 2008; 6
23. Watts JM, et al. Nature. 2009; 460
24. Nikolaitchik O, Rhodes TD, Ott D, Hu W-S. J. Virol. 2006; 80
25. Song R, Kafaie J, Laughrea M. Biochemistry. 2008; 47
26. Allain FH-T, Varani G. J. Mol. Biol. 1997; 267
27. Tolbert BS, et al. J. Biomol. NMR. 2010; 47
28. Materials and methods are available as supporting material.
29. Kerwood DJ, Cavaluzzi MJ, Borer PN. Biochemistry. 2001; 40
30. Yu ET, Hawkins A, Eaton J, Fabris D. Proc. Natl. Acad. Sci. USA. 2008; 105
31. The HIV Sequence Database. 2011. <<http://www.hiv.lanl.gov/content/sequence/NEW-ALIGN/align.html>>
32. Huthoff H, Berkhout B. RNA. 2001; 7
33. Summers MF, et al. Protein Sci. 1992; 1
34. Poon DT, Chertova EN, Ott DE. Virology. 2002; 293

Fig. 1. Structure of the HIV-1 5'-leader

(A–C), Secondary structure predictions showing AUG (green) in hairpin (B) and U5:AUG (C) conformations. (D) 5'-L forms a monomer at low ionic strength (LI buffer = 10 mM Tris-HCl, 10 mM NaCl, pH 7.0) ([RNA] = 60 μM). (E) 2D ^1H - ^{13}C HMQC NMR spectrum obtained for 5'-L containing ^{13}C -labeled AUG (5'-L^{AUG*}; LI buffer). (F) Overlay of spectra obtained for 5'-L^{AUG*} (gray) and oligo-AUG* hairpin (green). (G) 5'-L exists as a reversible monomer-dimer equilibrium in PI buffer ([5'-L] = 30 μM in left panel). The equilibrium shifts to the monomer upon subsequent dilution (100-fold) in PI buffer. (H) ^1H - ^{13}C HMQC NMR data obtained for 5'-L^{AUG*} in LI buffer (black) and immediately upon dissolution in PI buffer (red). Spectral changes correlate with an equilibrium shift toward the dimer. Residues 349, 351, 352 and 355 are unstructured and detectable, and residues 337 and 341 are rotationally restricted and undetectable, in the [5'-L]₂ dimer. (I) Similar spectral changes were observed upon titration of an isolated AUG* hairpin (black) with an unlabeled U5 oligonucleotide, except that all U5:AUG signals were detectable (red).

Fig. 2. AUG base pairs with U5 in the dimer

(A) Ir-AID mutations (red) designed to probe for predicted U5:AUG base pairing. Black arrows denote ^1H - ^1H NOEs. (B) Portions of 1D ^1H NMR spectra showing (top): the TAR A46-H2 signal of native $[5'\text{-L}]_2$; (middle): A46G substitution ($[5'\text{-L}^{\text{A46G}}]_2$) eliminates the TAR A46-H2 signal; (bottom): A338-H2 signal observed for Ir-AID substituted $[5'\text{-L}^{\text{A46G}}]_2$. (C,D) Similarities in NOE spectra observed for A46-H2 of an isolated TAR oligonucleotide (C) and native $[5'\text{-L}]_2$ (D). (E,F) Similar A338-H2 NOEs observed for an isolated Ir-AID U5:AUG oligonucleotide (E) and Ir-AID substituted $[5'\text{-L}^{\text{A46G}}]_2$ (F).

Fig. 3. U5:AUG formation promotes dimerization

(A) 5'-L Δ AUG (1 μ M) forms a stable monomer. (B) Addition of AUG-17 (10-fold molar excess) promotes dimerization of 5'-L Δ AUG. (C,D) Substitutions that enhance U5:DIS interactions favor the monomer. (E) Proposed U5:DIS base pairing and lr-AID sequences used for structural probing. (F) The A259-H2 signal is only observed when U5 and DIS both contain lr-AID substitutions. Top: 5'-L Δ AUG, A46G. Middle: 5'-L Δ AUG, A46G containing the lr-AID substitution in the DIS loop only. Bottom: 5'-L Δ AUG, A46G containing lr-AID substitution in both DIS and U5. (G) 2D NOE data for A259-H2 in lr-AID substituted 5'-L Δ AUG, A46G, assigned from the corresponding U5:DIS oligonucleotide control spectrum.

Fig. 4. U5:AUG formation promotes NC binding and packaging

(A) 5'-L^{AUG-HP} and 5'-L^{U5:AUG} form predominantly monomers and dimers, respectively, and native 5'-L adopts a 70:30 monomer:dimer equilibrium (PI buffer; [RNA] = 0.8 µM). MIX = 70:30 mixture of 5'-L^{AUG-HP} and 5'-L^{U5:AUG}. (B) ITC NC titration data for samples shown in (A). (C) Packaging of native HIV-1_{NL4-3} 5'-L, 5'-L^{U5:AUG} and 5'-L^{AUG-HP} RNAs under competition conditions (assayed by RNase protection; see Supplementary Methods). Lanes 2–8: 2: undigested probe; 3: RNA size standards; 4: native HIV-1_{NL4-3} helper versus native HIV-1_{NL4-3} test vector RNAs; 5: native HIV-1_{NL4-3} helper versus 5'-L^{U5:AUG} test vector; 6: native HIV-1_{NL4-3} helper versus 5'-L^{AUG-HP} test vector; 7: HIV-1_{NL4-3} helper

expressed without test RNA; 8: mock transfected-cells. Bands corresponding to native HIV-1_{NL4-3} helper RNA (helper) and co-packaged test RNAs (test) are labeled. **(D)** Nucleotide displacement mechanism regulates dimeric genome packaging.