Published in final edited form as:

Arch Pediatr Adolesc Med. 2011 December ; 165(12): 1133–1134. doi:10.1001/archpediatrics.2011.1046.

Conceptual and Methodological Issues in Neuroimaging Studies of the Effects of Child Maltreatment: Commentary on Edmiston et al

Philip A. Fisher, PhD and
University of Oregon and Oregon Social Learning Center

Jennifer H. Pfeifer, PhD
University of Oregon

The extensive scientific literature on child maltreatment (CM) has provided strong evidence of the globally negative effects of abuse, neglect, and emotional maltreatment on healthy development and of the remarkable resilience of individuals who manage to prosper even in the face of these adverse experiences.^{1,2} Yet, after many decades of CM research, the underlying mechanisms by which risk and resilience are conferred are just beginning to be understood. In recent years, there has been an acceleration of progress in this area, owing largely to advances in developmental cognitive neuroscience methodologies that are allowing researchers an unprecedented opportunity to understand the manner by which these experiences get “under the skin”³ to affect the development of neurobiological systems.^{4–6} These findings (from animal and human studies) document structural and functional changes in the developing brain associated with CM^{7,8} and yield strong support for the role of CM in subsequent psychopathology, in childhood and across the lifespan.⁹ Edmiston et al. followed this recent trend of neuroscientific investigations of CM but broke new ground in four important ways.

First, by examining adolescents, Edmiston et al. revealed *linkages between the neurobiological effects of CM previously observed in children and adults*. Specifically, Edmiston et al. showed that gray matter (GM) volume is decreased across several cortical, striatal, and limbic regions in adolescents reporting prior CM. The results cited in Edmiston et al. demonstrate GM volume decreases in both adults and children. However, whereas volumetric hippocampal reductions are the most frequent finding in previous research with adults with CM, this pattern is absent in children with CM, raising the question of whether it emerges in adolescence. Edmiston et al. suggested that such volumetric hippocampal reductions are present in some adolescents.

Second, whereas prior neuroscience researchers have treated CM as a unitary construct (without distinguishing subtypes or severity of CM) or have focused on a single type of CM (e.g., physical abuse), Edmiston et al. examined differences on measures of brain morphometry *according to CM subtypes*. For example, the above noted decreased hippocampal volume in the CM sample was primarily observed in those reporting emotional neglect. Similarly, the authors noted other regional effects that were common across CM subtypes (specifically, GM reductions in rostral prefrontal cortex [PFC]) and provided logical (albeit speculative) hypotheses about how these differences might explain parallel differences in previously reported behavioral sequelae of specific CM subtypes.

Third, Edmiston et al. examined *gender dimorphic CM effects* on brain morphometry. Specifically, the regions most impacted in males with prior CM appear to be more strongly associated with impulse control, including the caudate and rostral PFC. In contrast, the females showed alterations in multiple areas associated with emotion reactivity and regulation, including the amygdala, orbitofrontal cortex, subgenual cingulate cortex, insula, and hippocampus. Similar to the results for CM subtypes, the gender analyses provide evidence of concordance between previously observed behavioral phenomena (specifically, differential vulnerability to psychopathology) and underlying neurobiological effects.

Fourth, and perhaps the most noteworthy characteristic of the study, *the sample consisted of individuals who did not meet criteria for a mental health disorder*. The morphometric differences that were noted and were associated with particular CM subtypes existed among individuals who appeared behaviorally unscathed by their experiences of prior adversity. As Edmiston et al. noted, these results might be evidence of future vulnerabilities and/or evidence of adaptations that have allowed compensatory processes to facilitate healthy development in the face of adversity. Additional work in this area is particularly important to meeting the needs of this population because understanding future risk in the absence of behavioral difficulties would greatly assist in the development of screening and prevention programs. Furthermore, knowledge about compensatory processes that help to support resiliency could be leveraged in the development of treatment programs to mitigate CM effects.

Next Steps

Given the importance of Edmiston et al.'s study, we believe that an expansion of research on the developmental neurobiology of CM subtypes is warranted. One particularly relevant new mindset would consider the regional developmental trajectories of GM *and* white matter (WM) in a movement towards thinking about these effects at the level of networks rather than isolated regions. Anatomical integration between remote brain regions is accomplished by WM tracts, and measures of WM integrity such as diffusion tensor imaging have illustrated that WM development steadily increases into adulthood^{10,11} In particular, WM fibers connecting the PFC to subcortical limbic and striatal structures (including those found to be affected by CM in Edmiston et al.'s study) continue to mature into young adulthood.¹² Another relevant element of brain development is that of functional connectivity, which is complementary to, but not synonymous with, anatomical connectivity. Functional connectivity can be estimated in a task-independent fashion (i.e., at rest),¹³ and such methods have led to the identification of multiple networks—some of which are developed by puberty, although network metrics continue to evolve with further development.¹⁴

Therefore, to fully understand the impact of the GM reductions observed in individuals with CM across various cortical, striatal, and limbic regions (including those reported in Edmiston et al.'s study), it would be useful to consider the following:

- Are the precise locations of GM reductions in adults with CM consistent across studies? Meta-analyses conducted using tools such as GingerALE could provide insight beyond summary reporting of labeled regions.¹⁵
- Given the answer to the preceding question, does the trajectory of GM development in each of the regions impacted by CM predict the age during which GM reductions attributable to CM are first observed? The regionally specific patterns of maturation in the cortical and subcortical regions and the WM that connects them^{12,16,17} have been increasingly documented. Furthermore, changes in WM are known to be associated with changes in GM.^{18,19}

- Can we use typically developing patterns of anatomical or functional connectivity between cortical, limbic, and striatal regions to identify particular networks or additional regions that are placed at risk by CM in a “downstream” fashion, according to the earliest-emerging affected regions?
- Finally, can we use information about these networks’ functional contributions to better understand the behavioral consequences of these atypical patterns of neurodevelopment? Because any given brain region is responsible for multiple functions, it is important to be cautious about making strong claims that depend on reverse inference.^{20–21} Network-wide analyses and searches for selective associations should be more robust in regard to this issue.^{22–24}

Research in these above areas has great potential to address the needs for more effective prevention and treatments programs for individuals with specific CM subtypes.

Acknowledgments

Support for this article was provided by the following grants: MH078105, NIMH, U.S. PHS; DA023920 and DA021424, NIDA, U.S. PHS; R324A080026, USDE; HD045894, NICHD, U.S. PHS.

References

1. Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet*. 2009; 373(9657):68–81. [PubMed: 19056114]
2. Collinshaw S, Pickles A, Messer J, Rutter M, Shearer C, Maughan B. Resilience to adult psychopathology following childhood maltreatment: Evidence from a community sample. *Child Abuse Negl*. 2007; 31:211–229. [PubMed: 17399786]
3. Hyman SE. How adversity gets under the skin. *Nat Neurosci*. 2009; 12:241–243. [PubMed: 19238182]
4. Heim C, Shugart M, Craighead WE, Nemeroff CB. Neurobiological and psychiatric consequences of child abuse and neglect. *Dev Psychobiol*. 2010; 52:671–690. [PubMed: 20882586]
5. McCrory E, De Brito SA, Viding E. Research review: the neurobiology and genetics of maltreatment and adversity. *J Child Psychol Psychiatry*. 2010; 51:1079–1095. [PubMed: 20546078]
6. McGowan PO, Sasaki A, D’Alessio, et al. Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nat Neurosci*. 2009; 12:342–348. [PubMed: 19234457]
7. Watts-English T, Fortson BL, Gibler N, Hooper SR, De Bellis MD. The psychobiology of maltreatment in childhood. *J Soc Issues*. 2006; 62:717–736.
8. Sánchez MM, Ladd CO, Plotsky PM. Early adverse experience as a developmental risk factor for later psychopathology: evidence from rodent and primate models. *Dev Psychopathol*. 2001; 13:419–449. [PubMed: 11523842]
9. Woon FL, Hedges DW. Hippocampal and amygdala volumes in children and adults with childhood maltreatment-related posttraumatic stress disorder: a meta-analysis. *Hippocampus*. 2008; 18:729–736. [PubMed: 18446827]
10. Lenroot RK, Giedd JN. Brain development in children and adolescents: insights from anatomical magnetic resonance imaging. *Neurosci Biobehav Rev*. 2006; 30:718–729. [PubMed: 16887188]
11. Paus T. Mapping brain maturation and cognitive development during adolescence. *Trends Cogn Sci*. 2004; 9(2):60–68. [PubMed: 15668098]
12. Tamnes CK, Østby Y, Fjell AM, Westlye LT, Due-Tønnessen P, Walhovd KB. Brain maturation in adolescence and young adulthood: regional age-related changes in cortical thickness and white matter volume and microstructure. *Cereb Cortex*. 2010; 20:534–548. [PubMed: 19520764]
13. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Nat Acad Sci*. 2001; 98:676–682. [PubMed: 11209064]

14. Fair DA, Cohen AL, Dosenbach NUF, Church JA, Miezin FM, Barch DM, et al. The maturing architecture of the brain's default network. *Proc Nat Acad Sci*. 2008; 105:4028–4032. [PubMed: 18322013]
15. Laird AR, McMillan KM, Lancaster JL, Kochunov P, Turkeltaub PE, Pardo JV, et al. A comparison of label-based review and activation likelihood estimation in the Stroop task. *Hum Brain Mapping*. 2005; 25:6–21.
16. Østby Y, Tamnes CK, Fjell AM, Westlye LT, Due-Tønnessen P, Walhovd KB. Heterogeneity in subcortical brain development: a structural magnetic resonance imaging study of brain maturation from 8 to 30 years. *J Neurosci*. 2009; 29:11772–11782. [PubMed: 19776264]
17. Shaw P, Kabani NJ, Lerch JP, Eckstrand K, Lenroot R, Gogtay N, et al. Neurodevelopmental trajectories of the human cerebral cortex. *J Neurosci*. 2008; 28:3586–3594. [PubMed: 18385317]
18. Giorgio A, Watkins KE, Douaud G, James AC, James S, De Stefano N, et al. Changes in white matter microstructure during adolescence. *Neuro Image*. 2008; 39:52–61. [PubMed: 17919933]
19. Lebel C, Walker L, Leemans A, Phillips L, Beaulieu C. Microstructural maturation of the human brain from childhood to adulthood. *Neuro Image*. 2008; 40:1044–1055. [PubMed: 18295509]
20. Poldrack RA. Can cognitive processes be inferred from neuroimaging data? *Trends Cogn Sci*. 2006; 10:59–63. [PubMed: 16406760]
21. Christoff K, Owen AM. Improving reverse neuroimaging inference: cognitive domain versus cognitive complexity. *Trends Cogn Sci*. 2006; 10:352–353. [PubMed: 16843038]
22. McIntosh AR. Towards a network theory of cognition. *Neural Netw*. 2000; 13:861–870. [PubMed: 11156197]
23. Poldrack RA, Halchenko YO, Hanson SJ. Decoding the large-scale structure of brain function by classifying mental states across individuals. *Psychol Sci*. 2009; 20:1364–1372. [PubMed: 19883493]
24. Poldrack RA. Mapping mental function to brain structure: how can cognitive neuroimaging succeed? *Perspect Psychol Sci*. 2010; 5:753–761.