Detection of *Sst*I restriction site polymorphism in human *APOC3* by the polymerase chain reaction

J.E.Hixson*, D.T.Vernier and P.K.Powers

Department of Genetics, Southwest Foundation for Biomedical Research, PO Box 28147, San Antonio, TX 78228-0147, USA

Source and Description: Karathanasis *et al.* (1) reported cDNA sequences for two alleles of human *APOC3* that are distinguished by a polymorphic *Sst*I (position 371) in the 3' untranslated region. We have selected primers from this sequence for polymerase chain reaction (PCR) to amplify a 233 bp fragment that spans the *Sst*I polymorphism. The forward primer is 5'-GTGACCG-ATGGCTTCAGTTCCTG-3', the reverse primer is 5'-GGTAGGAGAGCACTGAGAATACTG-3'.

Protocol: In addition to the buffer and nucleotide components recommended by the supplier of *Taq* polymerase (Perkin-Elmer Cetus), each amplification reaction used 0.5 µg genomic DNA, 20 pMol of each primer, 10% dimethylsulfoxide, and 1 unit of *Taq* polymerase in a final reaction volume of 25 µl. Each reaction mixture was heated at 94° for 5 minutes for denaturation, and subjected to 30 cycles of PCR (primer annealing and elongation at 64° for 1.5 min, denaturation at 94° for 1 min). After amplification, 4 units of *Sst*I (Bethesda Research Laboratories) were added to each reaction (3 hrs at 37°). Digestion products were electrophoresed on nondenaturing 8% polyacrylamide gels (see Figure) or on 1.5% agarose gels.

Polymorphism: The A1 allele lacks the polymorphic *Sst*I site and produces a full-length PCR product of 233 bp. The A2 allele contains the *Sst*I site and produces 158 and 75 bp fragments.

Frequency: In 559 subjects, the frequency of the A1 allele was 0.88 and the frequency of the A2 allele was 0.12.

Chromosomal Localization: 11q23.

Mendelian Inheritance: Co-dominant segregation was shown in a three generation family by Sidoli *et al.* (2).

Acknowledgements: The DNA samples were from the multicentre study entitled 'Pathobiological Determinants of Atherosclerosis in Youth'. The work was supported by NIH grant HL39913.

References: 1) Karathanasis, S.K., Zannis, V.I. and Breslow, J.L. (1985) *J. Lipid Res.* **26**, 451-456. 2) Sidoli, A., Giudici, G., Soria, M. and Vergani, C. (1985) *Atherosclerosis* **62**, 81-87.

* To whom correspondence should be addressed

A *Pvu*II RFLP detected by pOB231 at CDC2 on human chromosome 10

N.L.Carson, S.Myers and N.E.Simpson*

Departments of Paediatrics and Biology, Queen's University, Kingston, Ontario, K7L 3N6, Canada

Source and Description: A 2.0 kb cDNA fragment cloned into the Okayama and Berg expression vector, pcD (1).

Polymorphism: A 2 allele co-dominant system identified in *Pvu*II digests. B1 is recognised as a 3.5 kb and B2 as a 4.5 kb band. Two constant bands are 10.0 kb and 5.5 kb.

Allele Frequencies: Estimated from 38 unrelated individuals. B1 0.68 B2 0.32.

Chromosomal Location: CDC2 was mapped to chromosome 10 in somatic cell hybrids (2) and localised to 10q21.1 by dosage studies (3). Linkage studies place the locus between D10S96 and EGR2/D10S19 (4).

Mendelian Inheritance: Co-dominant inheritance was observed in several MEN 2A families.

Not Polymorphic For: *Bgl*II, *Eco*RI, *Hinc*II, *Hind*III, *Msp*I, *Pst*I and *Stu*I.

Acknowledgements: We thank Drs. P.Nurse and M.Lee for providing pOB231. This study was supported by the Canadian Medical Research Council and an MRC studentship to N.L.C.

References: 1) Lee, M.G. and Nurse, P. (1987) *Nature* **327**, 31-35. 2) Spurr, N.K. *et al.* (1988) *Hum. Genet.* **78**, 333-337. 3) Carson, N.L. and Simpson, N.E. (1989) *Cytogenet. Cell Genet.* **51**, 974. 4) Wu, J. *et al.* (1990) *Genomics* (in press).

* To whom correspondence should be addressed