Published in final edited form as:

Periodontol 2000. 2012 June ; 59(1): 185–202. doi:10.1111/j.1600-0757.2011.00432.x.

Advanced regenerative technologies for periodontal tissue repair

Christoph A. Ramseier¹, Giulio Rasperini², Salvatore Batia², and William V. Giannobile^{3,4,*}

¹Department of Periodontology, School of Dental Medicine, University of Bern, Bern, Switzerland

²Unit of Periodontology, department of Surgical, Regenerative and Diagnostic Science, Foundation IRCCS Cà Granda Policlinico, University of Milan, Milan Italy ³Department of Periodontics and Oral Medicine and Michigan Center for Oral Health Research, University of Michigan, 1011 N. University Ave., Ann Arbor, MI 48109-1078, USA ⁴Department of Biomedical Engineering, College of Engineering, University of Michigan, 1011 N. University Ave., Ann Arbor, MI 48109-1078, USA

Keywords

periodontal regeneration; bone grafts; growth factors; periodontal disease; periodontal therapy

Regenerative periodontal therapy comprises techniques which are particularly designed to restore those parts of the tooth-supporting structures which have been lost due to periodontitis or gingival trauma. The term ‘regeneration’ is defined as a reconstruction of lost or injured tissues in such a way that both the original structures and their function are completely restored. Procedures aimed at restoring lost periodontal tissues favor the creation of new attachment, including the formation a new periodontal ligament with its fibers inserting in newly formed cementum and alveolar bone.

Deep infra-bony defects associated with periodontal pockets are the classic indication for periodontal regenerative therapy. Additionally, different degrees of furcation involvement in molars and upper first premolars are a further indication for regenerative approaches as the furcation area remains difficult to maintain through instrumentation and oral hygiene. A third group of indications for regenerative periodontal therapy are localized gingival recessions and root exposure since they may cause a significant esthetic concern for the patient. The denuding of a root surface with resultant root sensitivity represents a further indication to apply regenerative periodontal therapy in order to achieve both the reduction of root sensitivity and the improvement of esthetics.

Professional periodontal therapy and maintenance, combined with risk factor control, are shown to effectively reduce periodontal disease progression. In contrast to the conventional approaches of anti-inflammatory periodontal therapy, however, the regenerative procedures aimed at repairing lost periodontal tissues, including alveolar bone, periodontal ligament and root cementum, remain more challenging (22). Periodontal research in the past few decades has attempted to systematically determine predictably successful clinical procedures to regenerate periodontal tissues. Hence, various methods in combination with regenerative biomaterials, such as hard- and soft-tissue grafts, or cell occlusive barrier membranes used

*Corresponding author: William V. Giannobile, DDS, D.Med.Sc., University of Michigan School of Dentistry, Michigan Center for Oral Health Research, 24 Frank Lloyd Wright Drive, Lobby M, Box 422, Ann Arbor, MI 48106, USA, Tel: (734) 998-1468, Fax: (734) 998-7228, wgiannob@umich.edu.

in guided tissue regeneration procedures, have been pursued to regenerate lost tooth support (160).

Periodontal regeneration is assessed by probing measures, radiographic analysis, direct measurements of new bone and histology (129). Many cases that are considered clinically successful, including cases with significant re-growth of alveolar bone, may histologically still show an epithelial lining along the treated root surface instead of newly formed periodontal ligament and cementum (81). In general, however, the clinical outcome of periodontal regenerative techniques is shown to depend on 1) patient associated factors such as plaque control, smoking habits, residual periodontal infection, or membrane exposure in guided tissue regeneration procedures, 2) effects of occlusal forces that deliver intermittent loads in axial and transverse dimensions, as well as 3) factors associated with the clinical skills of the operator such as lack of primary closure of the surgical wound (90). Even though modified flap designs and microsurgical approaches are shown to positively affect the outcome of both soft and hard tissue regeneration, the clinical success for periodontal regeneration still remains limited in many cases. Moreover, the surgical protocols for regenerative procedures are skill-demanding and may therefore lack practicability for a number of clinicians. Consequently, both clinical and pre-clinical research continues to evaluate advanced regenerative approaches using either new barrier membrane techniques (67), cell-growth stimulating proteins (26, 42, 68) or gene delivery applications (125), respectively, in order to simplify and enhance the rebuilding of missing periodontal support. The aim of our review is to compare these advanced regenerative concepts for periodontal hard and soft tissue repair with conventional regenerative techniques (Table 1). While a focus will be given on clinical applications with the delivery of growth factors, the applications for gene delivery of tissue growth factors are also reviewed.

Periodontal wound healing

Research on periodontal wound healing in the past was able to provide the basic understanding of the mechanisms favoring periodontal tissue regeneration. A number of valuable findings at both the cellular and molecular levels was revealed and subsequently used for the engineering of regenerative biomaterials available in periodontal medicine today. In order to provide an overview of the cellular and molecular events and their relation to periodontal tissue regeneration, the course of periodontal wound healing is briefly reviewed in this article.

The biology and principles of periodontal wound healing have previously been reviewed (120). Based on observations following experimental incisions in periodontal soft tissues, after blood clot formation, the sequence of healing is commonly divided into the following phases: 1) soft tissue inflammation; 2) granulation tissue formation; and 3) intercellular matrix formation and remodeling (20, 146). Plasma proteins, mainly fibrinogen, dominate rapidly in the bleeding wound and provide an initial basis for the adherence of a fibrin clot (165). The inflammatory phase of healing in the soft tissue wound is initiated by polymorphonuclear leukocytes infiltrating the fibrin clot from the wound margins shortly followed by macrophages (111). The major function of the polymorphonuclear leukocytes is to debride the wound by removing bacterial cells and injured tissue particles through phagocytosis. The macrophages, in addition, have an important role to play in the initiation of tissue repair. The inflammatory phase progresses into its later stage as the polymorphonuclear leukocyte infiltrate gradually decreases while the macrophage influx continues. These macrophages contribute to the cleansing process by phagocytosis of used polymorphonuclear leukocytes and erythrocytes. Additionally, macrophages release a number of biologically active molecules such as inflammatory cytokines and tissue growth factors, which recruit further inflammatory cells as well as fibroblastic and endothelial cells,

thus playing an essential role in the transition of the wound from the inflammation into the granulation tissue formation. The influx of fibroblasts and budding capillaries from the gingival connective tissue and the periodontal ligament connective tissue initiate the phase of granulation tissue formation in the periodontal wound approximately two days after incision. At this stage, fibroblasts are responsible for the formation of a loose new matrix of collagen, fibronectin, and proteoglycans (11). Eventually, cells and matrix form cell-to-cell and cell-to-matrix links that generate a concerted tension resulting in tissue contraction. The phase of granulation tissue formation gradually develops into the final phase of healing in which the reformed, more cell-rich tissue undergoes maturation and sequenced re-modeling to meet functional needs (20, 146).

The morphology of a periodontal wound comprises 1) the gingival epithelium, 2) the gingival connective tissue, 3) the periodontal ligament, and the hard tissue components such as 4) alveolar bone and 5) cementum or dentin on the dental root surface (Fig. 1). This particular composition ultimately affects both the healing events in each tissue component as well as in the entire periodontal site. While the healing of gingival epithelia and their underlying connective tissues concludes in a number of weeks, the regeneration of periodontal ligament, root cementum and alveolar bone generally only occur within a number of weeks or months. Aiming for wound closure, the final outcome of wound healing in the epithelium is the formation of the junctional epithelium surrounding the dentition (15). The healing of gingival connective tissue on the other hand, results in a significant reduction of its volume, thus clinically creating both gingival recession and a reduction of the periodontal pocket. Periodontal ligament is shown to regenerate on newly formed cementum created by cementoblasts originated from periodontal ligament granulation tissue (71). Furthermore, alveolar bone modeling occurs following the stimulation of mesenchymal cells from the gingival connective tissue that are transformed into osteoprogenitor cells by locally expressed bone morphogenetic proteins (75, 149).

A series of classical animal studies were able to demonstrate that the tissue derived from alveolar bone or gingival connective tissue lack cells with the potential to produce a new attachment between the periodontal ligament and newly formed cementum (72, 109). Moreover, granulation tissue derived from the gingival connective tissue or alveolar bone result in root resorption or ankylosis when placed in contact with the root surface. It should be expected, therefore, that these complications would occur more frequently following regenerative periodontal surgery, particularly following those procedures which include the placement of grafting materials to stimulate bone formation. The reason for root resorption (which is rarely observed), however, may be that following the surgical intervention, the dento-gingival epithelium migrates apically along the root surface, forming a protective barrier towards the root surface (10, 73). The findings from these animal experiments revealed that ultimately the periodontal ligament tissue contains the cells with the potential to form a new connective tissue attachment (71).

Typically, the downgrowth of the epithelium along the tooth root surface reaches the level of the periodontal ligament before the latter has regenerated with new layers of cementum and newly inserting connective tissue fibers. Therefore, in order to enable and promote healing towards the rebuilding of cementum and periodontal ligament, the gingival epithelium must be prevented from forming a long junctional epithelium along the root surface down to the former level of the periodontal ligament (Fig. 2). This basic acquisition of knowledge has been the key for the engineering of standard clinical procedures for the placement of a fabricated membrane in guided tissue regeneration.

In summary, the principles of periodontal wound healing presented provide the basic understanding of the events following wounding in surgical interventions. In order to obtain

new connective tissue attachment, the granulation tissue derived from periodontal ligament cells has to be given both space and time to format and mature new cementum and periodontal ligament. The conventional guided tissue regeneration techniques in periodontal practice have shown their predictable, however limited, potential to regenerate lost periodontal support. Consequently, advanced regenerative technologies for periodontal tissue repair aim to increase the current gold standards for success of periodontal regeneration. In order to identify advanced repair of tooth-supporting periodontal tissues a number of combinations of conventional regenerative techniques have been evaluated:

Periodontal and alveolar bone repair

- Guided tissue regeneration & application of tissue growth factor(s)
- Guided tissue regeneration & hard tissue graft & application of tissue growth factor(s)
- Hard tissue graft & bio-modification of the tooth root surface
- Hard tissue graft & application of tissue growth factors

Advanced repair of alveolar bone defects

The morphology of the alveolar infra-bony defect was shown to play a significant role for the establishment of a predictable outcome of regeneration of periodontal attachment (121). Goldman and Cohen originally proposed a classification for infra-bony defects which referred to the number of osseous walls surrounding the defect, either being one-, two- or three-wall, respectively (48).

Hard tissue grafts

In a number of clinical trials and animal experiments, the periodontal flap approach was combined with the placement of bone grafts or implant materials into the curetted bony defects with the aim of favoring periodontal regeneration. The various graft and implant materials evaluated so far can be listed as follows:

1. Autogenous graft: graft transferred from one location to another within the same organism
2. Allogenic graft: graft transferred from one organism to another organism of the same species
3. Xenogenic graft: graft taken from an organism of a different species
4. Alloplastic material: synthetic or inorganic implant material used instead of the previously mentioned graft material.

The biologic rationale behind the use of bone grafts or alloplastic materials for regenerative approaches is the assumption that these materials may serve as a scaffold for bone formation (osteoconduction), contain the bone forming cells (osteogenesis) or bone inductive substances (osteinduction).

Histological studies in both humans and animals have demonstrated that grafting procedures often result in healing with a long junctional epithelium rather than a new connective tissue attachment (16, 81). Therefore, multiple studies have evaluated the use of hard tissue graft materials for periodontal regeneration in infra-bony defects when compared to the periodontal flap approach alone.

Bio-modification of the tooth root surface

A number of studies have focused on the modification of the periodontitis-involved root surface in order to advance the formation of a new connective tissue attachment. However, despite histological evidence of regeneration following root surface bio-modification with citric acid, the outcome of controlled clinical trials have failed to show any improvements in clinical conditions compared to non-acid treated controls (38, 88, 96).

In recent years, bio-modification of the root surface with enamel matrix proteins during periodontal surgery and following demineralization with ethylene-diamine-tetra-acetic acid (EDTA) has been introduced to promote periodontal regeneration. Based on the understanding of the biological model, the application of enamel matrix proteins (amelogenins) is seen to promote periodontal regeneration since it initiates events that occur during the growth of periodontal tissues (41, 52). The commercially available product Emdogain®, a purified acid extract of porcine origin contains enamel matrix derivatives, is reported to be able to enhance periodontal regeneration (Fig. 3). More basic research, in addition to the clinical findings, indicates that enamel matrix derivatives have a key role in periodontal wound healing (24, 30). Histological results from both animal and human studies have shown that the application of enamel matrix derivatives promotes periodontal regeneration and confidently influences periodontal wound healing (143). Thus far, enamel matrix derivatives either alone or in combination with grafts have demonstrated its potential to effectively treat intra-osseous defects and the clinical results appear to be stable long term (155).

Periodontal tissue growth factors

Wound healing approaches using growth factors to target restoration of tooth-supporting bone, periodontal ligament, and cementum has been shown to significantly advance the field of periodontal regenerative medicine. A major focus of periodontal research has studied the impact of tissue growth factor applications on periodontal tissue regeneration (Table 2) (3, 42, 101, 123). Advances in molecular cloning have made available unlimited quantities of recombinant growth factors for applications in tissue engineering. Recombinant growth factors known to promote skin and bone wound healing, such as platelet-derived growth factors (13, 44, 65, 106, 112, 136), insulin like growth factors (42, 44, 56, 84), fibroblast growth factors (47, 98, 145, 151, 152) and bone morphogenetic proteins (40, 57, 147, 162, 163) have been used in pre-clinical and clinical trials for the treatment of large periodontal or infra-bony defects, as well as around dental implants (34, 66, 106). The combined use of rhplatelet-derived growth factor-BB and peptide P-15 with a graft biomaterial has shown beneficial effects in intra-osseous defects (155). However, contrasting results were reported for growth factors such as platelet-rich plasma and graft combinations, or the use of BAs either alone or in association with graft or guided tissue regeneration for the treatment of furcation defects (155).

Biological effects of growth factors: platelet-derived growth factor—Platelet-derived growth factor is a member of a multifunctional polypeptide family that binds to two cell membrane tyrosine kinase receptors (platelet-derived growth factor- R_{α} and platelet-derived growth factor- R_{β}) and subsequently exerts its biological effects on cell proliferation, migration, extracellular matrix synthesis, and anti-apoptosis (54, 69, 134, 144). Platelet-derived growth factor- α and - β receptors are expressed in regenerating periodontal soft and hard tissues (116). In addition, platelet-derived growth factor initiates tooth-supporting periodontal ligament cell chemotaxis (108), mitogenesis (110), matrix synthesis (51), and attachment to tooth dentinal surfaces (170). More importantly, in vivo application of platelet-derived growth factor alone or in combination with insulin-like growth factor-1 results in partial repair of periodontal tissues (44, 45, 84, 85, 136). Platelet-derived growth

factor has been shown to have a significant regenerative impact on periodontal ligament cells as well as on osteoblasts (87, 89, 110, 112).

The clinical application of platelet-derived growth factor was shown to successfully advance alveolar bone repair and clinical attachment level gain. A first clinical study reported the successful repair of class II furcations using demineralized freeze-dried bone allograft saturated with rhplatelet-derived growth factor-BB (105). Subsequently, rhplatelet-derived growth factor-BB mixed with a synthetic beta-tricalcium phosphate matrix was shown to advance the repair of deep infra-bony pockets in a large multicenter randomized controlled trial (106). Both studies demonstrated that the use of rhplatelet-derived growth factor-BB was safe and effective in the treatment of periodontal osseous defects. In a follow-up trial, the same sample patients were again assessed 18 or 24 months following periodontal surgery. Substantial radiographic changes in the appearance of the defect fill were observed for patients treated with rhplatelet-derived growth factor-BB (91).

Biological effects of growth factors: bone morphogenetic proteins—Bone morphogenetic proteins are multifunctional polypeptides belonging to the transforming growth factor- β superfamily of proteins (167). The human genome encodes at least twenty bone morphogenetic proteins (128). Bone morphogenetic proteins bind to type I and II receptors that function as serine-threonine kinases. The type I receptor protein kinase phosphorylates intracellular signaling substrates called Smads (Sma gene in *Caenorhabditis elegans* and Mad gene in *Drosophila*). The phosphorylated bone morphogenetic protein-signaling Smads enter the nucleus and initiate the production of bone matrix proteins leading to bone morphogenesis. The most remarkable feature of bone morphogenetic proteins is the ability to induce ectopic bone formation (158). Bone morphogenetic proteins are not only powerful regulators of cartilage and bone formation during embryonic development and regeneration in postnatal life, but also participate in the development and repair of other organs such as the brain, kidney, and nerves (127).

Sigurdsson et al. (145) evaluated bone and cementum formation following regenerative periodontal surgery using recombinant human bone morphogenetic protein in surgically-created supra-alveolar defects in dogs (166). Histologic analysis showed significantly more cementum formation and regrowth of alveolar bone on bone morphogenetic protein treated sites as compared to the controls.

Studies have demonstrated the expression of bone morphogenetic proteins during tooth development and periodontal repair including alveolar bone (1, 2). Investigations in animal models have shown the potential repair of alveolar bony defects using recombinant human bone morphogenetic protein-12 (rhBMP-12) (163) or rhBMP-2 (83, 164). In a clinical trial by Fiorellini et al. (34), recombinant human bone morphogenetic protein-2 (rhBMP-2) delivered by a bioabsorbable collagen sponge revealed significant bone formation in a human buccal wall defect model following tooth extraction when compared to collagen sponge alone. Furthermore, bone morphogenetic protein-7, also known as osteogenic protein-1, stimulates bone regeneration around teeth, endosseous dental implants, and in maxillary sinus floor augmentation procedures (47, 137, 159).

Clinical applications of growth factors for use in periodontal regeneration

In general, the impact of a topical delivery of growth factors to periodontal wounds has shown to be promising, yet insufficient for the promotion of predictable periodontal tissue engineering (13, 21) (Fig. 4). Growth factor proteins, once delivered to the target site, tend to suffer from instability and quick dilution, presumably due to proteolytic breakdown, receptor-mediated endocytosis, and solubility of the delivery vehicle (3). Because their half-lives are significantly reduced, the period of exposure may not be sufficient to act on

osteoblasts, cementoblasts, or periodontal ligament cells. Therefore different methods of growth factor delivery need to be considered (4).

Investigations for periodontal bioengineering have examined a variety of methods combining delivery vehicles, such as scaffolds, with growth factors to target the defect site in order to optimize bioavailability (82). The scaffolds are designed to optimize the dosage of the growth factor and to control its release pattern which may be pulsatile, constant or time programmed (7). Additionally, the kinetics of the release and the duration of the exposure of the growth factor may be controlled (59).

A new polymeric system was reported in an animal study by Richardson et al. (133) enabling the tissue-specific delivery of two or more growth factors, with a controlled dose and rate of delivery. The dual delivery of vascular endothelial growth factor together with platelet-derived growth factor from a single, structural polymer scaffold results in the rapid formation of a mature vascular network (133).

Guided tissue regeneration

Histological findings from periodontal regeneration studies reveal that a new connective tissue attachment could be predicted if the cells from the periodontal ligament settle on the root surface during healing. Hence, the clinical applications of guided tissue regeneration in periodontics involve the placement of a physical barrier membrane to enable the previous periodontitis-affected tooth root surface to be repopulated with cells from the periodontal ligament. In the last decades, guided tissue regeneration has been applied in many clinical trials for the treatment of various periodontal defects, such as infra-bony defects (23), furcation involvements (70, 86), and localized gingival recessions (118). In a recent systematic review, the combinations of barrier membranes and grafting materials used in preclinical models have been summarized. The analysis of ten papers revealed that the combination of barrier membranes and grafting materials may result in histological evidence of periodontal regeneration, predominantly bone repair. No additional histological benefits of combination treatments were found in animal models of three wall intrabony, class II furcation, or fenestration defects. In supra-alveolar and two wall intrabony defect models of periodontal regeneration, the additional use of a grafting material gave superior histological results of bone repair to barrier membranes alone (141).

The types of barrier membranes evaluated in clinical studies vary regarding design, configuration, and composition. Non-resorbable membranes of expanded polytetrafluoroethylene have been used successfully in both animal experiments and human clinical trials. In recent years, natural or synthetic bio-absorbable barrier membranes have been used for guided tissue regeneration in order to eliminate the need for a follow-up surgery for membrane removal. Collagen membranes as well as barrier materials of polylactic acid or copolymers of polylactic acid and polyglycolic acid have been tested in animal and human studies.

Following therapy, guided tissue regeneration has a greater effect on probing measures of periodontal treatment than periodontal flap surgery alone, including increased attachment gain, reduction of probing depth, less gingival recession and more gain in hard tissue probing at surgical re-entry. Referring to the best available evidence today, however, it is difficult to draw general conclusions about the clinical benefit of guided tissue regeneration. Even as there is evidence that guided tissue regeneration can demonstrate a significant improvement over conventional open flap surgery, the factors affecting outcomes are unclear from the present literature since they might comprise study conduct issues such as bias (103).

In summary, guided tissue regeneration represents a very well documented regenerative procedure today to achieve periodontal regeneration in infra-bony defects and in degree II furcations. An added benefit may be achieved by the additional use of grafting materials (153).

Gene therapeutics for periodontal tissue repair

Although encouraging results for periodontal regeneration have been found in various clinical investigations using recombinant tissue growth factors, limitations exist with topical protein delivery such as transient biological activity, protease inactivation, and poor bioavailability from existing delivery vehicles. Therefore, newer approaches seek to develop methodologies that optimize growth factor targeting to maximize the therapeutic outcome of periodontal regenerative procedures. Genetic approaches in periodontal tissue engineering show early progress in achieving delivery of growth factor genes such as platelet-derived growth factor or bone morphogenetic protein to periodontal lesions (Fig. 5). Gene transfer methods may circumvent many of the limitations with protein delivery to soft tissue wounds (9, 43). It has been shown that the application of growth factors (35, 61, 63, 75) or soluble forms of cytokine receptors (150) by gene transfer provides a greater sustainability than that of single protein application. Thus, gene therapy may achieve greater bioavailability of growth factors within periodontal wounds and thus providing greater regenerative potential.

Methods for gene delivery in periodontal applications

Various gene delivery methods are available to administer growth factors to periodontal defects offering great flexibility for tissue engineering. The delivery method can be tailored to the specific characteristics of the wound site. For example, a horizontal one or two-walled defect may require the use of a supportive carrier like a scaffold. Other defect sites may be conducive to the use of an adenovirus-vector embedded in a collagen matrix.

More importantly from a clinical point of view is the risk associated with the use of gene therapy in periodontal tissue engineering (49). As with maximizing growth factor sustainability and accounting for specific characteristics of the wound site, both the DNA vector and delivery method need to be considered when assessing patient safety. In summary, studies examining the use of specific delivery methods and DNA vectors in periodontal tissue engineering reflect the aim to maximize the duration of growth factor expression, optimize delivery method to periodontal defect, and minimize patient risk.

A combination of an AAV-delivered angiogenic molecule such as vascular endothelial growth factor, bone morphogenetic protein signaling receptor (caALK2) and receptor activator of nuclear factor-kappa B ligand (RANKL) were demonstrated to promote bone allograft turnover and osteogenesis as a mode to enrich human bone allografts (60). To date, combinations of vascular endothelial growth factor/bone morphogenetic protein (117) and platelet-derived growth factor/vascular endothelial growth factor (133) have been performed with highly positive synergistic responses in bone repair.

Promising preliminary results from preclinical studies reveal that host modulation achieved through gene delivery of soluble proteins such as tumor necrosis factor receptor 1 (TNFR1:Fc) reduces tumor necrosis factor activity and therefore inhibits alveolar bone loss (150). These results are comparable to the findings in the research on rheumatoid arthritis where pathogenesis includes high tumor necrosis factor activity, and pathways for bone resorption are similar (124).

Preclinical studies evaluating growth factor gene therapy for periodontal tissue engineering

In order to overcome the short half-lives of growth factor peptides *in vivo*, gene therapy using a vector encoding the growth factor is advocated to stimulate tissue regeneration. So far, two main strategies of gene vector delivery have been applied to periodontal tissue engineering. Gene vectors can be introduced directly to the target site (*in vivo* technique) (61) or selected cells can be harvested, expanded, genetically transduced and then re-implanted (*ex vivo* technique) (63). *In vivo* gene transfer involves the insertion of the gene of interest directly into the body anticipating the genetic modification of the target cell. *Ex vivo* gene transfer includes the incorporation of genetic material into cells exposed from a tissue biopsy with subsequent re-implantation into the recipient. Using the *in vivo* technique, the potential inhibition of alveolar bone loss has been studied in an experimental periodontitis model evaluating the inhibition of osteoclastogenesis by administering human osteoprotegerin, a competitive inhibitor of the RANKL-derived osteoclast activation. Significant preservation of alveolar bone volume was observed among osteoprotegerin:Fc-treated animals compared to the controls. Systemic delivery of osteoprotegerin:Fc inhibits alveolar bone resorption in experimental periodontitis, suggesting that RANKL inhibition may represent an important therapeutic strategy for the prevention of progressive alveolar bone loss (62).

Platelet-derived growth factor gene delivery—The application of platelet-derived growth factor-gene transfer strategies to tissue engineering has originally been generated to improve healing in soft tissue wounds such as skin lesions (25). Both plasmid- (55) and adenovirus/platelet-derived growth factor gene delivery (122) have been evaluated in preclinical and human trials. However, the latter has been able to exhibit more safety favorable for clinical use (49). In a recent animal study reporting on safety and distribution profiles, adenovirus-platelet-derived growth factor-B applied for tissue engineering of tooth-supporting alveolar bone defects was well contained within the localized osseous defect area without viremia or distant organ involvement (17).

Early studies in dental applications using recombinant adenoviral vectors encoding platelet-derived growth factor demonstrated the ability of these vector constructs to potently transduce cells isolated from the periodontium (osteoblasts, cementoblasts, periodontal ligament cells, and gingival fibroblasts) (46, 171). These studies revealed the extensive and prolonged transduction of periodontal-derived cells. Both Chen & Giannobile (18) and Lin et al. (79) were able to demonstrate the effects of adenoviral delivery of platelet-derived growth factor for the better understanding of sustained platelet-derived growth factor signaling. Gene delivery of platelet-derived growth factor-B generally displays higher sustained signal transduction effects in human gingival fibroblasts when compared to cells treated with recombinant human platelet-derived growth factor-BB protein alone. Their data on platelet-derived growth factor gene delivery may contribute to an improved understanding of these pathways that are likely to play a role in the control of clinical outcomes of periodontal regenerative therapy.

In an *ex vivo* investigation by Anusaksathien et al. (6) it was shown that the expression of platelet-derived growth factor genes was prolonged for up to 10 days in gingival wounds. Adenovirus encoding platelet-derived growth factor-B (adenovirus/platelet-derived growth factor-B) transduced gingival fibroblasts and enhanced defect fill by induction of human gingival fibroblast migration and proliferation (6). On the other hand, continuous exposure of cementoblasts to platelet-derived growth factor-A had an inhibitory effect on cementum mineralization, possibly via the upregulation of osteopontin and subsequent enhancement of multinucleated giant cells in cementum engineered scaffolds. Moreover, adenovirus/platelet-

derived growth factor-1308 (a dominant-negative mutant of platelet-derived growth factor) inhibited mineralization of tissue-engineered cementum possibly due to downregulation of bone sialoprotein and osteocalcin with a persistence of stimulation of multinucleated giant cells. These findings suggest that continuous exogenous delivery of platelet-derived growth factor-A may delay mineral formation induced by cementoblasts, while platelet-derived growth factor is clearly required for mineral neogenesis (5).

Jin et al. (61) demonstrated that direct in vivo gene transfer of platelet-derived growth factor-B was able to stimulate tissue regeneration in large periodontal defects. Descriptive histology and histomorphometry revealed that human platelet-derived growth factor-B gene delivery promotes the regeneration of both cementum and alveolar bone, while platelet-derived growth factor-1308, a dominant negative mutant of platelet-derived growth factor-A, has minimal effects on periodontal tissue regeneration.

Bone morphogenetic protein gene delivery—An experimental study in rodents by Lieberman et al. (78) advanced gene therapy for bone regeneration with results revealing that the transduction of bone marrow stromal cells with rhBMP-2 lead to bone formation within an experimental defect comparable to skeletal bone. Another group was similarly able to regenerate skeletal bone by directly administering adenovirus5/BMP-2 into a bony segmental defect in rabbits (8). Further advances in the area of orthopedic gene therapy using viral delivery of bone morphogenetic protein-2 have provided further evidence for the ability of both in vivo and ex vivo bone engineering (19, 76, 77, 100). Franceschi et al. (35) investigated in vitro and in vivo adenovirus gene transfer of bone morphogenetic protein-7 for bone formation. Adenovirus transduced nonosteogenic cells were also found to differentiate into bone-forming cells and to produce bone morphogenetic protein-7 (75) or bone morphogenetic protein-2 (19) both in vitro and in vivo. In another study by Huang et al. (58), plasmid DNA encoding for bone morphogenetic protein-4 administered with a scaffold delivery system was found to enhance bone formation when compared with blank scaffolds.

In an early approach to regenerate alveolar bone in an animal model, it was demonstrated that the ex vivo delivery of adenovirus encoding murine bone morphogenetic protein-7 was found to promote periodontal tissue regeneration in large mandibular periodontal bone defects (63). Bone morphogenetic protein-7 gene transfer not only enhanced alveolar bone repair, it also stimulated cementogenesis and periodontal ligament fiber formation. Of interest, the alveolar bone formation was found to occur via a cartilage intermediate. However, when genes encoding the bone morphogenetic protein antagonist noggin were delivered, inhibition of periodontal tissue formation resulted (64). In a study by Dunn et al. (28), it was shown that direct in vivo gene delivery of adenovirus/bone morphogenetic protein-7 in a collagen gel carrier promoted successful regeneration of alveolar bone defects around dental implants. Furthermore, an in vivo synergism of adenoviral-mediated coexpression of bone morphogenetic protein-7 and insulin like growth factor-1 on human periodontal ligament cells in up-regulating alkaline phosphatase activity and mRNA levels of collagen type I and Runx2 was found (168). Implantation with scaffolds illustrated that the transduced cells exhibited osteogenic differentiation and formed bone-like structures. It was concluded that the combined delivery of bone morphogenetic protein-7 and insulin like growth factor-1 genes using an internal ribosome entry site-based strategy synergistically enhanced differentiation of human periodontal ligament cells (168).

These experiments provide promising evidence showing the feasibility of both in vivo and ex vivo gene therapy for periodontal tissue regeneration and peri-implant osseointegration.

Future perspectives: targeted gene therapy in vivo

Major advances have been made over the past decade in the reconstruction of complex periodontal and alveolar bone wounds that have resulted from disease or injury. Developments in scaffolding matrices for cell, protein and gene delivery have demonstrated significant potential to provide “smart” biomaterials that can interact with the matrix, cells and bioactive factors. The targeting of signaling molecules or growth factors (via proteins or genes) to periodontal tissue components has lead to significant new knowledge generation using factors that promote cell replication, differentiation, matrix biosynthesis and angiogenesis. A major challenge that has been less studied is the modulation of the exuberant host response to microbial contamination that plagues the periodontal wound microenvironment. For improvements in the outcomes in periodontal regenerative medicine, scientists will need to examine dual delivery of host modifiers or anti-infective agents to optimize the results of therapy. Further advancements in the field will continue to rely heavily on multidisciplinary approaches combining engineering, dentistry, medicine, and infectious disease specialists in repairing the complex periodontal wound environment.

Acknowledgments

This work was supported by NIH/NIDCR DE13397 and NIH/NCRR UL1RR-024986. The authors thank Mr. Chris Jung for his assistance with the figures.

References

1. Aberg T, Wozney J, Thesleff I. Expression patterns of bone morphogenetic proteins (BMPs) in the developing mouse tooth suggest roles in morphogenesis and cell differentiation. *Dev Dyn.* 1997; 210:383–396. [PubMed: 9415424]
2. Amar S, Chung KM, Nam SH, Karatzas S, Myokai F, Van Dyke TE. Markers of bone and cementum formation accumulate in tissues regenerated in periodontal defects treated with expanded polytetrafluoroethylene membranes. *J Periodontal Res.* 1997; 32:148–158. [PubMed: 9085226]
3. Anusaksathien O, Giannobile WV. Growth factor delivery to re-engineer periodontal tissues. *Curr Pharm Biotechnol.* 2002; 3:129–139. [PubMed: 12022256]
4. Anusaksathien O, Jin Q, Ma PX, Giannobile WV. Scaffolding in Periodontal Engineering. In: Ma, PX, Eliseeff, J., editors. *Scaffolding in Tissue Engineering*. Boca Raton, FL: CRC Press; 2005. p. 427–444.
5. Anusaksathien O, Jin Q, Zhao M, Somerman MJ, Giannobile WV. Effect of sustained gene delivery of platelet-derived growth factor or its antagonist (PDGF-1308) on tissue-engineered cementum. *J Periodontol.* 2004; 75:429–440. [PubMed: 15088882]
6. Anusaksathien O, Webb SA, Jin QM, Giannobile WV. Platelet-derived growth factor gene delivery stimulates ex vivo gingival repair. *Tissue Eng.* 2003; 9:745–756. [PubMed: 13678451]
7. Babensee JE, McIntire LV, Mikos AG. Growth factor delivery for tissue engineering. *Pharm Res.* 2000; 17:497–504. [PubMed: 10888299]
8. Baltzer AW, Lattermann C, Whalen JD, Wooley P, Weiss K, Grimm M, Ghivizzani SC, Robbins PD, Evans CH. Genetic enhancement of fracture repair: healing of an experimental segmental defect by adenoviral transfer of the BMP-2 gene. *Gene Ther.* 2000; 7:734–739. [PubMed: 10822299]
9. Baum BJ, Goldsmith CM, Kok MR, Lodde BM, van Mello NM, Voutetakis A, Wang J, Yamano S, Zheng C. Advances in vector-mediated gene transfer. *Immunol Lett.* 2003; 90:145–149. [PubMed: 14687716]
10. Bjorn H, Hollender L, Lindhe J. Tissue regeneration in patients with periodontal disease. *Odontol Revy.* 1965; 16:317–326. [PubMed: 5216253]
11. Briggs SL. The role of fibronectin in fibroblast migration during tissue repair. *J Wound Care.* 2005; 14:284–287. [PubMed: 15974416]

12. Camelo M, Nevins ML, Lynch SE, Schenk RK, Simion M, Nevins M. Periodontal regeneration with an autogenous bone-Bio-Oss composite graft and a Bio-Gide membrane. *Int J Periodontics Restorative Dent*. 2001; 21:109–119. [PubMed: 11829385]
13. Camelo M, Nevins ML, Schenk RK, Lynch SE, Nevins M. Periodontal regeneration in human Class II furcations using purified recombinant human platelet-derived growth factor-BB (rhPDGF-BB) with bone allograft. *Int J Periodontics Restorative Dent*. 2003; 23:213–225. [PubMed: 12854772]
14. Camelo M, Nevins ML, Schenk RK, Simion M, Rasperini G, Lynch SE, Nevins M. Clinical, radiographic, and histologic evaluation of human periodontal defects treated with Bio-Oss and Bio-Gide. *Int J Periodontics Restorative Dent*. 1998; 18:321–331. [PubMed: 12693419]
15. Caton J, Nyman S, Zander H. Histometric evaluation of periodontal surgery. II. Connective tissue attachment levels after four regenerative procedures. *J Clin Periodontol*. 1980; 7:224–231. [PubMed: 7000854]
16. Caton J, Zander HA. Osseous repair of an infrabony pocket without new attachment of connective tissue. *J Clin Periodontol*. 1976; 3:54–58. [PubMed: 815276]
17. Chang PC, Cirelli JA, Jin Q, Seol YJ, Sugai JV, D'Silva NJ, Danciu TE, Chandler LA, Sosnowski BA, Giannobile WV. Adenovirus encoding human platelet-derived growth factor-B delivered to alveolar bone defects exhibits safety and biodistribution profiles favorable for clinical use. *Hum Gene Ther*. 2009; 20:486–496. [PubMed: 19199824]
18. Chen QP, Giannobile WV. Adenoviral gene transfer of PDGF downregulates gas gene product PDGFalphaR and prolongs ERK and Akt/PKB activation. *Am J Physiol Cell Physiol*. 2002; 282:C538–C544. [PubMed: 11832339]
19. Cheng SL, Lou J, Wright NM, Lai CF, Avioli LV, Riew KD. In vitro and in vivo induction of bone formation using a recombinant adenoviral vector carrying the human BMP-2 gene. *Calcif Tissue Int*. 2001; 68:87–94. [PubMed: 11310352]
20. Clark RA. Biology of dermal wound repair. *Dermatol Clin*. 1993; 11:647–666. [PubMed: 8222349]
21. Cochran DL, Wozney JM. Biological mediators for periodontal regeneration. *Periodontol* 2000. 1999; 19:40–58. [PubMed: 10321215]
22. Cortellini P. Reconstructive periodontal surgery: a challenge for modern periodontology. *Int Dent J*. 2006; 56:250–255. [PubMed: 16972400]
23. Cortellini P, Bowers GM. Periodontal regeneration of intrabony defects: an evidence-based treatment approach. *Int J Periodontics Restorative Dent*. 1995; 15:128–145. [PubMed: 8593979]
24. Cortellini P, Tonetti MS. Clinical performance of a regenerative strategy for intrabony defects: scientific evidence and clinical experience. *J Periodontol*. 2005; 76:341–350. [PubMed: 15857066]
25. Crombleholme TM. Adenoviral-mediated gene transfer in wound healing. *Wound Repair Regen*. 2000; 8:460–472. [PubMed: 11208173]
26. Dereka XE, Markopoulou CE, Vrotsos IA. Role of growth factors on periodontal repair. *Growth Factors*. 2006; 24:260–267. [PubMed: 17381067]
27. Dryankova MM, Popova CL. Regenerative therapy of furcation defect. *Folia Med (Plovdiv)*. 2001; 43:64–68. [PubMed: 15354471]
28. Dunn CA, Jin Q, Taba M Jr, Franceschi RT, Bruce Rutherford R, Giannobile WV. BMP gene delivery for alveolar bone engineering at dental implant defects. *Mol Ther*. 2005; 11:294–299. [PubMed: 15668141]
29. Ellegaard B, Loe H. New attachment of periodontal tissues after treatment of intrabony lesions. *J Periodontol*. 1971; 42:648–652. [PubMed: 4937566]
30. Esposito M, Coulthard P, Thomsen P, Worthington HV. Enamel matrix derivative for periodontal tissue regeneration in treatment of intrabony defects: a Cochrane systematic review. *J Dent Educ*. 2004; 68:834–844. [PubMed: 15286106]
31. Esposito M, Coulthard P, Worthington HV. Enamel matrix derivative (Emdogain) for periodontal tissue regeneration in intrabony defects. *Cochrane Database Syst Rev*. 2003 CD003875.
32. Esposito M, Grusovin MG, Coulthard P, Worthington HV. Enamel matrix derivative (Emdogain) for periodontal tissue regeneration in intrabony defects. *Cochrane Database Syst Rev*. 2005 CD003875.

33. Fetner AE, Hartigan MS, Low SB. Periodontal repair using PerioGlas in nonhuman primates: clinical and histologic observations. *Compendium*. 1994; 15:932, 935–938. quiz 939. [PubMed: 7728821]
34. Fiorellini JP, Howell TH, Cochran D, Malmquist J, Lilly LC, Spagnoli D, Toljanic J, Jones A, Nevins M. Randomized study evaluating recombinant human bone morphogenetic protein-2 for extraction socket augmentation. *J Periodontol*. 2005; 76:605–613. [PubMed: 15857102]
35. Franceschi RT, Wang D, Krebsbach PH, Rutherford RB. Gene therapy for bone formation: in vitro and in vivo osteogenic activity of an adenovirus expressing BMP7. *J Cell Biochem*. 2000; 78:476–486. [PubMed: 10861845]
36. Francetti L, Del Fabbro M, Basso M, Testori T, Weinstein R. Enamel matrix proteins in the treatment of intra-bony defects. A prospective 24-month clinical trial. *J Clin Periodontol*. 2004; 31:52–59. [PubMed: 15058375]
37. Froum SJ, Thaler R, Scopp IW, Stahl SS. Osseous autografts. I. Clinical responses to bone blend or hip marrow grafts. *J Periodontol*. 1975; 46:515–521. [PubMed: 1099188]
38. Fuentes P, Garrett S, Nilveus R, Egelberg J. Treatment of periodontal furcation defects. Coronally positioned flap with or without citric acid root conditioning in class II defects. *J Clin Periodontol*. 1993; 20:425–430. [PubMed: 8349833]
39. Galgut PN, Waite IM, Brookshaw JD, Kingston CP. A 4-year controlled clinical study into the use of a ceramic hydroxylapatite implant material for the treatment of periodontal bone defects. *J Clin Periodontol*. 1992; 19:570–577. [PubMed: 1332987]
40. Gao Y, Yang L, Fang YR, Mori M, Kawahara K, Tanaka A. The inductive effect of bone morphogenetic protein (BMP) on human periodontal fibroblast-like cells in vitro. *J Osaka Dent Univ*. 1995; 29:9–17. [PubMed: 8935072]
41. Gestreluis S, Lyngstadaas SP, Hammarstrom L. Emdogain--periodontal regeneration based on biomimicry. *Clin Oral Investig*. 2000; 4:120–125.
42. Giannobile WV. Periodontal tissue engineering by growth factors. *Bone*. 1996; 19:23S–37S. [PubMed: 8830996]
43. Giannobile WV. What does the future hold for periodontal tissue engineering? *Int J Periodontics Restorative Dent*. 2002; 22:6–7. [PubMed: 11922219]
44. Giannobile WV, Finkelman RD, Lynch SE. Comparison of canine and non-human primate animal models for periodontal regenerative therapy: results following a single administration of PDGF/IGF-I. *J Periodontol*. 1994; 65:1158–1168. [PubMed: 7877089]
45. Giannobile WV, Hernandez RA, Finkelman RD, Ryan S, Kiritsy CP, D'Andrea M, Lynch SE. Comparative effects of platelet-derived growth factor-BB and insulin-like growth factor-I, individually and in combination, on periodontal regeneration in *Macaca fascicularis*. *J Periodontal Res*. 1996; 31:301–312. [PubMed: 8858534]
46. Giannobile WV, Lee CS, Tomala MP, Tejeda KM, Zhu Z. Platelet-derived growth factor (PDGF) gene delivery for application in periodontal tissue engineering. *J Periodontol*. 2001; 72:815–823. [PubMed: 11453245]
47. Giannobile WV, Ryan S, Shih MS, Su DL, Kaplan PL, Chan TC. Recombinant human osteogenic protein-1 (OP-1) stimulates periodontal wound healing in class III furcation defects. *J Periodontol*. 1998; 69:129–137. [PubMed: 9526911]
48. Goldman H, Cohen W. The infrabony pocket: classification and treatment. *J Periodontol*. 1958; 29:272–291.
49. Gu DL, Nguyen T, Gonzalez AM, Printz MA, Pierce GF, Sosnowski BA, Phillips ML, Chandler LA. Adenovirus encoding human platelet-derived growth factor-B delivered in collagen exhibits safety, biodistribution, and immunogenicity profiles favorable for clinical use. *Mol Ther*. 2004; 9:699–711. [PubMed: 15120331]
50. Gurinsky BS, Mills MP, Mellonig JT. Clinical evaluation of demineralized freeze-dried bone allograft and enamel matrix derivative versus enamel matrix derivative alone for the treatment of periodontal osseous defects in humans. *J Periodontol*. 2004; 75:1309–1318. [PubMed: 15562907]
51. Haase HR, Clarkson RW, Waters MJ, Bartold PM. Growth factor modulation of mitogenic responses and proteoglycan synthesis by human periodontal fibroblasts. *J Cell Physiol*. 1998; 174:353–361. [PubMed: 9462697]

52. Hammarstrom L. Enamel matrix, cementum development and regeneration. *J Clin Periodontol.* 1997; 24:658–668. [PubMed: 9310870]
53. Hartman GA, Arnold RM, Mills MP, Cochran DL, Mellonig JT. Clinical and histologic evaluation of anorganic bovine bone collagen with or without a collagen barrier. *Int J Periodontics Restorative Dent.* 2004; 24:127–135. [PubMed: 15119883]
54. Heldin P, Laurent TC, Heldin CH. Effect of growth factors on hyaluronan synthesis in cultured human fibroblasts. *Biochem J.* 1989; 258:919–922. [PubMed: 2543365]
55. Hijawi J, Mogford JE, Chandler LA, Cross KJ, Said H, Sosnowski BA, Mustoe TA. Platelet-derived growth factor B, but not fibroblast growth factor 2, plasmid DNA improves survival of ischemic myocutaneous flaps. *Arch Surg.* 2004; 139:142–147. [PubMed: 14769570]
56. Howell TH, Fiorellini JP, Paquette DW, Offenbacher S, Giannobile WV, Lynch SE. A phase I/II clinical trial to evaluate a combination of recombinant human platelet-derived growth factor-BB and recombinant human insulin-like growth factor-I in patients with periodontal disease. *J Periodontol.* 1997; 68:1186–1193. [PubMed: 9444594]
57. Huang KK, Shen C, Chiang CY, Hsieh YD, Fu E. Effects of bone morphogenetic protein-6 on periodontal wound healing in a fenestration defect of rats. *J Periodontal Res.* 2005; 40:1–10. [PubMed: 15613073]
58. Huang YC, Simmons C, Kaigler D, Rice KG, Mooney DJ. Bone regeneration in a rat cranial defect with delivery of PEI-condensed plasmid DNA encoding for bone morphogenetic protein-4 (BMP-4). *Gene Ther.* 2005; 12:418–426. [PubMed: 15647766]
59. Hutmacher DW, Teoh SH, Zein I, Ranawake M, Lau S. Tissue engineering research: the engineer's role. *Med Device Technol.* 2000; 11:33–39. [PubMed: 10915483]
60. Ito H, Koefoed M, Tiyyapatanaputi P, Gromov K, Goater JJ, Carmouche J, Zhang X, Rubery PT, Rabinowitz J, Samulski RJ, Nakamura T, Soballe K, O'Keefe RJ, Boyce BF, Schwarz EM. Remodeling of cortical bone allografts mediated by adherent rAAV-RANKL and VEGF gene therapy. *Nat Med.* 2005; 11:291–297. [PubMed: 15711561]
61. Jin Q, Anusaksathien O, Webb SA, Printz MA, Giannobile WV. Engineering of tooth-supporting structures by delivery of PDGF gene therapy vectors. *Mol Ther.* 2004; 9:519–526. [PubMed: 15093182]
62. Jin Q, Cirelli JA, Park CH, Sugai JV, Taba M, Kostenuik PJ, Giannobile WV. RANKL Inhibition Through Osteoprotegerin Blocks Bone Loss in Experimental Periodontitis. *J Periodontol.* 2007; 78:1300–1308. [PubMed: 17608585]
63. Jin QM, Anusaksathien O, Webb SA, Rutherford RB, Giannobile WV. Gene therapy of bone morphogenetic protein for periodontal tissue engineering. *J Periodontol.* 2003; 74:202–213. [PubMed: 12666709]
64. Jin QM, Zhao M, Economides AN, Somerman MJ, Giannobile WV. Noggin gene delivery inhibits cementoblast-induced mineralization. *Connect Tissue Res.* 2004; 45:50–59. [PubMed: 15203940]
65. Judith R, Nithya M, Rose C, Mandal AB. Application of a PDGF-containing novel gel for cutaneous wound healing. *Life Sci.*
66. Jung RE, Glauser R, Scharer P, Hammerle CH, Sailer HF, Weber FE. Effect of rhBMP-2 on guided bone regeneration in humans. *Clin Oral Implants Res.* 2003; 14:556–568. [PubMed: 12969359]
67. Jung RE, Zwahlen R, Weber FE, Molenberg A, van Lenthe GH, Hammerle CH. Evaluation of an in situ formed synthetic hydrogel as a biodegradable membrane for guided bone regeneration. *Clin Oral Implants Res.* 2006; 17:426–433. [PubMed: 16907774]
68. Kaigler D, Cirelli JA, Giannobile WV. Growth factor delivery for oral and periodontal tissue engineering. *Expert Opin Drug Deliv.* 2006; 3:647–662. [PubMed: 16948560]
69. Kaplan DR, Chao FC, Stiles CD, Antoniades HN, Scher CD. Platelet alpha granules contain a growth factor for fibroblasts. *Blood.* 1979; 53:1043–1052. [PubMed: 444648]
70. Karring T, Cortellini P. Regenerative therapy: furcation defects. *Periodontol 2000.* 1999; 19:115–137. [PubMed: 10321220]
71. Karring T, Isidor F, Nyman S, Lindhe J. New attachment formation on teeth with a reduced but healthy periodontal ligament. *J Clin Periodontol.* 1985; 12:51–60. [PubMed: 3855871]

72. Karring T, Nyman S, Lindhe J. Healing following implantation of periodontitis affected roots into bone tissue. *J Clin Periodontol.* 1980; 7:96–105. [PubMed: 6929795]
73. Karring T, Nyman S, Lindhe J, Sirirat M. Potentials for root resorption during periodontal wound healing. *J Clin Periodontol.* 1984; 11:41–52. [PubMed: 6363461]
74. Kimble KM, Eber RM, Soehren S, Shyr Y, Wang HL. Treatment of gingival recession using a collagen membrane with or without the use of demineralized freeze-dried bone allograft for space maintenance. *J Periodontol.* 2004; 75:210–220. [PubMed: 15068108]
75. Krebsbach PH, Gu K, Franceschi RT, Rutherford RB. Gene therapy-directed osteogenesis: BMP-7-transduced human fibroblasts form bone in vivo. *Hum Gene Ther.* 2000; 11:1201–1210. [PubMed: 10834621]
76. Lee JY, Musgrave D, Pelinkovic D, Fukushima K, Cummins J, Usas A, Robbins P, Fu FH, Huard J. Effect of bone morphogenetic protein-2-expressing muscle-derived cells on healing of critical-sized bone defects in mice. *J Bone Joint Surg Am.* 2001; 83-A:1032–1039. [PubMed: 11451972]
77. Lee JY, Peng H, Usas A, Musgrave D, Cummins J, Pelinkovic D, Jankowski R, Ziran B, Robbins P, Huard J. Enhancement of bone healing based on ex vivo gene therapy using human muscle-derived cells expressing bone morphogenetic protein 2. *Hum Gene Ther.* 2002; 13:1201–1211. [PubMed: 12133273]
78. Lieberman JR, Daluiski A, Stevenson S, Wu L, McAllister P, Lee YP, Kabo JM, Finerman GA, Berk AJ, Witte ON. The effect of regional gene therapy with bone morphogenetic protein-2-producing bone-marrow cells on the repair of segmental femoral defects in rats. *J Bone Joint Surg Am.* 1999; 81:905–917. [PubMed: 10428121]
79. Lin Z, Sugai JV, Jin Q, Chandler LA, Giannobile WV. Platelet-derived growth factor-B gene delivery sustains gingival fibroblast signal transduction. *J Periodontal Res.* 2008; 43:440–449. [PubMed: 18823454]
80. Lioubavina-Hack N, Karring T, Lynch SE, Lindhe J. Methyl cellulose gel obstructed bone formation by GBR: an experimental study in rats. *J Clin Periodontol.* 2005; 32:1247–1253. [PubMed: 16269002]
81. Listgarten MA, Rosenberg MM. Histological study of repair following new attachment procedures in human periodontal lesions. *J Periodontol.* 1979; 50:333–344. [PubMed: 381631]
82. Lutolf MP, Hubbell JA. Synthetic biomaterials as instructive extracellular microenvironments for morphogenesis in tissue engineering. *Nat Biotechnol.* 2005; 23:47–55. [PubMed: 15637621]
83. Lutolf MP, Weber FE, Schmoekel HG, Schense JC, Kohler T, Muller R, Hubbell JA. Repair of bone defects using synthetic mimetics of collagenous extracellular matrices. *Nat Biotechnol.* 2003; 21:513–518. [PubMed: 12704396]
84. Lynch SE, de Castilla GR, Williams RC, Kiritsy CP, Howell TH, Reddy MS, Antoniades HN. The effects of short-term application of a combination of platelet-derived and insulin-like growth factors on periodontal wound healing. *J Periodontol.* 1991; 62:458–467. [PubMed: 1920013]
85. Lynch SE, Williams RC, Polson AM, Howell TH, Reddy MS, Zappa UE, Antoniades HN. A combination of platelet-derived and insulin-like growth factors enhances periodontal regeneration. *J Clin Periodontol.* 1989; 16:545–548. [PubMed: 2778088]
86. Machtei EE, Schallhorn RG. Successful regeneration of mandibular Class II furcation defects: an evidence-based treatment approach. *Int J Periodontics Restorative Dent.* 1995; 15:146–167. [PubMed: 8593980]
87. Marcopoulou CE, Vavouraki HN, Dereka XE, Vrotsos IA. Proliferative effect of growth factors TGF-beta1, PDGF-BB and rhBMP-2 on human gingival fibroblasts and periodontal ligament cells. *J Int Acad Periodontol.* 2003; 5:63–70. [PubMed: 12887144]
88. Mariotti A. Efficacy of chemical root surface modifiers in the treatment of periodontal disease. A systematic review. *Ann Periodontol.* 2003; 8:205–226. [PubMed: 14971255]
89. Matsuda N, Lin WL, Kumar NM, Cho MI, Genco RJ. Mitogenic, chemotactic, and synthetic responses of rat periodontal ligament fibroblastic cells to polypeptide growth factors in vitro. *J Periodontol.* 1992; 63:515–525. [PubMed: 1625151]
90. McCulloch CA. Basic considerations in periodontal wound healing to achieve regeneration. *Periodontol 2000.* 1993; 1:16–25.

91. McGuire MK, Kao RT, Nevins M, Lynch SE. rhPDGF-BB promotes healing of periodontal defects: 24-month clinical and radiographic observations. *Int J Periodontics Restorative Dent*. 2006; 26:223–231. [PubMed: 16836164]
92. Meffert RM, Thomas JR, Hamilton KM, Brownstein CN. Hydroxylapatite as an alloplastic graft in the treatment of human periodontal osseous defects. *J Periodontol*. 1985; 56:63–73. [PubMed: 2984403]
93. Mellonig JT. Freeze-dried bone allografts in periodontal reconstructive surgery. *Dent Clin North Am*. 1991; 35:505–520. [PubMed: 1879574]
94. Mellonig JT. Human histologic evaluation of a bovine-derived bone xenograft in the treatment of periodontal osseous defects. *Int J Periodontics Restorative Dent*. 2000; 20:19–29. [PubMed: 11203545]
95. Minenna L, Herrero F, Sanz M, Trombelli L. Adjunctive effect of a polylactide/polyglycolide copolymer in the treatment of deep periodontal intra-osseous defects: a randomized clinical trial. *J Clin Periodontol*. 2005; 32:456–461. [PubMed: 15842259]
96. Moore JA, Ashley FP, Waterman CA. The effect on healing of the application of citric acid during replaced flap surgery. *J Clin Periodontol*. 1987; 14:130–135. [PubMed: 3549794]
97. Moses O, Pitaru S, Artzi Z, Nemcovsky CE. Healing of dehiscence-type defects in implants placed together with different barrier membranes: a comparative clinical study. *Clin Oral Implants Res*. 2005; 16:210–219. [PubMed: 15777331]
98. Murakami S, Takayama S, Kitamura M, Shimabukuro Y, Yanagi K, Ikezawa K, Saho T, Nozaki T, Okada H. Recombinant human basic fibroblast growth factor (bFGF) stimulates periodontal regeneration in class II furcation defects created in beagle dogs. *J Periodontal Res*. 2003; 38:97–103. [PubMed: 12558943]
99. Murphy KG, Gunsolley JC. Guided tissue regeneration for the treatment of periodontal intrabony and furcation defects. A systematic review. *Ann Periodontol*. 2003; 8:266–302. [PubMed: 14971257]
100. Musgrave DS, Bosch P, Ghivizzani S, Robbins PD, Evans CH, Huard J. Adenovirus-mediated direct gene therapy with bone morphogenetic protein-2 produces bone. *Bone*. 1999; 24:541–547. [PubMed: 10375195]
101. Nakashima M, Reddi AH. The application of bone morphogenetic proteins to dental tissue engineering. *Nat Biotechnol*. 2003; 21:1025–1032. [PubMed: 12949568]
102. Needleman I, Tucker R, Giedrys-Leeper E, Worthington H. A systematic review of guided tissue regeneration for periodontal infrabony defects. *J Periodontal Res*. 2002; 37:380–388. [PubMed: 12366862]
103. Needleman IG, Worthington HV, Giedrys-Leeper E, Tucker RJ. Guided tissue regeneration for periodontal infra-bony defects. *Cochrane Database Syst Rev*. 2006 CD001724.
104. Nery EB, LeGeros RZ, Lynch KL, Lee K. Tissue response to biphasic calcium phosphate ceramic with different ratios of HA/beta TCP in periodontal osseous defects. *J Periodontol*. 1992; 63:729–735. [PubMed: 1335498]
105. Nevins M, Camelo M, Nevins ML, Schenk RK, Lynch SE. Periodontal regeneration in humans using recombinant human platelet-derived growth factor-BB (rhPDGF-BB) and allogenic bone. *J Periodontol*. 2003; 74:1282–1292. [PubMed: 14584860]
106. Nevins M, Giannobile WV, McGuire MK, Kao RT, Mellonig JT, Hinrichs JE, McAllister BS, Murphy KS, McClain PK, Nevins ML, Paquette DW, Han TJ, Reddy MS, Lavin PT, Genco RJ, Lynch SE. Platelet-derived growth factor stimulates bone fill and rate of attachment level gain: results of a large multicenter randomized controlled trial. *J Periodontol*. 2005; 76:2205–2215. [PubMed: 16332231]
107. Nevins ML, Camelo M, Nevins M, King CJ, Oringer RJ, Schenk RK, Fiorellini JP. Human histologic evaluation of bioactive ceramic in the treatment of periodontal osseous defects. *Int J Periodontics Restorative Dent*. 2000; 20:458–467. [PubMed: 11203583]
108. Nishimura F, Terranova VP. Comparative study of the chemotactic responses of periodontal ligament cells and gingival fibroblasts to polypeptide growth factors. *J Dent Res*. 1996; 75:986–992. [PubMed: 8708140]

109. Nyman S, Karring T, Lindhe J, Planten S. Healing following implantation of periodontitis-affected roots into gingival connective tissue. *J Clin Periodontol.* 1980; 7:394–401. [PubMed: 6936413]
110. Oates TW, Rouse CA, Cochran DL. Mitogenic effects of growth factors on human periodontal ligament cells in vitro. *J Periodontol.* 1993; 64:142–148. [PubMed: 8433255]
111. Oberyshyn TM. Inflammation and wound healing. *Front Biosci.* 2007; 12:2993–2999. [PubMed: 17485276]
112. Ojima Y, Mizuno M, Kuboki Y, Komori T. In vitro effect of platelet-derived growth factor-BB on collagen synthesis and proliferation of human periodontal ligament cells. *Oral Dis.* 2003; 9:144–151. [PubMed: 12945596]
113. Owczarek B, Kiernicka M, Galkowska E, Wysokinska-Miszczuk J. The application of Bio-Oss and Bio-Gide as implant materials in the complex treatment of aggressive periodontitis. *Ann Univ Mariae Curie Sklodowska [Med].* 2003; 58:392–396.
114. Palti A, Hoch T. A concept for the treatment of various dental bone defects. *Implant Dent.* 2002; 11:73–78. [PubMed: 11915549]
115. Parashis A, Andronikaki-Faldami A, Tsiklakis K. Clinical and radiographic comparison of three regenerative procedures in the treatment of intrabony defects. *Int J Periodontics Restorative Dent.* 2004; 24:81–90. [PubMed: 14984149]
116. Parkar MH, Kuru L, Giouzei M, Olsen I. Expression of growth-factor receptors in normal and regenerating human periodontal cells. *Arch Oral Biol.* 2001; 46:275–284. [PubMed: 11165574]
117. Peng H, Wright V, Usas A, Gearhart B, Shen HC, Cummins J, Huard J. Synergistic enhancement of bone formation and healing by stem cell-expressed VEGF and bone morphogenetic protein-4. *J Clin Invest.* 2002; 110:751–759. [PubMed: 12235106]
118. Pini Prato G, Clauser C, Cortellini P, Tinti C, Vincenzi G, Pagliaro U. Guided tissue regeneration versus mucogingival surgery in the treatment of human buccal recessions. A 4-year follow-up study. *J Periodontol.* 1996; 67:1216–1223. [PubMed: 8959573]
119. Polimeni G, Koo KT, Qahash M, Xiropaidis AV, Albandar JM, Wikesjo UM. Prognostic factors for alveolar regeneration: effect of a space-providing biomaterial on guided tissue regeneration. *J Clin Periodontol.* 2004; 31:725–729. [PubMed: 15312093]
120. Polimeni G, Xiropaidis AV, Wikesjo UM. Biology and principles of periodontal wound healing/regeneration. *Periodontol 2000.* 2006; 41:30–47. [PubMed: 16686925]
121. Prichard J. Regeneration of bone following periodontal therapy; report of cases. *Oral Surg Oral Med Oral Pathol.* 1957; 10:247–252. [PubMed: 13407919]
122. Printz MA, Gonzalez AM, Cunningham M, Gu DL, Ong M, Pierce GF, Aukerman SL. Fibroblast growth factor 2-retargeted adenoviral vectors exhibit a modified biolocalization pattern and display reduced toxicity relative to native adenoviral vectors. *Hum Gene Ther.* 2000; 11:191–204. [PubMed: 10646650]
123. Raja S, Byakod G, Pudakalkatti P. Growth factors in periodontal regeneration. *Int J Dent Hyg.* 2009; 7:82–89. [PubMed: 19413545]
124. Ramamurthy NS, Greenwald RA, Celiker MY, Shi EY. Experimental arthritis in rats induces biomarkers of periodontitis which are ameliorated by gene therapy with tissue inhibitor of matrix metalloproteinases. *J Periodontol.* 2005; 76:229–233. [PubMed: 15974846]
125. Ramseier CA, Abramson ZR, Jin Q, Giannobile WV. Gene therapeutics for periodontal regenerative medicine. *Dent Clin North Am.* 2006; 50:245–263. ix. [PubMed: 16530061]
126. Rasperini G, Silvestri M, Ricci G. Long-term clinical observation of treatment of infrabony defects with enamel matrix derivative (Emdogain): surgical reentry. *Int J Periodontics Restorative Dent.* 2005; 25:121–127. [PubMed: 15839588]
127. Reddi AH. Bone morphogenetic proteins: from basic science to clinical applications. *J Bone Joint Surg Am.* 2001; 83-A(Suppl 1):S1–S6. [PubMed: 11263660]
128. Reddi AH. Role of morphogenetic proteins in skeletal tissue engineering and regeneration. *Nat Biotechnol.* 1998; 16:247–252. [PubMed: 9528003]
129. Reddy MS, Jeffcoat MK. Methods of assessing periodontal regeneration. *Periodontol 2000.* 1999; 19:87–103. [PubMed: 10321218]

130. Renvert S, Garrett S, Shallhorn RG, Egelberg J. Healing after treatment of periodontal intraosseous defects. III. Effect of osseous grafting and citric acid conditioning. *J Clin Periodontol.* 1985; 12:441–455. [PubMed: 3894434]
131. Reynolds MA, Aichelmann-Reidy ME, Branch-Mays GL, Gunsolley JC. The efficacy of bone replacement grafts in the treatment of periodontal osseous defects. A systematic review. *Ann Periodontol.* 2003; 8:227–265. [PubMed: 14971256]
132. Richardson CR, Mellonig JT, Brunsvold MA, McDonnell HT, Cochran DL. Clinical evaluation of Bio-Oss: a bovine-derived xenograft for the treatment of periodontal osseous defects in humans. *J Clin Periodontol.* 1999; 26:421–428. [PubMed: 10412845]
133. Richardson TP, Peters MC, Ennett AB, Mooney DJ. Polymeric system for dual growth factor delivery. *Nat Biotechnol.* 2001; 19:1029–1034. [PubMed: 11689847]
134. Rosenkranz S, Kazlauskas A. Evidence for distinct signaling properties and biological responses induced by the PDGF receptor alpha and beta subtypes. *Growth Factors.* 1999; 16:201–216. [PubMed: 10372961]
135. Rosing CK, Aass AM, Mavropoulos A, Gjermo P. Clinical and radiographic effects of enamel matrix derivative in the treatment of intrabony periodontal defects: a 12-month longitudinal placebo-controlled clinical trial in adult periodontitis patients. *J Periodontol.* 2005; 76:129–133. [PubMed: 15830647]
136. Rutherford RB, Niekrah CE, Kennedy JE, Charette MF. Platelet-derived and insulin-like growth factors stimulate regeneration of periodontal attachment in monkeys. *J Periodontal Res.* 1992; 27:285–290. [PubMed: 1640350]
137. Rutherford RB, Sampath TK, Rueger DC, Taylor TD. Use of bovine osteogenic protein to promote rapid osseointegration of endosseous dental implants. *Int J Oral Maxillofac Implants.* 1992; 7:297–301. [PubMed: 1289254]
138. Sanz M, Tonetti MS, Zabalegui I, Sicilia A, Blanco J, Rebelo H, Rasperini G, Merli M, Cortellini P, Suvan JE. Treatment of intrabony defects with enamel matrix proteins or barrier membranes: results from a multicenter practice-based clinical trial. *J Periodontol.* 2004; 75:726–733. [PubMed: 15212355]
139. Scher EL, Day RB, Speight PM. New bone formation after a sinus lift procedure using demineralized freeze-dried bone and tricalcium phosphate. *Implant Dent.* 1999; 8:49–53. [PubMed: 10356456]
140. Sculean A, Chiantella GC, Windisch P, Arweiler NB, Brex M, Gera I. Healing of intra-bony defects following treatment with a composite bovine-derived xenograft (Bio-Oss Collagen) in combination with a collagen membrane (Bio-Gide PERIO). *J Clin Periodontol.* 2005; 32:720–724. [PubMed: 15966877]
141. Sculean A, Nikolidakis D, Schwarz F. Regeneration of periodontal tissues: combinations of barrier membranes and grafting materials - biological foundation and preclinical evidence: a systematic review. *J Clin Periodontol.* 2008; 35:106–116. [PubMed: 18724845]
142. Sculean A, Pietruska M, Schwarz F, Willershausen B, Arweiler NB, Auschill TM. Healing of human intrabony defects following regenerative periodontal therapy with an enamel matrix protein derivative alone or combined with a bioactive glass. A controlled clinical study. *J Clin Periodontol.* 2005; 32:111–117. [PubMed: 15642068]
143. Sculean A, Schwarz F, Becker J, Brex M. The application of an enamel matrix protein derivative (Emdogain) in regenerative periodontal therapy: a review. *Med Princ Pract.* 2007; 16:167–180. [PubMed: 17409750]
144. Seppa H, Grotendorst G, Seppa S, Schiffmann E, Martin GR. Platelet-derived growth factor in chemotactic for fibroblasts. *J Cell Biol.* 1982; 92:584–588. [PubMed: 7061598]
145. Sigurdsson TJ, Lee MB, Kubota K, Turek TJ, Wozney JM, Wikesjo UM. Periodontal repair in dogs: recombinant human bone morphogenetic protein-2 significantly enhances periodontal regeneration. *J Periodontol.* 1995; 66:131–138. [PubMed: 7730964]
146. Singer AJ, Clark RA. Cutaneous wound healing. *N Engl J Med.* 1999; 341:738–746. [PubMed: 10471461]

147. Sorensen RG, Wikesjo UM, Kinoshita A, Wozney JM. Periodontal repair in dogs: evaluation of a bioresorbable calcium phosphate cement (Ceredex) as a carrier for rhBMP-2. *J Clin Periodontol.* 2004; 31:796–804. [PubMed: 15312104]
148. Stavropoulos A, Sculean A, Karring T. GTR treatment of intrabony defects with PLA/PGA copolymer or collagen bioresorbable membranes in combination with deproteinized bovine bone (Bio-Oss). *Clin Oral Investig.* 2004; 8:226–232.
149. Sykaras N, Opperman LA. Bone morphogenetic proteins (BMPs): how do they function and what can they offer the clinician? *J Oral Sci.* 2003; 45:57–73. [PubMed: 12930129]
150. Taba, M., Jr; Huffer, HH.; Shelburne, CE.; Kriegl, JM.; Goldstein, SA.; Lustig, KH.; Burnstein, H.; Giannobile, WV. Gene delivery of TNFR:Fc by adeno-associated virus vector blocks progression of periodontitis [abstract 677]; *Molecular Therapy, Abstracts from the American Society of Gene Therapy's 8th Annual Meeting*; Saint Louis, MO. 2005. p. 262
151. Takayama S, Murakami S, Shimabukuro Y, Kitamura M, Okada H. Periodontal regeneration by FGF-2 (bFGF) in primate models. *J Dent Res.* 2001; 80:2075–2079. [PubMed: 11808765]
152. Terranova VP, Odziemiec C, Tweden KS, Spadone DP. Repopulation of dentin surfaces by periodontal ligament cells and endothelial cells. Effect of basic fibroblast growth factor. *J Periodontol.* 1989; 60:293–301. [PubMed: 2550605]
153. Tonetti MS, Lang NP, Cortellini P, Suvan JE, Adriaens P, Dubravec D, Fonzar A, Fourmousis I, Mayfield L, Rossi R, Silvestri M, Tiedemann C, Topoll H, Vangsted T, Wallkamm B. Enamel matrix proteins in the regenerative therapy of deep intrabony defects. *J Clin Periodontol.* 2002; 29:317–325. [PubMed: 11966929]
154. Trejo PM, Weltman R, Caffesse R. Treatment of intraosseous defects with bioabsorbable barriers alone or in combination with decalcified freeze-dried bone allograft: a randomized clinical trial. *J Periodontol.* 2000; 71:1852–1861. [PubMed: 11156042]
155. Trombelli L, Farina R. Clinical outcomes with bioactive agents alone or in combination with grafting or guided tissue regeneration. *J Clin Periodontol.* 2008; 35:117–135. [PubMed: 18724846]
156. Trombelli L, Heitz-Mayfield LJ, Needleman I, Moles D, Scabbia A. A systematic review of graft materials and biological agents for periodontal intraosseous defects. *J Clin Periodontol.* 2002; 29(Suppl 3):117–135. discussion 160–112. [PubMed: 12787213]
157. Trombelli L, Minenna L, Farina R, Scabbia A. Guided tissue regeneration in human gingival recessions. A 10-year follow-up study. *J Clin Periodontol.* 2005; 32:16–20. [PubMed: 15642053]
158. Urist MR. Bone: formation by autoinduction. *Science.* 1965; 150:893–899. [PubMed: 5319761]
159. van den Bergh JP, ten Bruggenkate CM, Groeneveld HH, Burger EH, Tuinzing DB. Recombinant human bone morphogenetic protein-7 in maxillary sinus floor elevation surgery in 3 patients compared to autogenous bone grafts. A clinical pilot study. *J Clin Periodontol.* 2000; 27:627–636. [PubMed: 10983596]
160. Wang HL, Greenwell H, Fiorellini J, Giannobile W, Offenbacher S, Salkin L, Townsend C, Sheridan P, Genco RJ. Periodontal regeneration. *J Periodontol.* 2005; 76:1601–1622. [PubMed: 16171453]
161. Wikesjo UM, Lim WH, Thomson RC, Cook AD, Wozney JM, Hardwick WR. Periodontal repair in dogs: evaluation of a bioabsorbable space-providing macroporous membrane with recombinant human bone morphogenetic protein-2. *J Periodontol.* 2003; 74:635–647. [PubMed: 12816296]
162. Wikesjo UM, Qahash M, Thomson RC, Cook AD, Rohrer MD, Wozney JM, Hardwick WR. rhBMP-2 significantly enhances guided bone regeneration. *Clin Oral Implants Res.* 2004; 15:194–204. [PubMed: 15008931]
163. Wikesjo UM, Sorensen RG, Kinoshita A, Jian Li X, Wozney JM. Periodontal repair in dogs: effect of recombinant human bone morphogenetic protein-12 (rhBMP-12) on regeneration of alveolar bone and periodontal attachment. *J Clin Periodontol.* 2004; 31:662–670. [PubMed: 15257745]
164. Wikesjo UM, Xiropaidis AV, Thomson RC, Cook AD, Selvig KA, Hardwick WR. Periodontal repair in dogs: rhBMP-2 significantly enhances bone formation under provisions for guided tissue regeneration. *J Clin Periodontol.* 2003; 30:705–714. [PubMed: 12887339]

165. Wolberg AS. Thrombin generation and fibrin clot structure. *Blood Rev.* 2007; 21:131–142. [PubMed: 17208341]
166. Wozney JM. The potential role of bone morphogenetic proteins in periodontal reconstruction. *J Periodontol.* 1995; 66:506–510. [PubMed: 7562340]
167. Wozney JM, Rosen V, Celeste AJ, Mitsock LM, Whitters MJ, Kriz RW, Hewick RM, Wang EA. Novel regulators of bone formation: molecular clones and activities. *Science.* 1988; 242:1528–1534. [PubMed: 3201241]
168. Yang L, Zhang Y, Dong R, Peng L, Liu X, Wang Y, Cheng X. Effects of adenoviral-mediated coexpression of bone morphogenetic protein-7 and insulin-like growth factor-1 on human periodontal ligament cells. *J Periodontal Res.*
169. Yeo YJ, Jeon DW, Kim CS, Choi SH, Cho KS, Lee YK, Kim CK. Effects of chitosan nonwoven membrane on periodontal healing of surgically created one-wall intrabony defects in beagle dogs. *J Biomed Mater Res B Appl Biomater.* 2005; 72:86–93. [PubMed: 15389496]
170. Zaman KU, Sugaya T, Kato H. Effect of recombinant human platelet-derived growth factor-BB and bone morphogenetic protein-2 application to demineralized dentin on early periodontal ligament cell response. *J Periodontal Res.* 1999; 34:244–250. [PubMed: 10567947]
171. Zhu Z, Lee CS, Tejeda KM, Giannobile WV. Gene transfer and expression of platelet-derived growth factors modulate periodontal cellular activity. *J Dent Res.* 2001; 80:892–897. [PubMed: 11379891]

Fig. 1.
Periodontal wound following flap surgery: 1) gingival epithelium, 2) gingival connective tissue, 3) periodontal ligament, 4) alveolar bone and 5) cementum or dentin on the dental root surface

Fig. 2.

A) Regular healing process following the periodontal flap adaptation with significant reduction of the attachment apparatus.

B) In order to enable and promote the healing towards the rebuilding of cementum and periodontal ligament, the gingival epithelium must be prevented from forming a long junctional epithelium along the root surface down to the former level of the periodontal ligament (e.g. by placement of a bioresorbable membrane).

Fig. 3.

Periodontal regeneration of a 3 wall infrabony defect using Emdogain

A: 32 year old patient, male, non smoker with severe periodontitis. Tooth 13 shows a probing pocket depth of 10 mm disto-buccally and clinical attachment loss of 14 mm.

B: Pre-treatment radiograph shows the infra-bony defect distal to the tooth 13.

C: After the buccal incision of the papilla, the interdental tissue is preserved attached to the palatal flap. After the debridement of the granulation tissue and the root surface debridement, the infra-bony defect is classified and measured: the predominant component is a 7 mm deep 3-wall defect.

D: 1-year after surgical intervention the distal site of the tooth #13 shows a probing pocket depth of 2 mm and clinical attachment loss of 7 mm. If compared with the initial measurements a probing pocket depth gain of 8 mm and a clinical attachment loss gain of 7 mm has been achieved.

E: Radiograph 1 year post-surgery showing filling of the defect.

Fig. 4.

Periodontal regeneration using platelet-derived growth factor and bone graft materials

A: 27-year old patient at the reevaluation visit after the initial non-surgical therapy presented 3 sites with probing pocket depth > 6 mm, one of those distal to the tooth #44 shows a probing pocket depth of 7 mm and no gingival recession.

B: The periapical radiograph shows a deep 1-wall defect distal to tooth #44 and a lesion between the teeth #45 and 46.

C: The measurement of the 1-wall defect shows an infra-bony component of 6 mm

D: The Grafting material (GEM 21S®) is mixed with particles of autogenous bone chips collected in the surgical area with a Rhodes instrument and with the liquid component of the GEM 21S® (platelet-derived growth factor).

E: The liquid platelet-derived growth factor is placed in the defect together with the graft to rebuild the lost bone.

F: A second internal mattress suture is performed with a 7- 0 Gore Tex® suture, to allow for optimal adaptation of the flap margin without the interference of the epithelium. The two internal mattress sutures are tied and the knots are performed only after a perfect free tension closure of the wound. Two additional interrupted 7-0 sutures are placed to ensure stable contact between the connective tissues of the edges of the flaps. The mesial and distal papillae are stabilized with additional simple interrupted sutures

G: 9 months after surgery, the probing pocket depth is 2 mm.

H: 9 months after surgery, the periapical radiograph shows a good bone fill of the 1 wall bony defect.

I: 9 months after surgery, the surgical re-entry shows the new bone formation

Fig. 5.

Advanced approaches for regenerating tooth-supporting structures.

A) Application of a graft material (e.g. bone ceramic) and growth factor into an infra-bony defect covered by a bioresorbable membrane.

B) Application of gene vectors for the transduction of growth factors producing target cells.

Table 1

Currently available regenerative biomaterials in periodontology

Regenerative biomaterials	Trade name(s)	References
Bone autogenous grafts (Autografts)		
Intraoral autografts	n/a	Renvert et al. 1985 (130) Ellegaard & L��e 1971 (29)
Extraoral autografts	n/a	Froum et al. 1975 (37)
Bone allogenic grafts (Allografts)		
Freeze dried bone allograft	Grafton��, Lifenet��	Mellonig et al. 1991 (93)
Demineralized freeze dried bone allograft	Transplant Foundation��	Gurinsky et al. 2004 (50) Kimble et al. 2004 (74) Trejo et al. 2000 (154)
Bone xenogenic grafts (Xenografts)		
Bovine mineral matrix	Bio-Oss, OsteoGraf��, Pep-Gen P-15��	Hartman et al. 2004 (53) Camelo et al. 2001 (12) Mellonig et al. 2000 (94) Nevins et al. 2000 (107) Richardson et al. 1999 (132)
Bone alloplastic grafts (Alloplasts)		
Hydroxyapatite (dense, porous, resorbable)	Osteogen��, Periograft��, ProOsteone��	Meffert et al. 1985 (92) Galgut et al. 1992 (39)
Beta Tricalcium phosphate	Sunthgraft��, alpha-BSM��	Palti et al. 2002 (114) Scher et al. 1999 (139) Nery et al. 1992 (104)
Hard tissue replacement polymers	Bioplant��	Dryankova et al. 2001 (27)
Bioactive glass (SiO ₂ , CaO, Na ₂ O, P ₂ O ₅)	PerioGlas��, BioGran��	Sculean et al. 2005 (142) Reynolds et al. 2003 (131) Trombelli et al. 2002 (156) Fetner et al. 1994 (33)
Coral-derived calcium carbonate	Biocoral��	Polimeni et al. 2004 (119)
Polymer and Collagen sponges		
Collagen	Helistat��, Collacote��, Colla-Tec��, Gelfoam��	
Poly lactide-copolyglycolide barrier membranes		
Methylcellulose	n/a	Lioubavina-Hack et al. 2005 (80)
Hyaluronic acid ester	Hy��	Wikesj�� et al. 2003 (161)
Chitosan	n/a	Yeo et al. 2005 (169)
Synthetic hydrogel		
Polyethylene glycol	n/a	Jung et al. 2007 (67)
Non-resorbable cell occlusive barrier membranes		
Polytetrafluorethylene	Gore-Tex��	Trombelli et al. 2005 (157) Moses et al. 2005 (97) Murphy et al. 2003 (99) Needleman et al. 2002 (102)
Resorbable cell occlusive barrier membranes		
Polyglycolide/Poly lactide (synthetic)	Ossix��	Minenna et al. 2005 (95) Stavropoulos et al. 2004 (148) Parashis et al. 2003 (115)
Collagen membrane (xenogen)	Bio-Gide��	Sculean et al. 2005 (140) Owczarek et al. 2003 (113) Camelo et al. 1998 (14)
Growth Factors		

Regenerative biomaterials	Trade name(s)	References
Enamel matrix derivative	Emdogain®	Rasperini et al. 2005 (126) Rosing et al. 2005 (135) Sanz et al. 2004 (138) Francetti et al. 2004 (36) Tonetti et al. 2002 (153) Esposito et al. 2003 (31) Esposito et al. 2004 (30) Esposito et al. 2005 (32)
Platelet-derived growth factor	Gem 21S®	Nevins et al. 2005 (106)
Bone morphogenetic protein	Infuse®	Fiorellini et al. 2005 (34)

Table 2

Effects of growth factors used for periodontal tissue engineering

Growth Factor	Effects
Platelet-derived growth factor	Migration, proliferation and non-collagenous matrix synthesis of mesenchymal cells
Bone morphogenetic protein	Proliferation, differentiation of osteoblasts, differentiation of periodontal ligament cells into osteoblasts
Enamel matrix derivative	Proliferation, protein synthesis and mineral nodule formation in periodontal ligament cells, osteoblasts and cementoblasts
Transforming growth factor Beta	Proliferation of cementoblasts and periodontal ligament fibroblasts
Insulin-like growth factor-1	Cell migration, proliferation, differentiation and matrix synthesis
Fibroblast growth factor-2	Proliferation and attachment of endothelial cells and periodontal ligament cells