Published in final edited form as:

European J Org Chem. 2011 December ; 2011(34): 6843–6846. doi:10.1002/ejoc.201101235.

Alkenenitrile Transmissive Olefination: Synthesis of the Putative Lignan "Morinol I"

Fraser F. Fleming^{a,*}, Wang Liu^b, Lihua Yao^a, Bhaskar Pitta^a, Matthew Purzycki^a, and P. C. Ravikumar^c

^aDepartment of Chemistry and Biochemistry, Duquesne University, Pittsburgh, PA 15282-1530, USA

^bDepartment of Chemistry, Ohio State University, Ohio, USA.

Department of Chemistry, Indian Institute of Technology, Mandi, India.

Abstract

Grignard reagents trigger an addition-elimination with α' -hydroxy acrylonitriles to selectively generate *Z*-alkenenitriles. The modular assembly of *Z*-alkenenitriles from a Grignard reagent, acrylonitrile, and an aldehyde is ideal for stereoselectively synthesizing alkenes as illustrated in the synthesis of the putative lignan "morinol I."

Keywords

Nitrile; Rearrangement; Olefination; Allylic compounds

Introduction

Six-electron transition structures undergird many extremely powerful synthetic methods.^[1] Diels-Alder cycloadditions, electrocyclic reactions, and [3,3]-sigmatropic rearrangements all proceed through privileged six-electron, six-atom transition structures. A remarkable diversity of atom substitutions are tolerated within the six-membered ring; Cope, Claisen, aza-Cope reactions incorporate carbon, oxygen, and nitrogen ring atoms, respectively. Equally powerful are heteroatom-metal combinations proceeding through six-membered Zimmerman-Traxler transition structures that efficiently and stereoselectively allow aldol condensations, allylations, and intramolecular hydride reductions.^[2]

Harnessing the facility of six-electron reactions to drive the formal displacement of hydroxide is rare – only high-valent titanium and cerium promote the elimination.^[3] Typically, allylic alcohols require activation prior to displacement because hydroxide is a poor leaving group.^[4] Conceptually, hydroxide ejection from hydroxyalkenenitrile **1** (Scheme 1) could be promoted through a concerted, six-electron, anionic rearrangement (**2** → **3**) and by forming a particularly stable metal oxide. Preferential rearrangement through conformer **2''** would assemble the α -substituted *Z*-alkenenitrile **3** whose stereoselective synthesis is challenging.^[4a,5] The value of alkenenitriles lies in their versatility as synthetic intermediates^[6] and as potent pharmacophores embedded within numerous bioactive natural products^[7] and pharmaceuticals.^[8] Generally only one *E/Z* stereoisomer is bioactive, or at

Fax: +1-412-396-5683, flemingf@duq.edu.

Supporting information for this article is available on the WWW under <http://www.eurjoc.org/> or from the author.

least significantly more active, underscoring the need for rapid, modular, stereoselective syntheses of alkenenitriles.^[9]

Results and Discussion

Exploratory forays focused on a magnesium oxide-driven rearrangement because of the strength of the Mg=O bond,^[10] the prevalence of commercial Grignards, and ready access to activated allylic alcohols through Baylis-Hillman condensations.^[11] After some optimization of temperature and stoichiometry, the addition of excess *i*-PrMgCl to hydroxyalkenenitrile **1a** was found to readily afford the *Z*-alkenenitrile **3a** as a single diastereomer^[12] (Table 1, entry 1).

The transmissive olefination smoothly generates an array of *Z*-alkenenitriles **3** from the corresponding hydroxyalkenenitriles **1** (Table 1). The extremely compact nitrile group resists nucleophilic attack by the proximal Grignard reagent, instead channeling a stereoselective^[12] alkyl addition-elimination to the alkenenitrile **3**. For example, addition of BuMgCl to nitrile **1a** forms the perfumery agent **3c** as a single stereoisomer (Table 1, entry 3).^[13] Sterically demanding, secondary, and tertiary Grignard reagents add with efficacies comparable to those of primary Grignards (Table 1, compare entries 2 and 3 with entries 1, 4, and 5). Similarly, no difference in efficiency is observed between electron rich and electron deficient aryl hydroxynitriles (Table 1, compare entries 3, 10, and 11). Although acetylenic Grignards do not engage in the rearrangement, vinyl magnesium halides effectively generate skipped dienes without detectable olefin isomerization (Table 1, entries 6 and 9).

Modest oxygenation within the Grignard reagent is tolerated (Table 1, entry 7), although attempts to use (1,3-dioxan-2-ylethyl)magnesium bromide were unsuccessful. The transmissive olefination with (3-methoxyphenethyl)magnesium bromide employed one equivalent of *i*-PrMgBr as a sacrificial base (Table 1, entry 7), reducing the amount of the Grignard reagent required for alkyl transfer to two equivalents. Using this procedure the alkenenitrile **3g** was prepared in 62% yield.

A carbinol substituent capable of stabilizing charge is required for the rearrangement.^[14] Presumably the sp² center promotes the rearrangement by aligning a p-orbital with the C-O bond undergoing cleavage. In addition to benzylic alcohols (Table 1, entries 1- 11), the cinnamyl alcohol **1d** readily affords conjugated dienenitriles^[9,15] **3l-3n** (Table 1, entries 12-14).

The addition-elimination requires 3.5 equivalents of Grignard reagent suggesting rearrangement via the more nucleophilic magnesiate **4** (Scheme 2) rather than an alkylmagnesium alkoxide (cf. **2** Scheme 1).^[16] Consistent with the intermediacy of a magnesiate, no rearrangement is observed on treating **1a** with one equivalent of dibutylmagnesium. Magnesium cations appear essential as the sequential addition of an equivalent of RMgX followed excess RLi affords only trace amounts of alkenenitrile **3**. Formation of the *Z*-alkenenitrile **3** is consistent with a concerted,¹⁷ exocyclic intramolecular rearrangement through the magnesiate **4''** in which the sterically miniscule nitrile group^[18] eclipses the carbinol substituent R¹. The model also accounts for the higher selectivity of aryl over alkenyl substituents R¹ assuming that the latter orients^[14] to position a proton toward the eclipsing inside substituent (H* in **4'** and CN in **4''**).

Rapid access to *Z*-alkenenitriles stimulated applying the transmissive olefination to the synthesis of the neolignan morinol I. Isolated during a bioactivity guided isolation of the Chinese medicinal herb *Morina chinensis*,^[19] morinol I possesses a skipped diene unique

among this class of neolignan metabolites (**6a**, Scheme 3).^[20] Vinylmagnesium bromide-initiated transmissive olefination with the veratraldehyde-derived nitrile **1b** effectively provided the requisite *Z*-dienenitrile **3o** as a single stereoisomer.^[11b] Subsequent exposure of **3o** to 3, 4-dimethoxyboronic acid in the presence of Pd(OAc)₂ and Ag₂CO₃ affords dienenitrile **5** with full control over the olefin stereochemistry (Scheme 3). Conventional *i*-Bu₂AlH reduction of nitrile **5**,^[21] hydrolysis, and aldehyde reduction afforded an 8:1 ratio of *E/Z* enal diastereomers that were reduced to the target alcohol (*7Z*, *7'E*)-**6a** and diastereomer (*7E*, *7'E*)-**6b**. Neither alcohol (*7Z*, *7'E*)-**6a** nor the C7-C8 *E*-diastereomer (*7E*, *7'E*)-**6b** exhibit spectral data matching the reported spectral listing of the natural product. Unfortunately samples and copies of the spectral data exhibited by the natural lignin are not available to resolve the ambiguous HMBC and NOE data that are inconsistent with the skipped diene structure proposed for morinol I.^[22] Although the precise structure of morinol I is elusive, the five-step synthesis of the putative structure of morinol I demonstrates the ability of the transmissive olefination strategy to rapidly assemble this type of molecular scaffold.

Conclusions

The transmissive olefination of hydroxy alkenenitriles efficiently and stereoselectively assembles *Z*-alkenenitriles from three readily available components: an aldehyde, acrylonitrile, and a Grignard reagent. The strategy directly employs Baylis-Hillman alcohols without prior hydroxyl activation by harnessing a unique elimination of magnesium oxide. Simply adding excess Grignard reagent to an α' -hydroxyalkenenitrile triggers a transmissive olefination to efficiently provide a diverse array of *Z*-alkenenitriles. The versatility of the transmissive olefination is illustrated in the rapid synthesis of the structure proposed for the lignan morinol I.

Experimental Section

General Transmissive Olefination Procedure

An ethereal solution of the Grignard reagent (3.5 equiv) was slowly added to a stirred 0 °C, THF solution of **1**. After 50 min the mixture was quenched with 0.1 M HCl solution and then the resulting mixture was stirred vigorously for 5 min. The crude product was extracted with ether, dried (MgSO₄), concentrated, and purified by radial chromatography to afford analytically pure material.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support from the National Institutes of Health (2R15AI051352-03) and in part from the National Science Foundation (CHE 0808996; NMR: 0614785, X-ray: 024872, HRMS: 0421252) is gratefully acknowledged.

References

1. Yang, J. Six-Membered Transition States in Organic Synthesis. Hoboken N.J.: Wiley; 2008.
2. Braun, M. Modern Aldol Reactions. Mahrwald, R., editor. Vol. 1. Weinheim, Germany: Wiley-VCH; 2004. p. 1-61.
3. For related reactions see: Belardi JK, Micalizio GC. J. Am. Chem. Soc. 2008; 130:16870–16872. [PubMed: 19053421] Bartoli G, Bellucci MC, Bosco M, Dalpozzo R, De Nino A, Sambri L, Tagarelli A. Eur. J. Org. Chem. 2000;99–104. Dalpozzo R, De Nino A, Tagarelli A, Bartoli G, Bellucci MC, Bosco M, Sambri L. J. Org. Chem. 1998; 63:9559–9560.

4. For related strategies built on prior hydroxyl activation see: Mandal SK, Paira M, Roy SC. J. Org.Chem. 2008; 73:3823–3827. [PubMed: 18426237] Ramachandran PV, Gurchardt TE, Reddy MVR. Tetrahedron Lett. 2005; 46:2121–2124. Das B, Banerjee J, Mahender G, Majhi A. Org. Lett. 2004; 6:3349–3352. [PubMed: 15355049] Hbaieb S, Ben Ayed T, Amri H. Synth. Commun. 1997; 27:2825–2832. Basavaiah D, Pandiaraju S, Padmaja K. Synlett. 1996:393–395. Basavaiah D, Pandiaraju S. Tetrahedron. 1996; 52:2261–2268.
5. a) Mauzé B, Miginiac L. Synth. Commun. 1992; 22:2229–2235. b) Takayanagi H, Kitano Y, Morinaka Y. Tetrahedron Lett. 1990; 31:3317–3320. c) Matsuda I, Okada H, Izumi Y. Bull. Chem. Soc. Jpn. 1983; 56:528–532.
6. Fatiadi, AJ. Ch 26. In: Patai, S.; Rappoport, Z., editors. The Chemistry of Functional Groups, Supplement C. New York, N.Y.: Wiley; 1983.
7. Fleming FF. Nat. Prod. Rep. 1999; 16:597–606.
8. Fleming FF, Yao L, Ravikumar PC, Funk L, Shook BC. J. Med. Chem. 2010; 53:7902–7917. [PubMed: 20804202]
9. a) Josien-Lefebvre D, Le Drian C. Helv. Chim. Acta. 2007; 90:19–30. b) Krishna V, Maitra S, Dev RV, Mukkanti K, Iqbal J. Tetrahedron Lett. 2006; 47:6103–6106. c) Nagamitsu T, Takano D, Fukuda T, Otoguro K, Kuwajima I, Harigaya Y, Omura S. Org. Lett. 2004; 6:1865–1867. [PubMed: 15151434] d) Smith AB III, Friestad GK, Barbosa J, Bertounesque E, Duan JJ-W, Hull KG, Iwashima M, Qiu Y, Spoors PG, Salvatore BA. J. Am. Chem. Soc. 1999; 121:10478–10486.
10. Roden, BA.; Shook, BC. Magnesium Oxide. Wiley: e-EROS Encyclopedia of Reagents for Organic Synthesis; 2001.
11. For general reviews to Baylis-Hillman reactions see: Basavaiah D, Reddy B, Sekhara B, Satpal S. Chem. Rev. 2010; 110:5447–5674. [PubMed: 20735052] Basavaiah D, Rao KV, Reddy RJ. Chem. Soc. Rev. 2007; 36:1581–1588. [PubMed: 17721583] Basavaiah D, Rao AJ, Satyanarayana T. Chem. Rev. 2003; 103:811–891. [PubMed: 12630854] For ultrasound assisted reactions see: Coelho F, Almeida WP, Veronese D, Mateus CR, Silva Lopes EC, Rossi RC, Silveira GPC, Pavam CH. Tetrahedron. 2002; 58:7437–7447.
12. ¹H NMR analysis of the crude reaction mixture failed to identify any of the *E*-alkenenitrile with aryl carbinols **1a-1c**.
13. Guan S-I, Zhou G-h. Wuhan Huagong Xueyuan Xuebao. 2005; 27:5–7.
14. Alkyl-substituted carbinols react sluggishly and afford numerous uncharacterized materials.
15. a) Vong BG, Kim SH, Abraham S, Theodorakis EA. Angew. Chem., Int. Ed. 2004; 43:3947–3951. b) Duffey MO, LeTiran A, Morken JP. J. Am. Chem. Soc. 2003; 125:1458–1459. [PubMed: 12568588] c) Hanessian S, Yang Y, Giroux S, Mascitti V, Ma J, Raeppe F. J. Am. Chem. Soc. 2003; 125:13784–13792. [PubMed: 14599218] d) Carless HAJ, Dove Y. Tetrahedron Asymm. 1996; 7:649–652. e) Takayanagi H, Sugiyama S, Morinaka Y. J. Chem. Soc., Perkin Trans. 1995; 1:751–756. f) Takayanagi H, Kitano Y, Morinaka Y. J. Org.Chem. 1994; 59:2700–2706. g) Kodama M, Yoshio S, Yamaguchi S, Fukuyama Y, Takayanagi H, Morinaka Y, Usui S, Fukazawa Y. Tetrahedron Lett. 1993; 34:8453–8456.
16. For the enhanced reactivity of magnesiates over alkylmagnesiums see: Hatano M, Matsumura T, Ishihara K. Org. Lett. 2005; 7:573–576. [PubMed: 15704897]
17. Although the rearrangement could potentially proceed through a non-concerted addition-elimination, the stereoselectivity is typically low with alkenes not having *b*-substituents whereas the transmissive olefination exclusively generates *Z*-alkenenitriles: Hoffmann RW. Chem. Rev. 1989; 89:1841–1860.
18. Eliel, EL.; Wilen, SH.; Mander, LN. Stereochemistry of Organic Compounds. NY: Wiley; 1994. p. 597–606.
19. Su B-N, Takaishi Y, Kusumi T. Tetrahedron. 1999; 55:14571–14586.
20. a) Zhu Y, Lu Z-P, Xue C-B, Wu W-S. Helv. Chim. Acta. 2009; 92:536–545. b) Morikawa T, Ando S, Matsuda H, Kataoka S, Muraoka O, Yoshikawa M. Chem. Pharm. Bull. 2005; 53:625–630. [PubMed: 15930771] c) Su B-N, Takaishi Y, Duan H-Q, Chen B. J. Nat. Prod. 1999; 62:1363–1366. [PubMed: 10543893]

21. Winterfeldt E. *Synthesis*. 1975:617–630.. The ratio and efficiency of the two-step reduction was variable with pure (7*Z*, 7'*E*)-**6a** best prepared through an alternative hydrolysis-acylation-reduction sequence.
22. Although other metabolites appear to be derivatives of a precursor having the structure proposed for morinol I, the HMBC and NOE data are inconsistent with the skipped diene structure of **6a**. The ambiguities and the lack of further reports leave synthesis as the best avenue for determining the correct structure. Full details of the transmissive-olefination approach to morinol I alkene diastereomers and structural isomers will be published in a separate account.

Scheme 1.
Transmissive Olefination Route to *Z*-Alkenenitriles.

Scheme 2.
Transmissive Olefination Mechanism.

Scheme 3.
Transmissive Olefination Route to Morinol I.

Table 1

 α' -Hydroxy Alkenenitrile Transmissive Olefination^[a]

Entry	Hydroxynitrile	Grignard	Alkenenitrile	Yield (%)
1	1a	<i>i</i> -PrMgCl	3a	56
2	1a	MeMgCl	3b	62
3	1a	BuMgCl	3c	77
4	1a	<i>t</i> -BuMgCl	3d	54
5	1a	Me ₃ SiCH ₂ MgCl	3e	52
6	1a	BrMg	3f	66
7	1a	MgBr	3g	62
8	1b	<i>i</i> -PrMgCl	3h	60
9	1b	BrMg	3i	63
10	1b	BuMgCl	3j	83

Entry	Hydroxynitrile	Grignard	Alkenenitrile	Yield (%)
11	1c	BuMgCl	3k	68
12	1d	MeMgCl	3l	54 (4:1)
13	1d	BuMgCl	3m	50 (6:1)
14	1d	Me ₃ SiCH ₂ MgCl	3n	52 (6:1)

^[a] An excess of the Grignard reagent was added to the hydroxynitrile **1** at 0 °C and after 50 min. saturated, aqueous NH₄Cl was added.