MspI RFLP for microtubule associated protein-2 (MAP2)

M.J.Alberts, R.S.Kandt, M.A.Pericak-Vance, J.Bebout, M.C.Speer, T.S.Siddique, L.Yamaoka, W.-Y.Hung, P.C.Gaskell and A.D.Roses

The Joseph and Kathleen Bryan Alzheimer's Disease Research Center, Division of Neurology, Department of Medicine, Duke University Medical Center, Durham, NC 27710, USA

Source/Description: The MAP2 probe (KN7) used was a 2.4 kb partial cDNA in the EcoRI site of pBR322. The probe was supplied by R. Neve (1).

Polymorphism: A two allele RFLP was detected after digestion of genomic DNA with MspI. Band sizes of 1.95 kb (A1) and 1.8 kb (A2) were produced.

Frequency: Analysis of 43 unrelated individuals from 22 different families showed these frequencies:

A1 = 0.23

A2 = 0.77

Other Enzymes: Enzymes that did not demonstrate a polymorphism with random placental DNA blots were: EcoRI, HindIII, BglI, BglII, SacI, RsaI, PvuII, HincII, TaqI, PstI and BamHI.

Chromosomal Location: MAP2 has been localized using somatic cell hybrids to the 2q34-q35 region (1).

Mendelian Inheritance: Co-dominant inheritance has been shown based on genotypes of 239 individuals from 24 families.

Probe Availability: Contact Dr. Neve at the Department of Psychobiology, University of California (Irvine).

Other: Double digestion of genomic DNA is recommended to avoid incomplete digestion with MspI.

Acknowledgements: The authors wish to thank Dr. Rachel Neve for supplying the MAP2 probe, and John Thomas, Shelly Bankston, and Mark Nellist for expert technical assistance. This work was supported in part by grants AG 05128 and AG 07922 from the National Institute of Aging, and grant NS 01241 from the National Institute of Neurologic Disorders and Stroke (NINDS). Dr. Alberts is a recipient of a Clinical Investigator Development Award from NINDS.

References: 1) Neve,R.L., Harris,P., Kosik,K.S. *et al.* Identification of cDNA clones for the human microtubule-associated protein tau and chromosomal localization of the genes for tau and microtubule-associated protein 2. (1986) *Mol. Brain Res.* 1, 271-280.

A two allele XbaI RFLP at the catalase 2 locus

M.-A.Colman, S.Shibahara¹, B.Kwon², T.Jenkins and M.Ramsay

MRC Human Ecogenetics Research Unit, Department of Human Genetics, School of Pathology, South African Institute for Medical Research and University of the Witwatersrand, Johannesburg, South Africa, ¹Department of Applied Physiology, Tohoku University School of Medicine, Sendai, Japan and ²Department of Microbiology and Immunology, School of Medicine, Indiana University, IN, USA

Source/Description: pMT4 is a mouse cDNA clone which was isolated from a B16 melanoma cDNA library (1). pMT4 codes for part of the b protein at the b locus (CAS2: catalase 2) in the mouse (2). The human counterpart has been cloned as a 2 kb EcoRI cDNA fragment in the vector pGEM (B. Kwon, unpublished data).

Polymorphism: The mouse and human clones both identify a two allele XbaI RFLP in human genomic DNA, with polymorphic fragment sizes of 10 kb and 8.6 kb, and a constant fragment of 11 kb.

Frequency: Determined in 18 random Negroid individuals and 48 Negroid individuals who are parents of albino children (the frequencies between the two were not significantly different and they are therefore presented together). Number of chromosomes studied is 132.

Allele	Size (kb)	Frequency
1	10.0	0.09 (0.02)
2	8.6	0.91 (0.02)

Not Polymorphic For: StuI, BglII, HpaI, HaeII, PvuII, BamHI, PstI, HindIII, EcoRI, TaqI, AvaII or MspI in ten unrelated individuals.

Chromosome Localization: pMT4 is a clone at or close to the b (catalase 2) locus which has been mapped to mouse chromosome 4 (2). The human syntenic region is on chromosome 9, and the human catalase 2 cDNA is presently being mapped.

Mendelian Inheritance: Co-dominant segregation has been demonstrated in 8 Negroid families.

Probe Availability: Contact S.S. for pMT4 and B.K. for the human catalase 2 cDNA.

Other Comments: For pMT4 the washing stringency is 1×SSC at 65°C.

References: 1) Shibahara *et al.* (1986) *Nucl. Acids Res.* 14, 2413-2427. 2) Jackson (1988) *Proc. Natl. Acad. Sci. USA* 85, 4392-4396.