Published in final edited form as:

J Hypertens. 2008 April ; 26(4): 729–736. doi:10.1097/HJH.0b013e3282f524b4.

Genetic Influences on Blood Pressure Response to the Cold Pressor Test: Results from the HAPI Heart Study

Marie-Hélène Roy-Gagnon^a, Matthew R. Weir^b, John D. Sorkin^{b,c,d}, Kathleen A. Ryan^b, Paul A. Sack^b, Scott Hines^b, Lawrence F. Bielak^e, Patricia A. Peyser^e, Wendy Post^{f,g}, Braxton D. Mitchell^b, Alan R. Shuldiner^{b,c}, and Julie A. Douglas^a

^aDepartment of Human Genetics, University of Michigan, Ann Arbor, MI

^bDepartment of Medicine, University of Maryland, Baltimore, MD

^cGeriatrics Research and Education Clinical Center, Baltimore Veterans Administration Medical Center, Baltimore, MD

^dClaude D. Pepper Older Americans Independence Center, University of Maryland School of Medicine

^eDepartment of Epidemiology, University of Michigan, Ann Arbor, MI

^fDivision of Cardiology, Department of Medicine, Johns Hopkins University, Baltimore, MD

^gDepartment of Epidemiology, Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD

Abstract

Blood pressure (BP) response to the cold pressor test (CPT) has been found to predict the development of hypertension and cardiovascular disease in prospective studies. However, the determinants of BP response to the CPT, including the role of genetic factors, are largely unknown. Additionally, to our knowledge, no study has examined the genetics of BP recovery from the CPT, including whether shared genetic factors influence both reactivity and recovery. As part of the Heredity and Phenotype Intervention (HAPI) Heart Study, we administered a 2½-minute hand CPT to 835 participants from 18 extended Amish families. We estimated the heritability of BP reactivity and recovery (measured by the incremental area under the curve) and the genetic correlations between baseline, reactivity, and recovery BP phenotypes. After adjustment for relevant covariates, including baseline BP, the heritability estimates for both systolic BP (SBP) and diastolic BP (DBP) reactivity and recovery differed significantly from zero ($P < 0.01$), with 12–25% of the total variation in BP response attributable to additive genetic effects. The genetic correlations between baseline DBP and response phenotypes were not significantly different from 0, while the genetic correlation between DBP reactivity and recovery (0.74) was significantly different from 0 and one ($P < 0.005$). The genetic correlation between SBP reactivity and recovery was similar (0.81, $P < 0.05$). We conclude that, independent of baseline BP, BP response to the CPT is heritable, and that both shared and unshared genetic factors influence BP reactivity and recovery, thus stressing the importance of identifying genetic variants that influence both traits.

Address Correspondence to: Julie A. Douglas, Ph.D., University of Michigan School of Medicine, Department of Human Genetics, 1241 E. Catherine St., Buhl Building, Room 5912, Ann Arbor, MI 48109-0618, Tel.: (734) 615-2616, Fax: (734) 763-3784, jddoug@umich.edu.

Conflict of Interest/Disclosure: None

Keywords

cardiovascular reactivity; cold pressor test; blood pressure; genetics; heritability

Introduction

Hypertension is a major risk factor for both stroke and coronary heart disease, the leading causes of death and premature, permanent disability in the United States [1]. Genetic influences on blood pressure (BP) have been clearly established, with most reported heritability estimates of 20–40%, although environmental factors also contribute greatly to BP variation [2–6]. It is likely that the relative importance and types of genes influencing BP variation differ by environmental context. Characterizing genetic influences on BP variation in response to a specific environmental stressor may help focus the search for hypertension susceptibility genes relevant to cardiovascular disease and its treatment and prevention.

In 1932, Hines and Brown [7, 8] introduced the Cold Pressor Test (CPT) as a tool to measure cardiovascular reactivity, specifically BP changes, in response to stress. The CPT stimulus involves both a cold and pain component, which induces a thermoregulatory reflex and global sympathetic activation, producing several physiological responses, including vasoconstriction and BP elevation [9, 10]. Prospective studies have demonstrated that normotensive individuals with a heightened BP reactivity response to the CPT have higher ambulatory BP 3 years later [11], are at a 2–4-fold increased risk of developing hypertension [12–15], and are also more likely to develop clinical coronary heart disease [16]. Recently, the importance of measuring BP during recovery from the CPT has also been emphasized [17, 18], and higher BP levels during recovery have been found to predict increases in BP levels three years later, independently of BP reactivity [19].

Twin studies have suggested a genetic basis for BP reactivity to the CPT [20–22], and a recent family study of BP reactivity reported heritability estimates of 12–37% for systolic BP (SBP) and 2–8% for diastolic BP (DBP) [4]. In addition, results from the latter study suggested that the genetic determinants of baseline and reactivity SBP to the CPT might be different. To our knowledge, the genetics of BP recovery from the CPT, including whether shared genetic factors influence both reactivity and recovery, has not been studied. Such studies would provide insight into the pathophysiology of BP response to the CPT and guidance for future genetic studies involving the CPT.

As part of the Heredity and Phenotype Intervention (HAPI) Heart Study, we administered a 2½-minute hand CPT to 835 participants from 18 extended, Old Order Amish families from Lancaster County, Pennsylvania. The overall goal of the HAPI Heart Study is to identify genes that interact with specific environmental exposures to influence risk factors for cardiovascular disease. In this study, we examined the heritability of phenotypes measuring BP reactivity to and recovery from the CPT. In addition, we investigated whether the same or different genes influence baseline, reactivity, and recovery BP phenotypes.

Methods

Participants

Between 2003 and 2006, we recruited study participants aged 20 years and older from Amish families living in Lancaster County, Pennsylvania. We initially selected participants on the basis of their prior participation in earlier family studies of bone mineral density and coronary artery calcification. Recruitment guidelines were subsequently broadened to allow other interested individuals from the Amish community and their family members to

participate. Consequently, the resulting sample of HAPI Heart Study participants represents a community-based convenience sample of relatively healthy Amish individuals aged 20 to 80 years old (see health exclusions below).

Women were excluded from the HAPI Heart Study if they were currently pregnant. However, they were eligible to participate one year after delivery. Other study-wide exclusion criteria included severe hypertension (SBP/DBP greater than 180/105 mm Hg), co-existing malignancy, creatinine level greater than 2 mg/dl, Aspartate transaminase (AST) or Alanine transaminase (ALT) greater than twice the normal values, Hematocrit less than 32%, and/or abnormal levels of thyroid stimulating hormone. We also excluded individuals if they were taking medications potentially affecting any outcome measures that could not be discontinued for one week prior to initiation of the protocol. These medications included all anti-hypertensive medications, nitrates, lipid lowering agents, anti-diabetic medications (with the exception of insulin), adrenergic or cholinergic acting agents (including cold formulas and anti-depressants), and diet or weight loss agents. Anti-platelet agents and vitamins were discontinued 14 and 7 days, respectively, prior to starting the study. In addition to these study-wide exclusion criteria, participants were excluded from the CPT intervention if they had Raynaud's disease.

Eight hundred sixty-eight participants were enrolled in the HAPI Heart Study. Of these, 835 completed the CPT and were included in our analyses. These participants were distributed across 518 nuclear families. For the purpose of these analyses, we used the extensive genealogical information available from the Amish Genealogical Database [23] to combine nuclear families into 18 extended families by including genealogical information on grandparents of the participants even if they did not participate in the study. All participants gave written, informed consent as approved by the Institutional Review Board at the University of Maryland and collaborating institutions prior to participation.

Study Design

Variables assessing medical history and lifestyle were obtained during a home visit by field staff through questionnaires. During the participant's first visit at the Amish Research Clinic, a physical examination was performed. Height (cm), weight (kg), waist and hip circumference (cm) were measured. Body mass index (BMI) was calculated as weight (kg) divided by height (m) squared. Radial pulse (bpm) was measured three times, and resting heart rate was calculated as the average of the last two measurements.

The CPT was performed during the first clinic visit as described by Corretti et al. (1995) [24]. Stimuli other than the CPT (e.g., pharmacological agents) can be used to trigger cardiovascular response, and several measures of cardiovascular response (e.g., forearm blood flow or epicardial coronary artery dysfunction assessed by quantitative coronary angiography) exist. We chose to measure BP response to the CPT because it is well suited for the rural clinic setting of the HAPI Heart Study.

All participants were studied in a temperature-controlled room (22 °C) in the resting, supine state in the morning (following a 12-hour fast). Participants were instructed not to engage in exercise or any vigorous activity on the morning of the study. BP and heart rate were measured with a properly fitted automated BP cuff (Datascop Acutor 3SAT). BP measurements were made before the CPT every 5 minutes for at least 20 minutes or until a stable baseline BP was obtained (i.e., no more than 3mm Hg difference between 3 consecutive readings for both SBP and DBP). The right hand and wrist were immersed in ice water for 2½ minutes and then withdrawn. Nine blood pressure measurements were taken during and after the cold pressor stimulus at minutes 1, 2, 3, 4, 5, 7.5, 10, 15, and 20.

Statistical Analysis

All analyses were carried out separately for SBP and DBP. We used the average of the last two BP measurements taken before the CPT as baseline BP, denoted as time 0. Over 90% of participants reached their maximum change in BP by 2 minutes, and the majority had a stable BP after 5 minutes. Therefore, measurements at 1 and 2 minutes were considered to reflect reactivity to the CPT, while measurements at 3, 4, and 5 minutes were considered to reflect recovery. To measure BP response, we calculated the incremental area under the curve (iAUC) separately for the reactivity and recovery periods. Within each phase, the iAUC was defined as the difference between the area under the response curve and the area below baseline and was calculated by first subtracting baseline from all subsequent measurements before calculating the area (see Figure 1) [25, 26]. We used the trapezoidal rule to calculate the areas [27].

We performed heritability analyses using maximum likelihood variance components methods as implemented in the program SOLAR [28]. For a particular phenotype y , the value of y for individual i is modeled as $y_i = \mu + \sum \beta_j X_{ij} + g_i + e_i$, where μ is the mean of y , X_{ij} is the j -th covariate with associated regression coefficient β_j , g_i is an additive genetic effect normally distributed with mean 0 and variance σ_g^2 , and e_i is a random residual effect normally distributed with mean 0 and variance σ_e^2 . Any non-additive genetic and unmeasured non-genetic effects (as well as random errors) are incorporated into e_i . The heritability of the phenotype is estimated by the ratio of the variance attributable to additive genetic effects, σ_g^2 , to the total phenotypic variance. We used likelihood-ratio tests to assess the significance of a parameter of interest by comparing the log-likelihood of the model in which the parameter is estimated to that of the model in which the parameter is fixed at 0 [29]. We adjusted all heritability estimates for covariates that were associated with at least one of the phenotypes (at a $P < 0.1$). We performed covariate screening in SOLAR. Significant covariates were age, age², sex, age- and age²-by-sex interactions, BMI, and resting heart rate, a surrogate for cardiovascular fitness (that was measured during the clinical examination). For the response traits, we also included baseline BP as a covariate. We estimated the total proportion of variance explained by additive genetic effects as the product of the heritability estimate and one minus the proportion of the variance explained by covariates.

We also performed bivariate analyses using maximum likelihood methods as implemented in the program SOLAR [30–32] in order to assess the presence of pleiotropic genetic effects influencing baseline and reactivity, baseline and recovery, and reactivity and recovery. We partitioned the phenotypic correlation (ρ_P) between two traits into components attributable to shared genetic effects (genetic correlation, ρ_G) and shared environmental factors (environmental correlation, ρ_E). If h_1^2 and h_2^2 are the heritabilities of the two traits, then

$$\rho_P = \rho_G \sqrt{h_1^2 h_2^2} + \rho_E \sqrt{(1 - h_1^2)(1 - h_2^2)}$$

Using likelihood-ratio tests, we tested two different hypotheses: $\rho_G = 0$ and $\rho_G = 1$ (or -1). Rejecting $\rho_G = 0$ provides evidence that shared genetic effects exist, while rejecting $\rho_G = 1$ (or -1) indicates that one or more genetic factors are specific to each trait. We also tested the hypothesis that $\rho_E = 0$ to assess the presence of shared environmental factors affecting the two traits. Covariates included in the bivariate analyses were the same as in the univariate analyses. Baseline BP was included as a covariate when the bivariate analysis included both response traits.

After adjustment for covariates, the BP phenotypes were close to normally distributed except for the presence of outliers. We assessed the impact of outliers on the estimates of heritability and genetic and environmental correlation by examining the change in the estimates when extreme values were excluded. All analyses (except where noted above) were conducted using version 8.2 of the Statistical Analysis System programming language (SAS Institute).

Results

The 835 HAPI Heart participants (387 women) who underwent the CPT intervention had a mean age of 44 ± 14 years ($\pm$ standard deviation). Selected characteristics of these participants are shown in Table 1 by sex. Mean body mass index (BMI) was 25.6 ± 3.2 kg/m² for men and 27.8 ± 5.5 kg/m² for women, and mean resting heart rate was 62 ± 7 bpm for men and 66 ± 7 bpm for women. The proportion of participants with hypertension (defined by baseline SBP > 140 mm Hg or DBP > 90 mm Hg) was 3.6% for men and 11.1% for women (data not shown). Less than 4% of participants reported that they currently smoke (16 men and 0 women, data not shown). The participants were distributed across 518 nuclear families from 18 extended families, including 581 sibling pairs, 302 parent-offspring pairs, and 10,113 more distantly related pairs.

Maximum BP response was attained within 2 minutes for over 90% of participants. On average, participants' BP returned to baseline levels by 5 minutes. Few participants (4.1%) had a negative BP response to the CPT, meaning that no BP measurement during the reactivity part of the CPT was greater than or equal to their baseline BP. Of note, with the exception of self-reported diabetes, there were no significant differences in baseline clinical or demographic characteristics (e.g., age, sex, body size, baseline BP, and hypertension status) between negative and non-negative responders ($P > 0.05$). In fact, after comparing 17 characteristics, only the proportion of individuals who self-reported diabetes (in total, 3 negative and 11 non-negative responders) was significantly different between the two groups ($P = 0.02$ for Fisher's exact test). In addition, the majority of the negative responders were not closely related to one another (data not shown).

Mean baseline SBP and DBP were 117 ± 13 mm Hg and 71 ± 8 mm Hg, respectively. Mean reactivity, as measured by the iAUC from 0 to 2 minutes, was 28 ± 17 (mm Hg)² for SBP and 16 ± 12 (mm Hg)² for DBP. Descriptive statistics of baseline BP and the response phenotypes are shown in Table 2 by sex. For most phenotypes, BP response was greater in men than in women and was positively associated with age and BMI and negatively associated with heart rate and baseline BP ($P < 0.05$, data not shown). As an illustration of both the familial resemblance of response patterns and the variation in the magnitude of response, Figure 2 shows SBP response profiles for two different father-son pairs from the HAPI Heart Study.

After adjustment for covariates, heritability estimates for baseline SBP and DBP were 0.27 and 0.21, respectively ($P < 0.001$, see Table 3). After adjustment for covariates, including baseline BP, all heritability estimates of BP responses (as measured by the iAUC) were also significantly different from zero ($P < 0.01$), ranging from 0.14 (for SBP recovery) to 0.27 (for DBP reactivity). Compared to measured covariates, additive genetic factors accounted for a greater proportion of the total variance in DBP response and for a similar proportion of the variance in SBP response. For example, additive genetic effects and covariates accounted for 25% and 5%, respectively, of the total variance in DBP reactivity and for 13% and 14% of the total variance in SBP reactivity (see Table 3).

After adjustment for covariates, the estimates of phenotypic correlation (ρ_P) between baseline and response BP phenotypes were negative and ranged from -0.13 (SBP reactivity)

to -0.23 (DBP recovery). Table 4 shows the partition of ρ_P into genetic and environmental correlations. The genetic correlation (ρ_G) between baseline and response DBP phenotypes was significantly different from one ($P < 0.001$) but not 0 ($P > 0.4$), suggesting that different sets of genes may be influencing baseline and response DBP phenotypes. For SBP, ρ_G estimates between baseline and response phenotypes were 0.43 and 0.86 for reactivity and recovery, respectively, and neither was significantly different from one ($P > 0.08$). Environmental correlation (ρ_E) estimates between all baseline and response BP phenotypes were negative and differed significantly from 0 ($\rho_E = -0.25$, $P < 0.001$), except for baseline DBP and reactivity ($\rho_E = -0.13$, $P = 0.08$). Hence, unmeasured environmental factors may influence both baseline and response BP phenotypes in opposite directions.

The reactivity and recovery periods of the BP response to the CPT were positively correlated ($\rho_P = 0.73$ for SBP and 0.68 for DBP). Estimates of the genetic correlation between reactivity and recovery were significantly different from 0 and one ($\rho_G = 0.81$, $P < 0.05$ for SBP; $\rho_G = 0.74$, $P < 0.005$ for DBP), providing evidence that shared (i.e., pleiotropic) and unique genetic effects influence both reactivity and recovery. Estimates of the environmental correlation differed significantly from 0 ($\rho_E = -0.70$, $P < 0.0001$), indicating that unmeasured environmental factors influence both reactivity and recovery in the same direction.

We repeated all heritability and genetic and environmental correlation analyses after removing individuals with values more extreme than 3 standard deviations from the mean (3 to 14 participants). All heritability and correlation estimates from these analyses were within one standard error of the original estimates.

Discussion

Our study is the largest family study of BP response to the CPT to date and is the only family study to obtain multiple measurements before, during, and after the CPT. To our knowledge, we are also the first to demonstrate that BP recovery from the CPT is heritable and to estimate the genetic correlation between the reactivity and recovery periods of BP response to the CPT. We found that both BP reactivity and recovery were significantly heritable and estimated that additive genetic effects explained 12–25% of the total variance in BP response. In other words, while no inferences can be made regarding specific individuals or families in the present study, these estimates suggest that in the population from which the pedigrees were sampled, genetic differences among individuals explain 1/8th to 1/4th of the observed variation in the CPT responses. Both SBP and DBP reactivity were previously found to be heritable in twin studies [20–22], while only SBP reactivity was found to be heritable in the family study by Choh et al. [4].

Our results suggest that DBP response to the CPT captures a physiological phenomenon that is different from baseline DBP, i.e., there may be genetic factors relevant to the etiology of cardiovascular disease that influence CPT-derived BP response but not baseline BP. We also found evidence that both shared and unique genetic factors influence DBP reactivity and recovery. For SBP response, we could not reject the hypothesis of complete pleiotropy of genetic effects on baseline and response SBP phenotypes ($P = 0.09$). However, the genetic correlation between baseline and reactivity SBP (0.43) was also not significantly different from 0 ($P = 0.18$). Of note, Choh et al. [4] found no evidence for pleiotropic effects of genes on baseline and reactivity SBP to the CPT, in agreement with two previous twin studies [21, 22]. Further investigation is needed to determine if both shared and unshared genetic factors influence baseline, reactivity, and recovery SBP phenotypes.

We found that a large proportion of the variation in BP response was left unexplained after accounting for measured covariates and unmeasured additive genetic effects. Part of this

unexplained variation could be due to measurement error, although previous studies have found that CPT-derived BP responses were reproducible, especially when the responses were derived from several measurements during the test [33–35]. Other possible factors that may explain the residual variation in BP response include dietary intake [36] and physical activity/cardiovascular fitness [37, 38]. However, variation in dietary intake is likely to be limited in the Amish due to their communal lifestyle. In addition, resting heart rate (a measure of cardiovascular fitness) was included in our analyses but explained less than 3% and 1% of the variation in SBP and DBP response, respectively.

The Old Order Amish is a Caucasian population with a relatively homogeneous lifestyle and similar environmental exposures. For example, the Amish rarely use prescription medications or smoke. The homogeneity of environmental exposures in the Amish represents a strength of our study by minimizing the effects of important, potential confounders. It also permits a more controlled investigation of the role of genetic factors in BP response to the CPT than has been possible in previous studies. Since the Amish are derived from the same overall gene pool as U.S. and European Caucasian populations, we believe that the genes influencing BP response in the Amish will be relevant to the general population.

Differences in BP response to the CPT are likely to be influenced by several physiological pathways and different from those controlling baseline BP. For example, in addition to physiological pathways related to global sympathetic activation, endothelial function and psychological components (related to the perception of pain and the degree of difficulty of the test) may influence BP response [39, 40]. In addition, Busjahn et al. [22] suggested that long-term regulatory systems influence baseline BP while systems controlling acute sympathetic activation are involved in BP reactivity. Our results provide new information implicating both shared and unique pathways/genes in the reactivity and recovery periods of BP response to the CPT. For example, systems involved in catecholamine reuptake are likely to be specific to BP recovery [41]. Still, at present, little is known about the specific pathways involved in and the genetic factors that contribute to and/or distinguish BP reactivity and recovery.

Importantly, BP response to the CPT predicts the development of hypertension and coronary heart disease, even in normotensive individuals [12–16]. Moreover, as described above, we found that a significant proportion of the variation in BP response to the CPT is independent of baseline BP and likely due to a heritable (genetic) component that is also independent of baseline BP. In fact, the BP response profiles in our sample were remarkably similar between individuals who would be classified [42] as normotensive (baseline SBP and DBP less than 120 and 80 mm Hg, respectively) versus hypertensive (baseline SBP or DBP greater than or equal to 140 or 90 mm Hg, respectively). For example, after adjustment for relevant covariates, including baseline BP, mean BP reactivity was not significantly different between normotensive and hypertensive individuals ($P = 0.44$ and $P = 0.23$ for SBP and DBP reactivity, respectively). Thus, there are normotensive individuals who are at increased risk of developing hypertension and coronary heart disease on the basis of their BP reactivity to the CPT yet who would not come to clinical attention on the basis of their baseline BP alone. Together, these findings emphasize the importance of understanding how genetic factors influence BP response to the CPT as this information may provide valuable insight into the pathophysiology of BP, i.e., insight that could not be surmised by simply studying baseline BP.

Conclusion

Our results provide new evidence for a genetic basis for BP recovery from the CPT and add to the existing evidence that BP reactivity to the CPT is heritable. Our results also

emphasize the importance of considering both reactivity to and recovery from the CPT since both shared and unique genetic factors may influence the reactivity and recovery periods of the BP response to the CPT. Findings from this study will inform and guide our search for the specific genes that underlie BP response to the CPT. Identifying genes that influence BP response to the CPT is clinically relevant given the accumulating evidence that BP measured under environmental stressors may predict the risk of developing hypertension. In turn, understanding the genetic basis of cardiovascular reactivity may facilitate future efforts to improve cardiovascular health.

Acknowledgments

Sources of Funding

This study was supported in part by NIH grants U01 HL72515, T32 AG000219 and the American Diabetes Association. Funding and support was also provided by the University of Maryland General Clinical Research Center (M01 RR 16500), the NIDDK Clinical Nutrition Research Unit of Maryland (NIH P30 DK072488), the Department of Veterans Affairs and Veterans Affairs Medical Center Baltimore Geriatric Research, Education and Clinical Center (GRECC), and the University of Maryland Claude D. Pepper Older Americans Independence Center (NIA P60AG 12583).

We thank the Amish Research Clinic Staff for their energetic efforts in study subject recruitment and characterization and Drs. Alejandro Schäffer and Richa Agarwala for assistance with pedigree construction. This study would not have been possible without the outstanding cooperation and support of the Amish community.

References

1. Rosamond W, Flegal K, Friday G, Furie K, Go A, Greenlund K, et al. Heart disease and stroke statistics--2007 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2007; 115:e69–171. [PubMed: 17194875]
2. Mitchell BD, Kammerer CM, Blangero J, Mahaney MC, Rainwater DL, Dyke B, et al. Genetic and Environmental Contributions to Cardiovascular Risk Factors in Mexican Americans: The San Antonio Family Heart Study. *Circulation*. 1996; 94:2159–2170. [PubMed: 8901667]
3. Gu C, Borecki I, Gagnon J, Bouchard C, Leon AS, Skinner JS, et al. Familial resemblance for resting blood pressure with particular reference to racial differences: preliminary analyses from the HERITAGE Family Study. *Hum Biol*. 1998; 70:77–90. [PubMed: 9489236]
4. Choh AC, Czerwinski SA, Lee M, Demerath EW, Wilson AF, Towne B, et al. Quantitative genetic analysis of blood pressure response during the cold pressor test. *Am J Hypertens*. 2005; 18:1211–1217. [PubMed: 16182112]
5. Levy D, DeStefano AL, Larson MG, O'Donnell CJ, Lifton RP, Gavvas H, et al. Evidence for a Gene Influencing Blood Pressure on Chromosome 17: Genome Scan Linkage Results for Longitudinal Blood Pressure Phenotypes in Subjects From the Framingham Heart Study. *Hypertension*. 2000; 36:477–483. [PubMed: 11040222]
6. Perusse L, Rice T, Bouchard C, Vogler GP, Rao DC. Cardiovascular risk factors in a French-Canadian population: resolution of genetic and familial environmental effects on blood pressure by using extensive information on environmental correlates. *Am J Hum Genet*. 1989; 45:240–251. [PubMed: 2757030]
7. Hines J, Brown GE. A standard stimulant for measuring vasomotor reactions: its application in the study of hypertension. *Proc Staff Meet Mayo Clin*. 1932; 7:332–335.
8. Hines J, Brown GE. The cold pressor test for measuring the reactivity of the blood pressure: Data concerning 571 normal and hypertensive subjects. *Am Heart J*. 1936; 11:1–9.
9. Peckerman A, Hurwitz BE, Saab PG, Llabre MM, McCabe PM, Schneiderman N. Stimulus dimensions of the cold pressor test and the associated patterns of cardiovascular response. *Psychophysiology*. 1994; 31:282–290. [PubMed: 8008792]
10. Lovallo W. The cold pressor test and autonomic function: a review and integration. *Psychophysiology*. 1975; 12:268–282. [PubMed: 1153632]

11. Knox SS, Hausdorff J, Markovitz JH. Reactivity as a predictor of subsequent blood pressure: racial differences in the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Hypertension*. 2002; 40:914–919. [PubMed: 12468579]
12. Wood DL, Sheps SG, Elveback LR, Schirger A. Cold pressor test as a predictor of hypertension. *Hypertension*. 1984; 6:301–306. [PubMed: 6735451]
13. Menkes MS, Matthews KA, Krantz DS, Lundberg U, Mead LA, Qaqish B, et al. Cardiovascular reactivity to the cold pressor test as a predictor of hypertension. *Hypertension*. 1989; 14:524–530. [PubMed: 2807514]
14. Kasagi F, Akahoshi M, Shimaoka K. Relation Between Cold Pressor Test and Development of Hypertension Based on 28-Year Follow-up. *Hypertension*. 1995; 25:71–76. [PubMed: 7843757]
15. Matthews KA, Katholi CR, McCreath H, Whooley MA, Williams DR, Zhu S, et al. Blood pressure reactivity to psychological stress predicts hypertension in the CARDIA study. *Circulation*. 2004; 110:74–78. [PubMed: 15210592]
16. Keys A, Taylor HL, Blackburn H, Brozek J, Anderson JT, Simonson E. Mortality and coronary heart disease among men studied for 23 years. *Arch Intern Med*. 1971; 128:201–214. [PubMed: 5564199]
17. Linden W, Earle TL, Gerin W, Christenfeld N. Physiological stress reactivity and recovery: Conceptual siblings separated at birth? *J of Psychosom Res*. 1997; 42:117–135. [PubMed: 9076640]
18. Llabre MM, Spitzer SB, Saab PG, Schneiderman N. Piecewise latent growth curve modeling of systolic blood pressure reactivity and recovery from the cold pressor test. *Psychophysiology*. 2001; 38:951–960. [PubMed: 12240671]
19. Stewart JC, France CR. Cardiovascular recovery from stress predicts longitudinal changes in blood pressure. *Biol Psychol*. 2001; 58:105–120. [PubMed: 11600240]
20. McIlhany ML, Shaffer JW, Hines EA Jr. The heritability of blood pressure: an investigation of 200 pairs of twins using the cold pressor test. *Johns Hopkins Med J*. 1975; 136:57–64. [PubMed: 1167918]
21. Ditto B. Familial influences on heart rate, blood pressure, and self-report anxiety responses to stress: results from 100 twin pairs. *Psychophysiology*. 1993; 30:635–645. [PubMed: 8248455]
22. Busjahn A, Faulhaber HD, Viken RJ, Rose RJ, Luft FC. Genetic influences on blood pressure with the cold-pressor test: a twin study. *J Hypertens*. 1996; 14:1195–1199. [PubMed: 8906518]
23. Agarwala R, Biesecker LG, Schaffer AA. Anabaptist genealogy database. *Am J Med Genet C Semin Med Genet*. 2003; 121:32–37. [PubMed: 12888984]
24. Corretti MC, Plotnick GD, Vogel RA. Correlation of cold pressor and flow-mediated brachial artery diameter responses with the presence of coronary artery disease. *Am J Cardiol*. 1995; 75:783–787. [PubMed: 7717279]
25. Le Floch JP, Escuyer P, Baudin E, Baudon D, Perlemuter L. Blood glucose area under the curve. Methodological aspects. *Diabetes Care*. 1990; 13:172–175. [PubMed: 2351014]
26. Allison DB, Paultre F, Maggio C, Mezzitis N, Pi-Sunyer FX. The use of areas under curves in diabetes research. *Diabetes Care*. 1995; 18:245–250. [PubMed: 7729306]
27. Matthews JN, Altman DG, Campbell MJ, Royston P. Analysis of serial measurements in medical research. *BMJ*. 1990; 300:230–235. [PubMed: 2106931]
28. Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet*. 1998; 62:1198–1211. [PubMed: 9545414]
29. Self SG, Liang KY. Asymptotic Properties of Maximum Likelihood Estimators and Likelihood Ratio Tests Under Nonstandard Conditions. *J Am Statist Assoc*. 1987; 82:605–610.
30. Hopper JL, Mathews JD. Extensions to multivariate normal models for pedigree analysis. *Ann Hum Genet*. 1982; 46:373–383. [PubMed: 6961886]
31. Lange K, Boehnke M. Extensions to pedigree analysis. IV. Covariance components models for multivariate traits. *Am J Med Genet*. 1983; 14:513–524. [PubMed: 6859102]
32. Boehnke M. Estimating the power of a proposed linkage study: a practical computer simulation approach. *Am J Hum Genet*. 1986; 39:513–527. [PubMed: 3464203]

33. Durel LA, Kus LA, Anderson NB, McNeilly M, Llabre MM, Spitzer S, et al. Patterns and stability of cardiovascular responses to variations of the cold pressor test. *Psychophysiology*. 1993; 30:39–46. [PubMed: 8416061]
34. Llabre MM, Saab PG, Hurwitz BE, Schneiderman N, Frame CA, Spitzer S, et al. The stability of cardiovascular parameters under different behavioral challenges: one-year follow-up. *Int J Psychophysiol*. 1993; 14:241–248. [PubMed: 8340242]
35. Kelsey RM, Ornduff SR, Alpert BS. Reliability of cardiovascular reactivity to stress: Internal consistency. *Psychophysiology*. 2007; 44:216–225. [PubMed: 17343705]
36. Monahan KD, Wilson TE, Ray CA. Omega-3 fatty acid supplementation augments sympathetic nerve activity responses to physiological stressors in humans. *Hypertension*. 2004; 44:732–738. [PubMed: 15452023]
37. Jackson EM, Dishman RK. Hemodynamic responses to stress among black women: fitness and parental hypertension. *Med Sci Sports Exerc*. 2002; 34:1097–1104. [PubMed: 12131247]
38. Dishman RK, Jackson EM, Nakamura Y. Influence of fitness and gender on blood pressure responses during active or passive stress. *Psychophysiology*. 2002; 39:568–576. [PubMed: 12236322]
39. Kamarck TW, Lohvallo WR. Cardiovascular Reactivity to Psychological Challenge: Conceptual and Measurement Considerations. *Psychosom Med*. 2003; 65:9–21. [PubMed: 12554812]
40. Peckerman A, Saab PG, McCabe PM, Skyler JS, Winters RW, Llabre MM, et al. Blood pressure reactivity and perception of pain during the forehead cold pressor test. *Psychophysiology*. 1991; 28:485–495. [PubMed: 1758926]
41. Haynes SN, Gannon LR, Orimoto L, O'Brien WH. Psychophysiological assessment of poststress recovery. *Psychological Assessment*. 1991; 3:356–365.
42. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003; 42:1206–52. [PubMed: 14656957]

Figure 1.

Response profile from one HAPI Heart Study participant illustrating the blood pressure (BP) phenotypes derived from the Cold Pressor Test. Baseline BP is shown by the bold, horizontal line. The reactivity (0–2 minutes) and recovery (2–5 minutes) periods are delimited by vertical lines at times 0, 2, and 5 minutes. We used the incremental area under the curve (iAUC) to measure the two periods of BP response, i.e., reactivity and recovery, separately. Within each period, the iAUC was defined as the difference between the area under the response curve and the area below baseline. Hence, reactivity iAUC corresponds to area A (vertical lines), while recovery iAUC is represented by the difference between areas B and C (horizontal lines).

Figure 2. Familial resemblance of response profiles: individual profiles of systolic blood pressure response to the Cold Pressor Test for two different father-son pairs from the HAPI Heart Study.

Table 1

Characteristics of the 835 HAPI Heart Study participants who underwent the Cold Pressor Test

	Men (n = 448)		Women (n = 387)	
	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range
Age (years)	42 $\pm$ 13	[20, 80]	45 $\pm$ 14	[20, 80]
Height (cm)	173 $\pm$ 6	[147, 194]	161 $\pm$ 6	[141, 175]
Weight (kg)	77 $\pm$ 11	[54, 110]	72 $\pm$ 14	[42, 122]
Body Mass Index (kg/m ²)	25.6 $\pm$ 3.2	[18.4, 37.9]	27.8 $\pm$ 5.5	[17.2, 46.8]
Resting heart rate (bpm)	62 $\pm$ 7	[41, 92]	66 $\pm$ 7	[52, 96]

SD = standard deviation

Table 2

Summary of baseline blood pressure and blood pressure response to the Cold Pressor Test

	Men (n = 448)		Women (n = 387)	
	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range
<u>Systolic blood pressure</u>				
Baseline (mm Hg)	116 $\pm$ 11	[80, 173]	117 $\pm$ 15	[89, 175]
iAUC (mm Hg) ² :				
Reactivity (0 – 2 min.)	27 $\pm$ 16	[–13, 88]	29 $\pm$ 17	[–19, 84]
Recovery (2 – 5 min.)	31 $\pm$ 29	[–52, 186]	25 $\pm$ 28	[–44, 124]
<u>Diastolic blood pressure</u>				
Baseline (mm Hg)	71 $\pm$ 7	[52, 103]	70 $\pm$ 8	[55, 95]
iAUC (mm Hg) ² :				
Reactivity (0 – 2 min.)	17 $\pm$ 13	[–17, 69]	15 $\pm$ 11	[–14, 61]
Recovery (2 – 5 min.)	13 $\pm$ 19	[–45, 79]	6 $\pm$ 16	[–42, 48]

SD = standard deviation

iAUC = incremental Area Under the Curve

Table 3

Heritability estimates for baseline blood pressure and blood pressure response to the Cold Pressor Test

Trait*	Heritability Estimate (Standard Error)	P value	% of variance explained by	
			Measured Covariates*	Additive Genetic Effects
<u>Systolic blood pressure</u>				
Baseline	0.27 (0.08)	< 0.0001	26	20
Response (iAUC):				
Reactivity (0 – 2 min.)	0.15 (0.07)	0.005	14	13
Recovery (2 – 5 min.)	0.14 (0.06)	0.006	15	12
<u>Diastolic blood pressure</u>				
Baseline	0.21 (0.08)	0.0008	23	17
Response (iAUC):				
Reactivity (0 – 2 min.)	0.27 (0.08)	< 0.0001	5	25
Recovery (2 – 5 min.)	0.20 (0.07)	0.0003	10	18

iAUC = incremental Area Under the Curve

* Adjusted for age, age², sex, age- and age²-by-sex interactions, body mass index, resting heart rate, and for response phenotypes only the baseline blood pressure.

Genetic (ρ_G) and environmental (ρ_E) correlation between baseline blood pressure and blood pressure response phenotypes to the Cold Pressor Test

Table 4

Pairs of traits	ρ_G (Standard Error)	P value		ρ_E (Standard Error)	P value $H_0: \rho_E = 0$
		$H_0: \rho_G = 0$	$H_0: \rho_G = 1$ (or -1)		
<u>Systolic blood pressure</u>					
Baseline - Reactivity iAUC [*]	0.43 (0.35)	0.18	0.087	-0.26 (0.07)	0.0002
Baseline - Recovery iAUC [*]	0.86 (0.45)	0.013	0.39	-0.38 (0.06)	< 0.0001
Reactivity iAUC - Recovery iAUC ^{**}	0.81 (0.14)	0.024	0.041	0.72 (0.03)	< 0.0001
<u>Diastolic blood pressure</u>					
Baseline - Reactivity iAUC [*]	-0.18 (0.23)	0.45	0.0007	-0.13 (0.07)	0.084
Baseline - Recovery iAUC [*]	-0.13 (0.26)	0.62	0.0007	-0.25 (0.07)	0.0004
Reactivity iAUC - Recovery iAUC ^{**}	0.74 (0.12)	0.002	0.003	0.66 (0.04)	< 0.0001

iAUC = incremental Area Under the Curve

* Adjusted for age, age², sex, age-by-sex interactions, body mass index, and resting heart rate.

** Adjusted for age, age², sex, age-by-sex interactions, body mass index, resting heart rate, and baseline blood pressure.

Note – The genetic and environmental correlation between a pair of traits (expressed as ρ_G and ρ_E , respectively) captures the extent to which the same genes and the same environmental factors influence both traits. Rejection of the hypothesis that $\rho_G = 0$ suggests that one or more of the same genetic factors influences both traits. Rejection of the hypothesis that $\rho_G = 1$ (or -1) suggests that one or more unique genetic factors influences one trait but not the other (i.e., all of the genetic factors that influence one trait are not the same as all of the genetic factors that influence the other trait). Rejection of the hypothesis that $\rho_E = 0$ indicates that one or more of the same environmental factors influences both traits.