TaqI polymorphism at D5S205, located within 5q11.2 – 13.3

L.E. Bernard and S. Wood*

Department of Medical Genetics, Room 312 Wesbrook Bldg, University of British Columbia, 6174 University Boulevard, Vancouver BC V6T 1W5, Canada

Source/Description: p5PCR1LIC-1 is a 2.2 kb inter-Alu fragment subcloned into XbaI digested pUC19 by LIC tailing (1). This fragment was prepared by Alu primed PCR from phage 5PCR1 (2) isolated from LA05NS01, a complete EcoRI digest chromosome 5 library obtained from Los Alamos National Laboratory.

Polymorphism: TaqI identifies a four allele polymorphism with bands at 9 kb (A1), 6 kb (A2), 3.2 kb (A3) or 11 kb (A4).

Allele Frequencies: Calculated from 80 parents in the CEPH panel
A1 allele (9 kb): 0.12
A2 allele (6 kb): 0.74
A3 allele (3.2 kb): 0.12
A4 allele (11 kb): 0.02

Observed PIC Value: 0.39.

Not Polymorphic For: BamHI, BglII, EcoRI, HindIII, PstI, PvuII.

Chromosomal Localization: Located within 5q11.2–q13.3 by hybridization to somatic cell hybrid DNA (2).

Mendelian Inheritance: Codominant segregation was observed in 29 CEPH families.

Probe Availability: Submitted to ATCC.

Acknowledgements: This work was supported by a grant from the B.C. Health Care Research Foundation. L.E.B. is a recipient of a Medical Research Council of Canada studentship.

References: 1) Aslanidis, C. and deJong, P.J. (1990) *Nucl. Acids Res.* **18**, 6069–6074. 2) Bernard, L.E. *et al.* (1990) *Genomics*, in press.

Figure 1. CEPH Family 1349 generation III sibship showing alleles A1 and A2.

Dinucleotide repeat polymorphism at the human c-fms proto-oncogene for the CFS-1 receptor (CFS1R)

M.H. Polymeropoulos, H. Xiao, D.S. Rath and C.R. Merrill
National Institute of Mental Health Neuroscience Center, St Elizabeths Hospital, Room 131, 2700 Martin Luther King Avenue, Washington, DC 20032, USA

Source/Description: The polymorphic (GT)_n repeat begins at base pair 4938 in intron II of the human c-fms proto-oncogene for the CFS-1 receptor on chromosome 5q33.3–34 (1). The polymorphism can be typed using the polymerase chain reaction (PCR) as described previously (2). The predicted length of the amplified sequence was 117 bp.

Primer Sequences: TGTGTCCAGCCTTAGTGTGCA (GT strand); TCATCACTTCCAGAATGTGC (CA strand).

Frequency: Estimated from 40 chromosomes of unrelated individuals. Observed heterozygosity = 86%. PIC = 0.85.

Allele (bp)	Frequency	Allele (bp)	Frequency
C1 127	0.03	C5 117	0.10
C2 125	0.07	C6 115	0.08
C3 123	0.03	C7 113	0.22
C4 121	0.12	C8 111	0.05
C4 119	0.20	C9 95	0.10

Mendelian Inheritance: Co-dominant segregation was observed in two informative families.

Chromosomal Localization: The human c-fms proto-oncogene for the CFS-1 receptor has been assigned to chromosome 5q33.3–34 (1).

Other Comments: The PCR reaction was performed on 80 ng of genomic DNA using 100 pmoles of each oligonucleotide primer. The samples were processed as described (3) except that the denaturation cycle at 94°C was extended to 1.4 minutes. The dinucleotide repeat was based on a (GT)₂₆ sequence.

References: 1) Hampe, A. *et al.* (1989) *Oncogene Res.* **4**, 9–17. 2) Weber, J.L. and May, P.E. (1989) *Am. J. Hum. Genet.* **44**, 388–396. 3) Weber, J.L. *et al.* (1990) *Nucl. Acids Res.* **18**, 4637.

* To whom correspondence should be addressed