Consumption of Rice Bran Increases Mucosal Immunoglobulin A Concentrations and Numbers of Intestinal *Lactobacillus* spp.

Angela J. Henderson,¹ Ajay Kumar,² Brittany Barnett,³ Steven W. Dow,^{1,2} and Elizabeth P. Ryan²

Departments of ¹Microbiology, Immunology, and Pathology and ²Clinical Sciences and
³Center for Rhizosphere Biology, Colorado State University, Fort Collins, Colorado, USA

ABSTRACT Gut-associated lymphoid tissue maintains mucosal homeostasis by combating pathogens and inducing a state of hyporesponsiveness to food antigens and commensal bacteria. Dietary modulation of the intestinal immune environment represents a novel approach for enhancing protective responses against pathogens and inflammatory diseases. Dietary rice bran consists of bioactive components with disease-fighting properties. Therefore, we conducted a study to determine the effects of whole dietary rice bran intake on mucosal immune responses and beneficial gut microbes. Mice were fed a 10% rice bran diet for 28 days. Serum and fecal samples were collected throughout the study to assess total immunoglobulin A (IgA) concentrations. Tissue samples were collected for cellular immune phenotype analysis, and concentrations of native gut *Lactobacillus* spp. were enumerated in the fecal samples. We found that dietary rice bran induced an increase in total IgA locally and systemically. In addition, B lymphocytes in the Peyer's patches of mice fed rice bran displayed increased surface IgA expression compared with lymphocytes from control mice. Antigen-presenting cells were also influenced by rice bran, with a significant increase in myeloid dendritic cells residing in the lamina propria and mesenteric lymph nodes. Increased colonization of native *Lactobacillus* was observed in rice bran-fed mice compared with control mice. These findings suggest that rice bran-induced microbial changes may contribute to enhanced mucosal IgA responses, and we conclude that increased rice bran consumption represents a promising dietary intervention to modulate mucosal immunity for protection against enteric infections and induction of beneficial gut bacteria.

KEY WORDS: • antibody • dendritic cell • innate immunity • mucosal immunity • prebiotic

INTRODUCTION

RICE IS A STAPLE FOOD for a large portion of the world's population. Rice is typically polished to make white rice for reasons of stability and consumer acceptability. The removal of the bran results in the loss of many prebiotic components and beneficial nutrients, including various polyphenols, essential fatty acids, and numerous antioxidants.^{1–3} Dietary rice bran intake has been shown to aid in protection against gastrointestinal cancers,^{1,4,5} improve lipid metabolism in diabetic rats,⁶ and significantly lower cholesterol levels in humans^{7,8} and hamsters.⁹ Also, rice bran-derived prebiotics suppress inflammatory bowel disease^{10,11} by decreasing ulceration and inducing beneficial effects on the intestinal mucosa through the production of anti-inflammatory cytokines.

Prebiotics are defined as nondigestible food ingredients that have a beneficial effect through their selective metabolism in the intestinal tract.¹² Prebiotics promote the growth of beneficial bacteria, commonly referred to as probiotics.^{13,14} In

addition, the fermentation of prebiotic components via probiotic bacteria elicits the production of short-chain fatty acids. This results in the acidification of the colonic environment,^{15–17} which is detrimental to pathogenic bacteria and advantageous for colonic epithelial cells. Probiotic bacteria tend to be classified as noncolonizing members of the microbiota with the exception of *Lactobacillus*, found in low concentration in the intestines.¹⁸ The mechanisms by which probiotics benefit the host include strengthening of the mucosal surface,¹⁹ modulating the immune response, and antagonizing pathogens via competition or antimicrobial agents.²⁰

Changes in the gut microbiota have been shown to significantly affect the mucosal immune system,²¹ most notably through the increased production and secretion of immunoglobulin A (IgA).^{22,23} For example, mice administered oral *Lactobacillus acidophilus* were shown to have increased production of IgA by B cells in the Peyer's patches.²⁴ Also, probiotic bacteria have been associated with the enhancement of innate immunity through increased macrophage recruitment and phagocytic activity,^{13,25,26} as well as pro-inflammatory cytokine production.²⁷

Emerging evidence supports the beneficial effects of single rice bran components,²⁸ enzyme-treated rice bran,¹¹ and fermented rice bran^{3,10} on the immune system. Although these studies have uncovered important information, whole

Manuscript received 15 August 2011. Revision accepted 18 November 2011.

Address correspondence to: Elizabeth P. Ryan, Animal Cancer Center, Department of Clinical Sciences, Colorado State University, Fort Collins, CO 80523, E-mail: e.p.ryan@colostate.edu

dietary rice bran demonstrates greater feasibility and may be more practical for public health intervention in the developing and developed world when compared with extract preparations. Given the strong potential for dietary inclusion of whole rice bran as a functional food ingredient for chronic disease control and prevention,^{1,7,9,29} studies focused on the effects of dietary rice bran intake on the gut mucosal immune response and the microbiota are warranted. We investigated whether a 10% rice bran intake when compared with a control diet led to changes in cellular and humoral immune responses as well as alterations in the concentration of native gut *Lactobacillus* spp.

MATERIALS AND METHODS

Animals and feeding schedule

Specific pathogen-free 4–6-week-old female ICR mice were purchased from Harlan Laboratories (Indianapolis, IN, USA). Specific pathogen-free mice have a normal composition of intestinal bacteria and are devoid of known mouse pathogens.³⁰ All mice were provided water *ad libitum* and fed a maintenance diet (AIN-93M; Harlan Teklad, Madison, WI, USA) for 1 week prior to randomization. Mice were placed on either the control diet (AIN-93M) or the 10% rice bran diet for 28 days. The Institutional Animal Care and Use Committee at Colorado State University approved all protocols involving the animal experiments described in this study.

Diet composition

The Neptune rice variety was chosen based on its availability for rice grown in the United States. Control (AIN-93M) and rice bran-containing diets were formulated to match macronutrients across diet groups, and differences in total starch and fiber contents were balanced using purified diet components. The composition of rice bran-containing diet was calculated based on published reports^{31,32} that demonstrated chronic disease-fighting activity of rice bran and were stored at -20°C until fed to the mice. Diet formulations are shown in Table 1.

Sampling procedures

Mice were fed the experimental rice bran diet for 28 days. Fecal and serum samples were collected from all mice on day 0, 4, 7, 14, 18, 21, 25, and 28 following commencement of experimental diets. Mice were bled from the tail vein (50–75 μL per mouse), and serum samples were obtained using serum separator tubes (BD Biosciences, San Diego, CA, USA). Fecal extracts were made by using methods previously described.³³

Quantification of total IgA and *Lactobacillus*-specific IgA antibody

Total IgA antibody titers were assessed in mouse feces and serum by enzyme-linked immunosorbent assay (ELISA). Nunc-Immuno™ plates (Thermo Scientific, Waltham, MA, USA) were coated overnight at 4°C with purified rat anti-

TABLE 1. COMPOSITION OF EXPERIMENTAL DIETS

Component	Level (g/kg)	
	Control diet ^a	10% rice bran diet ^b
Casein	140.0	140.0
L-Cysteine	1.8	1.8
Cornstarch	465.7	422.7
Maltodextrin	155.0	155.0
Sucrose	100.0	100.3
Corn oil	40.0	19.0
Cellulose	50.0	29.0
Mineral Mix	35.0	35.0
Vitamin Mix	10.0	10.0
Choline bitartrate	2.5	2.5
TBHQ, antioxidant	0.008	0.008
Rice bran	—	100.0

^aControl diet (AIN-93M) produced by Harlan.

^b10% rice bran diet produced by Harlan using AIN-93M in addition to rice bran from Neptune rice variety, adjusted to match control diet in total fat, protein, and carbohydrates.

TBHQ, *tert*-butylhydroquinone.

mouse IgA (C10-3; BD) at a concentration of 5 $\mu\text{g/mL}$ in carbonate-bicarbonate buffer (pH 9.6). Nonspecific protein binding sites were blocked with phosphate-buffered saline (PBS) containing 3% bovine serum albumin (Sigma, St. Louis, MO, USA). Dilutions (1% bovine serum albumin) of the samples and standard were applied to the plate (50 μL per well) and incubated for 1.5 hours at room temperature. Plates were incubated with biotin rat anti-mouse IgA (C10-1; BD) at a dilution of 1:1,000 for 1 hour at room temperature. Next, plates were incubated with peroxidase-conjugated streptavidin (Jackson ImmunoResearch, West Grove, PA, USA) at a concentration of 1:1,500 for 20 minutes at room temperature. Plates were developed with 3,3',5,5'-tetramethylbenzidine (Sigma). The reaction was stopped using 1 *N* HCl, and the optical densities were read at 450 nm. Purified mouse IgA (eBioscience, San Diego, CA, USA) was used to create the standard curve for the ELISA.

The *Lactobacillus* IgA ELISA used a similar protocol as the total IgA ELISA. The *Lactobacillus* spp. used to coat the plate were isolated from fecal samples collected on day 18 following commencement of dietary rice bran. The lactobacilli was expanded in MRS broth prior to undergoing heat-killing as described previously.³⁴ ELISA plates were coated overnight at 4°C with 10^8 colony-forming units/mL heat-killed lactobacilli in carbonate-bicarbonate buffer. Nonspecific binding sites were blocked with PBS containing 5% nonfat dried milk. A positive control sample was obtained by vaccinating intraperitoneally an ICR mouse with the heat-killed lactobacilli and collecting serum after 14 days.

Tissue harvest and immune cell preparation

After the 28-day feeding period, mice were killed following anesthetization by intraperitoneal injection of ketamine (100 mg/kg) with xylazine (10 mg/kg). The mesenteric lymph nodes and Peyer's patches were collected, mechanically disrupted through a mesh screen, and treated with

NH₄Cl. Next, a 10-cm section of the intestine (ileum) was collected, and the intestinal lamina propria lymphocytes were prepared as previously described.³⁵ The intestinal tissue was disrupted through a mesh screen prior to being run through a density gradient (Optiprep, Accurate Chemical & Scientific Corp., Westbury, NY, USA). All cell preparations were resuspended in complete RPMI 1640 medium (Invitrogen, Carlsbad, CA, USA) containing 10% fetal bovine serum (Gemini Bio-Products, West Sacramento, CA, USA), 2 mM L-glutamine (Invitrogen), 1× nonessential amino acids (Invitrogen), 0.075% sodium bicarbonate (Fisher Scientific, Pittsburgh, PA, USA), 100 U/mL penicillin, and 100 µg/mL streptomycin (Invitrogen). Cells were kept on ice prior to immunostaining and analysis.

Flow cytometric analysis

Directly conjugated antibodies used for these analyses were purchased from eBioscience or BD Pharmingen (San Jose, CA, USA). The following antibodies were used in various combinations: anti-CD4 (GK1.5), anti-CD8b (H35-17.2), anti-CD5 (53-7.3), anti-CD27 (LG.7F9), anti-CD11c (N418), anti-CD11b (M1/70), anti-CD45R (clone RA3-6B2), and anti-mouse IgA (C10-1). Immunostaining was completed as described previously.³⁶ Analysis was performed with a Cyan ADP flow cytometer (Beckman Coulter, Fort Collins, CO, USA), and data were analyzed using FlowJo software (Tree Star, Ashland, OR, USA).

Fecal *Lactobacillus* levels

On day 0, 4, 7, 14, 18, 21, 25, and 28, following commencement of experimental diets, fecal samples were collected from all mice. Approximately five or six fresh fecal pellets (0.1 g) were collected per mouse and rehydrated in 1 mL of sterile PBS for 15 minutes. The samples were vortex-mixed, diluted in PBS, plated on MRS agar (enrichment agar for the genus *Lactobacillus*), and placed in a 37°C incubator containing 5% CO₂ for 48 hours. All colonies grown on MRS agar were confirmed as *Lactobacillus* spp. using real-time polymerase chain reaction as previously described.³⁷

Statistical analyses

Statistical analysis was performed using Prism version 5.0 software (GraphPad, La Jolla, CA, USA). Two-tailed nonparametric (Mann–Whitney) *t* tests were performed for comparisons between the two groups. A repeated-measures (mixed model) two-way analysis of variance with a Bonferroni *post hoc* test was performed to compare the two groups over time. Differences were considered statistically significant for $P < .05$ for all comparisons.

RESULTS

Dietary rice bran intake induces local and systemic IgA production

The effect of daily rice bran intake on the mucosal immune system was first investigated in mice for changes in

serum, intestinal, and fecal IgA. Control and 10% rice bran diets were fed to mice for 28 days, and the total IgA antibody levels were measured throughout this time period. The concentration of total IgA in the feces of the rice bran-fed mice (Fig. 1A) was significantly increased on day 18 and remained high through day 21 in comparison with mice on the control diet. A similar difference was detected systemically in the serum (Fig. 1B) with the total IgA concentration peaking at day 18.

We assessed the expression of IgA on the surface of intestinal B cells to elucidate the modulating IgA response following dietary rice bran intake. Increased expression of surface IgA indicates activation of local B cells as well as an

FIG. 1. Effect of dietary rice bran on local and systemic total immunoglobulin A (IgA) responses. A total IgA enzyme-linked immunosorbent assay was performed on serial dilutions of (A) fecal and (B) serum samples collected for 28 days. Antibody concentrations were determined through comparison with a standard control. Data are mean $\pm$ SEM values ($n = 12$ mice per group). Results were pooled from two independent experiments. Significant differences (***) were determined by a repeated-measures (mixed model) two-way analysis of variance followed by a Bonferroni *post hoc* test.

TABLE 2. INCREASED EXPRESSION OF SURFACE IMMUNOGLOBULIN A ON B220⁺ B CELLS IN PEYER'S PATCHES FOLLOWING 14 DAYS OF RICE BRAN DIET CONSUMPTION

Cell population	Lamina propria		Peyer's patches		Mesenteric lymph nodes	
	Control	10% RB	Control	10% RB	Control	10% RB
B220 ⁺ IgA ⁺ cells (%)	1.4±0.3	1.6±0.3	1.0±0.3	1.1±0.3	0.6±0.1	0.7±0.2
IgA (MFI) on B220 ⁺ B cells	193±36	194±51	98.7±23.1	166.7±19.7*	81.3±24	88.6±28

Data are mean±SEM values, pooled from two independent experiments ($n=10$).

* $P<.05$ as determined by comparing control diet versus 10% rice bran (RB) diet in each cell population using a nonparametric Mann-Whitney t test.

MFI, mean fluorescence intensity.

increased potential to respond to commensal and pathogenic bacteria in a T cell-independent fashion.²³ Mice were euthanized after 14 days of rice bran consumption, and the B220⁺ B cells were analyzed for surface IgA expression (Table 2). A significant difference was observed in the number of IgA molecules expressed on the B cell surface between control and rice bran-fed mice, determined by mean fluorescence intensity. These results are consistent with the idea that dietary rice bran induces mucosal immunity via systemic and local IgA production as well as increased activation of IgA⁺ B cells found in the Peyer's patches.

Increased mucosal CD11c⁺CD11b⁺ dendritic cells induced by rice bran

In order to determine whether dietary rice bran intake influences antigen-presenting cells as a mechanism for IgA induction, we examined cellular immune phenotypes previously shown to be required for IgA class switching.^{38,39} Following 14 days of daily 10% rice bran intake, mice were sacrificed, and the cellular responses were assessed in the lamina propria, Peyer's patches, and mesenteric lymph nodes (Table 3). The percentages of myeloid dendritic cells (CD11c⁺CD11b⁺) in both the lamina propria and the mesenteric lymph nodes were significantly increased in mice fed the 10% rice bran diet in comparison with the mice

on control diet. No significant differences were detected in T or B cell populations (Table 3). These data suggested that rice bran-mediated induction of dendritic cell recruitment into the mucosal tissues resulted in increased antigen presentation and subsequent IgA production.

Dietary rice bran increased *Lactobacillus* colonization

Given the associations of probiotic bacteria with the modulation of the mucosal immune system,²¹ we next determined the titers of the native gut *Lactobacillus* spp. in the rice bran-fed mice compared with the control group. *Lactobacillus* was chosen for its aerobic growth conditions as well as its ability to protect against a variety of infectious diseases.⁴⁰ For these experiments, we processed and plated fresh fecal pellets for the presence of *Lactobacillus* spp. over a 28-day period. On days 11, 18, 21, and 25 the numbers of *Lactobacillus* spp. in the feces were significantly higher from the rice bran-fed mice in comparison with a more cyclical pattern seen in the mice fed control diet (Fig. 2). The nearly 500% increase in numbers of *Lactobacillus* spp. (in colony-forming units per gram of feces) became apparent on day 11 and remained significantly high for 14 days. Because of the ability of multiple *Lactobacillus* species to induce protection against intestinal pathogens,⁴⁰⁻⁴³ real-time polymerase chain reaction was performed to confirm a greater than 900% increase in DNA levels from the genus

TABLE 3. INFLUENCE OF DIETARY RICE BRAN ON CELLULAR POPULATIONS IN THE INTESTINAL IMMUNE COMPARTMENTS FOLLOWING 14 DAYS OF DIET CONSUMPTION

Cell population (%)	Lamina propria		Peyer's patches		Mesenteric lymph nodes	
	Control	10% RB	Control	10% RB	Control	10% RB
T cells						
CD4 ⁺	7.4±0.8	9.6±1.0	20.3±1.5	19.7±1.9	66.8±1.7	67.6±1.5
CD8 ⁺	16.5±3.3	18.4±3.3	6.3±0.8	4.5±0.4	17.1±0.6	16.3±0.9
B cells						
B220 ⁺	33.7±5.9	31.3±3.1	66.1±3.6	66.4±4.1	10.8±3.6	20.4±0.9
B220 ⁺ CD5 ⁺	0.9±0.1	1.3±0.1	3.3±0.7	3.8±0.8	7.4±1.5	8.7±0.8
B220 ⁺ CD27 ⁺	0.8±0.2	1.0±0.3	0.2±0.04	0.2±0.09	0.3±0.05	0.3±0.05
CD11c ⁺ CD11b ⁺ DCs	6.8±0.9	11.1±0.9**	1.1±0.1	1.2±0.1	0.6±0.1	0.7±0.04*

Data are mean±SEM values, pooled from two independent experiments ($n=10$).

* $P<.05$, ** $P<.01$ as determined by comparing control diet versus 10% RB diet in each cell population using a nonparametric Mann-Whitney t test.

DC, dendritic cell.

Lactobacillus between day 0 and day 14 in the mice fed the rice bran diet (data not shown). Both the control and rice bran diets did not contain any lactobacilli. Next, we determined if the increased IgA titers were *Lactobacillus*-specific. A *Lactobacillus* IgA ELISA was performed and revealed low to undetectable *Lactobacillus*-specific IgA (data not shown). A positive control was in place to ensure the reliability and sensitivity of the *Lactobacillus* IgA ELISA as described in Materials and Methods. Therefore, these results indicate a potential role for dietary rice bran in modulating the intestinal microbiota through eliciting increased concentrations of beneficial bacteria.

DISCUSSION

The unique health-promoting properties and chemical composition of rice bran make it a promising candidate for dietary supplementation, for nutritional therapy, and for prevention of chronic disease.^{1,2,4-9,28,44-47} In this study we found the effects of whole dietary rice bran intake on the mucosal immune system to be the induction of IgA (Fig. 1) and the enhancement of the innate immune response (Table 3). In addition, the ability of dietary rice bran to promote increased intestinal colonization of native *Lactobacillus* spp. (Fig. 2) highlights a novel role for rice bran prebiotic components to influence the intestinal microbiota.^{2,48}

Understanding the effect of dietary rice bran on the innate immune system and antigen presentation is crucial to elucidating the mechanisms by which rice bran may induce protective responses at mucosal surfaces. The increased percentages of myeloid dendritic cells in both the lamina propria and the mesenteric lymph nodes following rice bran consumption speak to the targeted effect of rice bran on the key cells involved in shaping an immune response. Intestinal dendritic cells have the unique ability to induce IgA production from B cells through secretion of retinoic acid, interleukin-5, and interleukin-6.⁴⁹ Therefore, the enhanced

dendritic cell population can help in the initiation of an adaptive immune response through the activation of B cells and subsequent IgA class-switching. Also, the increased presence of dendritic cells in the lamina propria enhances the mucosal innate immune response at the site where pathogens invade and reveals a potential protective mechanism induced by dietary rice bran.

Evidence for the modulation of the mucosal IgA response associated with the consumption of dietary components has been limited to fructooligosaccharides and pectin.^{50,51} Consistent with these findings, this study demonstrated the ability of dietary rice bran to induce increased IgA concentrations systemically and locally. Various bioactive and prebiotic components present in rice bran are hypothesized to be involved in the enhancement of immunity. These phytochemicals include, but are not limited to, γ -oryzanol, polyphenols, and fatty acids, as well as some essential amino acids and micronutrients.² Understanding the mechanism of IgA induction is of considerable importance when evaluating the dietary capacity of rice bran to elicit protection against enteric infections. Current evidence suggests that the majority of IgA molecules in the gut are induced by commensal bacteria and that there is a role for this transient population to induce a specific IgA response.^{23,52-57} A recent study performed by Hapfelmeier *et al.*⁵⁴ confirms the role of commensal bacteria, but also describes the specific IgA response to be less robust and have a slower onset, consistent with our findings. Another relevant finding by Macpherson *et al.*²³ showed induction of commensal-specific IgA antibodies to be mostly T cell independent but dependent largely on the B1 peritoneal B cells, a potential angle that should be evaluated in rice bran-fed mice.

Based on the emerging evidence for commensal involvement in shaping the mucosal IgA population, the beneficial effect of dietary rice bran on the native gut *Lactobacillus* spp. was shown to be a potential mechanism of immune modulation (Fig. 2). The antibody-enhancing ability of *Lactobacillus* was described previously in a study showing increased production of rotavirus-specific antibodies when fermented milk was administered during the acute phase of a rotavirus infection.⁵⁸ The low *Lactobacillus*-specific IgA antibodies in our model may speak to the unique ability of rice bran to enhance antibody responses specific for other bacteria, or it may reflect the fact that dietary rice bran contains growth substrates for resident bacteria. As a result, the specificity of the increased IgA would be heterogeneous, making the *Lactobacillus* titer too low for detection by ELISA. We hypothesize that rice bran-induced modulation of multiple commensal bacteria resulted in increased luminal IgA concentrations in the intestine. Future research directions should include elucidating the effect of dietary rice bran on the gastrointestinal microbiota using 454 pyrosequencing or other high-throughput microbiome analytical approaches.

The ability of dietary rice bran to promote the growth of native gut *Lactobacillus* spp. and enhance mucosal immune cell populations offers numerous health-promoting and disease-fighting possibilities. For example, the potential use

FIG. 2. Effect of dietary RB on fecal titers of *Lactobacillus*. Fresh fecal pellets were collected, processed, and plated on MRS agar every 3–4 days in order to determine the Log₁₀ *Lactobacillus* titer per gram of feces. Data are mean $\pm$ SEM values ($n = 10$ mice per group). Results were pooled from two independent experiments. Significant differences ($*P < .05$, $***P < .001$) were determined by a repeated-measures (mixed model) two-way analysis of variance followed by a Bonferroni *post hoc* test.

of rice bran as a dietary vaccine adjuvant is highlighted by the recent application of *Lactobacillus fermentum* in an intramuscular influenza vaccine.⁵⁹ Also, the beneficial effects of rice bran on inflammatory diseases such as inflammatory bowel disease^{10,11} and type 1 diabetes^{6,60} emphasize a unique research avenue to study the immune-enhancing effects of dietary rice bran on these nutritionally relevant diseases. In summary, the ability of whole dietary rice bran to modulate the immune system as well as promote the growth of native gut *Lactobacillus* spp. holds great promise in protection against enteric pathogens and modulation of chronic inflammatory diseases.

ACKNOWLEDGMENTS

We would like to thank Dr. Anna McClung from the U.S. Department of Agriculture rice research unit for providing rice bran from the single Neptune variety. This work was supported by a Grand Explorations in Global Health grant (OPP1015267) from the Bill and Melinda Gates Foundation and the Shipley Foundation and a grant from the National Institutes of Health (R03CA150070).

AUTHOR DISCLOSURE STATEMENT

The authors disclose no conflicts of interest.

REFERENCES

- Hudson EA, Dinh PA, Kokubun T, Simmonds MSJ, Gescher A: Characterization of potentially chemopreventive phenols in extracts of brown rice that inhibit the growth of human breast and colon cancer cells. *Cancer Epidemiol Biomarkers Prev* 2000; 9:1163–1170.
- Ryan EP: Bioactive food components and health properties of rice bran. *J Am Vet Med Assoc* 2011;238:593–600.
- Ryan EP, Heuberger AL, Weir TL, Barnett B, Broeckling CD, Prenni JE: Rice bran fermented with *Saccharomyces boulardii* generates novel metabolite profiles with bioactivity. *J Agric Food Chem* 2011;59:1862–1870.
- Cai H, Al-Fayez M, Tunstall RG, et al.: The rice bran constituent tricin potentially inhibits cyclooxygenase enzymes and interferes with intestinal carcinogenesis in ApcMin mice. *Mol Cancer Ther* 2005;4:1287–1292.
- Norazalina S, Norhaizan ME, Hairuszah I, Norashareena MS: Anticarcinogenic efficacy of phytic acid extracted from rice bran on azoxymethane-induced colon carcinogenesis in rats. *Exp Toxicol Pathol* 2010;62:259–268.
- Chou TW, Ma CY, Cheng HH, Chen YY, Lai MH: A rice bran oil diet improves lipid abnormalities and suppress hyperinsulinemic responses in rats with streptozotocin/nicotinamide-induced type 2 diabetes. *J Clin Biochem Nutr* 2009;45:29–36.
- Gerhardt AL, Gallo NB: Full-fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *J Nutr* 1998;128:865–869.
- Hegsted M, Windhauser MM, Morris SK, Lester SB: Stabilized rice bran and oat bran lower cholesterol in humans. *Nutr Res* 1993;13:387–398.
- Kahlon TS, Chow FI, Sayre RN, Betschart AA: Cholesterol-lowering in hamsters fed rice bran at various levels, defatted rice bran and rice bran oil. *J Nutr* 1992;122:513–519.
- Kataoka K, Ogasa S, Kuwahara T, et al.: Inhibitory effects of fermented brown rice on induction of acute colitis by dextran sulfate sodium in rats. *Dig Dis Sci* 2008;53:1601–1608.
- Komiyama Y, Andoh A, Fujiwara D, et al.: New prebiotics from rice bran ameliorate inflammation in murine colitis models through the modulation of intestinal homeostasis and the mucosal immune system. *Scand J Gastroenterol* 2011;46:40–52.
- Gibson GR, Probert HM, Loo JV, Rastall RA, Roberfroid MB: Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutr Res Rev* 2004;17:259–275.
- Isolauri E, Sütas Y, Kankaanpää P, Arvilommi H, Salminen S: Probiotics: effects on immunity. *Am J Clin Nutr* 2001;73:444S–450S.
- Sanders ME: Probiotics: considerations for human health. *Nutr Rev* 2003;61:91–99.
- Cummings JH, Macfarlane GT: Role of intestinal bacteria in nutrient metabolism. *Clin Nutr* 1997;16:3–11.
- Reilly KJ, Rombeau JL: Metabolism and potential clinical applications of short-chain fatty acids. *Clin Nutr* 1993;12:S97–S105.
- van Hylckama Vlieg JET, Veiga P, Zhang C, Derrien M, Zhao L: Impact of microbial transformation of food on health—from fermented foods to fermentation in the gastro-intestinal tract. *Curr Opin Biotechnol* 2011;22:211–219.
- Makarova K, Slesarev A, Wolf Y, et al.: Comparative genomics of the lactic acid bacteria. *Proc Natl Acad Sci USA* 2006;103:15611–15616.
- Ulluwishewa D, Anderson RC, McNabb WC, Moughan PJ, Wells JM, Roy NC: Regulation of tight junction permeability by intestinal bacteria and dietary components. *J Nutr* 2011;141:769–776.
- Corr SC, Li Y, Riedel CU, O'Toole PW, Hill C, Gahan CGM: Bacteriocin production as a mechanism for the anti-infective activity of *Lactobacillus salivarius* UCC118. *Proc Natl Acad Sci USA* 2007;104:7617–7621.
- Forsythe P, Bienenstock J: Immunomodulation by commensal and probiotic bacteria. *Immunol Invest* 2010;39:429–448.
- He B, Xu W, Santini PA, et al.: Intestinal bacteria trigger T cell-independent immunoglobulin A2 class switching by inducing epithelial-cell secretion of the cytokine APRIL. *Immunity* 2007;26:812–826.
- Macpherson AJ, Gatto D, Sainsbury E, Harriman GR, Hengartner H, Zinkernagel RM: A primitive T cell-independent mechanism of intestinal mucosal IgA responses to commensal bacteria. *Science* 2000;288:2222–2226.
- Torii A, Torii S, Fujiwara S, Tanaka H, Inagaki N, Nagai H: *Lactobacillus acidophilus* strain L-92 regulates the production of Th1 cytokine as well as Th2 cytokines. *Allergol Int* 2007;56:293–301.
- Perdigon G, de Macias ME, Alvarez S, Oliver G, de Ruiz Holgado AA: Effect of perorally administered lactobacilli on macrophage activation in mice. *Infect Immun* 1986;53:404–410.
- Perdigon G, de Macias ME, Alvarez S, Oliver G, de Ruiz Holgado AP: Systemic augmentation of the immune response in mice by feeding fermented milks with *Lactobacillus casei* and *Lactobacillus acidophilus*. *Immunology* 1988;63:17–23.
- Galdeano CM, Perdigon G: The probiotic bacterium *Lactobacillus casei* induces activation of the gut mucosal immune system through innate immunity. *Clin Vaccine Immunol* 2006;13:219–226.
- Sierra S, Lara-Villoslada F, Olivares M, Jimenez J, Boza J, Xaus J: Increased immune response in mice consuming rice bran oil. *Eur J Nutr* 2005;44:509–516.

29. Jariwalla RJ: Rice-bran products: phytonutrients with potential applications in preventive and clinical medicine. *Drugs Exp Clin Res* 2001;27:17–26.
30. Macpherson AJ, Harris NL: Interactions between commensal intestinal bacteria and the immune system. *Nat Rev Immunol* 2004;4:478–485.
31. Bird AR, Hayakawa T, Marsono Y, *et al.*: Coarse brown rice increases fecal and large bowel short-chain fatty acids and starch but lowers calcium in the large bowel of pigs. *J Nutr* 2000;130:1780–1787.
32. Cara L, Dubois C, Borel P, *et al.*: Effects of oat bran, rice bran, wheat fiber, and wheat germ on postprandial lipemia in healthy adults. *Am J Clin Nutr* 1992;55:81–88.
33. Haneberg B, Kendall D, Amerongen HM, Apter FM, Kraehenbuhl JP, Neutra MR: Induction of specific immunoglobulin A in the small intestine, colon-rectum, and vagina measured by a new method for collection of secretions from local mucosal surfaces. *Infect Immun* 1994;62:15–23.
34. Lauw FN, Simpson AJ, Prins JM, *et al.*: The CXC chemokines gamma interferon (IFN-gamma)-inducible protein 10 and monokine induced by IFN-gamma are released during severe melioidosis. *Infect Immun* 2000;68:3888–3893.
35. Vajdy M, Kosco-Vilbois MH, Kopf M, Kohler G, Lycke N: Impaired mucosal immune responses in interleukin 4-targeted mice. *J Exp Med* 1995;181:41–53.
36. Bosio CM, Dow SW: *Francisella tularensis* induces aberrant activation of pulmonary dendritic cells. *J Immunol* 2005;175:6792–6801.
37. Malinen E, Rinttila T, Kajander K, *et al.*: Analysis of the fecal microbiota of irritable bowel syndrome patients and healthy controls with real-time PCR. *Am J Gastroenterol* 2005;100:373–382.
38. Fayette J, Dubois B, Vandenabeele S, *et al.*: Human dendritic cells skew isotype switching of CD40-activated naive B cells towards IgA1 and IgA2. *J Exp Med* 1997;185:1909–1918.
39. Macpherson AJ: IgA adaptation to the presence of commensal bacteria in the intestine. In: *Current Topics in Microbiology and Immunology*, Vol. 308: *Gut-Associated Lymphoid Tissues* (Honjo T, Melchers F, eds.). Springer, Berlin, 2006, pp. 117–136.
40. Casas IA, Dobrogosz WJ: Validation of the probiotic concept: *Lactobacillus reuteri* confers broad-spectrum protection against disease in humans and animals. *Microbial Ecol Health Dis* 2000;12:247–285.
41. Gill H, Shu Q, Lin H, Rutherford K, Cross M: Protection against translocating *Salmonella typhimurium* infection in mice by feeding the immuno-enhancing probiotic *Lactobacillus rhamnosus* strain HN001. *Med Microbiol Immunol* 2001;190:97–104.
42. Oksanen PJ, Salminen S, Saxelin M, *et al.*: Prevention of travellers diarrhoea by *Lactobacillus GG*. *Ann Med* 1990;22:53–56.
43. Valladares R, Sankar D, Li N, *et al.*: *Lactobacillus johnsonii* N6.2 mitigates the development of type 1 diabetes in BB-DP rats. *PLoS One* 2010;5:e10507.
44. Cholujova D, Jakubikova J, Sedlak J: BioBran-augmented maturation of human monocyte-derived dendritic cells. *Neoplasma* 2009;56:89–95.
45. Chotimarkorn C, Benjakul S, Silalai N: Antioxidant components and properties of five long-grained rice bran extracts from commercial available cultivars in Thailand. *Food Chem* 2008;111:636–641.
46. Ghoneum M, Matsuura M: Augmentation of macrophage phagocytosis by modified arabinoxylan rice bran (MGN-3/biobran). *Int J Immunopathol Pharmacol* 2004;17:283–292.
47. Ghoneum M, Matsuura M, Gollapudi S: Modified arabinoxylan rice bran (MGN3/Biobran) enhances intracellular killing of microbes by human phagocytic cells in vitro. *Int J Immunopathol Pharmacol* 2008;21:87–95.
48. Heuberger AL, Lewis MR, Chen MH, Brick MA, Leach JE, Ryan EP: Metabolomic and functional genomic analyses reveal varietal differences in bioactive compounds of cooked rice. *PLoS One* 2010;5:e12915.
49. Mora JR, Iwata M, Eksteen B, *et al.*: Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science* 2006;314:1157–1160.
50. Hosono A, Ozawa A, Kato R, *et al.*: Dietary fructooligosaccharides induce immunoregulation of intestinal IgA secretion by murine Peyer's patch cells. *Biosci Biotechnol Biochem* 2003;67:758–764.
51. Lim BO, Yamada K, Nonaka M, Kuramoto Y, Hung P, Sugano M: Dietary fibers modulate indices of intestinal immune function in rats. *J Nutr* 1997;127:663–667.
52. Bos N, Bun J, Popma S, *et al.*: Monoclonal immunoglobulin A derived from peritoneal B cells is encoded by both germ line and somatically mutated VH genes and is reactive with commensal bacteria. *Infect Immun* 1996;64:616–623.
53. Bos NA, Jiang H-Q, Cebra JJ: T cell control of the gut IgA response against commensal bacteria. *Gut* 2001;48:762–764.
54. Hapfelmeier S, Lawson MAE, Slack E, *et al.*: Reversible microbial colonization of germ-free mice reveals the dynamics of IgA immune responses. *Science* 2010;328:1705–1709.
55. Macpherson AJ, Uhr T: Induction of protective IgA by intestinal dendritic cells carrying commensal bacteria. *Science* 2004;303:1662–1665.
56. Shroff K, Meslin K, Cebra J: Commensal enteric bacteria engender a self-limiting humoral mucosal immune response while permanently colonizing the gut. *Infect Immun* 1995;63:3904–3913.
57. Talham GL, Jiang H-Q, Bos NA, Cebra JJ: Segmented filamentous bacteria are potent stimuli of a physiologically normal state of the murine gut mucosal immune system. *Infect Immun* 1999;67:1992–2000.
58. Kaila M, Isolauri E, Soppi E, Virtanen E, Laine S, Arvilommi H: Enhancement of the circulating antibody secreting cell response in human diarrhea by a human *Lactobacillus* strain. *Pediatr Res* 1992;32:141–144.
59. Olivares M, Díaz-Ropero MP, Sierra S, *et al.*: Oral intake of *Lactobacillus fermentum* CECT5716 enhances the effects of influenza vaccination. *Nutrition* 2007;23:254–260.
60. Kim TH, Kim EK, Lee MS, *et al.*: Intake of brown rice lees reduces waist circumference and improves metabolic parameters in type 2 diabetes. *Nutr Res* 2011;31:131–138.