Dinucleotide repeat polymorphism at the human cysteine-proteinase inhibitor pseudogene (CSTP1)

M.H.Polymeropoulos, H.Xiao, D.S.Rath and C.R.Merrill
National Institute of Mental Health Neuroscience Center,
St Elizabeths Hospital, Room 131, 2700 Martin Luther
King Avenue, Washington, DC 20032, USA

Source/Description: The polymorphic (GT)_n repeat begins at base pair 2533 in intron A of the human cysteine-proteinase inhibitor pseudogene (CSTP1) on chromosome 20 (1). The polymorphism can be typed using the polymerase chain reaction (PCR) as described previously (2). The predicted length of the amplified sequence was 129 bp.

Primer Sequences: ATGTGACTGATGTGGGTCAG (GT strand); CATCTGCACTCATGCTCCAT (CA strand).

Frequency: Estimated from 50 chromosomes of unrelated individuals. Observed heterozygosity = 61%. PIC = 0.58.

Allele (bp)	Frequency	Allele (bp)	Frequency
A1 141	0.04	A5 129	0.20
A2 137	0.02	A6 127	0.06
A3 133	0.02	A7 125	0.58
A4 131	0.02	A8 123	0.06

Mendelian Inheritance: Co-dominant segregation was observed in two informative families.

Chromosomal Localization: CSTP1 has been assigned to chromosome 20 (3).

Other Comments: The PCR reaction was performed on 80 ng of genomic DNA using 100 pmoles of each oligonucleotide primer. The samples were processed as described (4) except that the denaturation cycle at 94°C was extended to 1.4 minutes. The dinucleotide repeat was based on a (GT)₁₅ sequence.

References: 1) Saitoh, E. *et al.* (1987) *Gene* **61**, 329–338. 2) Weber, J.L. and May, P.E. (1989) *Am. J. Hum. Genet.* **44**, 388–396. 3) Saitoh, E. *et al.* (1989) *Biochem. Biophys. Res. Commun.* **162**, 1324–1331. 4) Weber, J.L. *et al.* (1990) *Nucl. Acids Res.* **18**, 4637.

HinfI polymorphism within the 3' untranslated region of the candidate Wilms tumour gene

Paul R.Hoban* and Anna M.Kelsey
Department of Pathology, Royal Manchester Childrens
Hospital (RMCH), Pendlebury, Manchester M27 1HA, UK

Source/Description: Oligonucleotide primers were used to amplify a 952 bp genomic fragment from within the 3' untranslated region of the cDNA from the candidate Wilms tumour gene (1).

PCR Primers:

5' GCCTGGAAGAGTTGGTCTCT 3'

5' ACACAGTAATTTCAAGCAACGG 3'.

Method: 1 µg of genomic DNA was amplified using 0.1 µg of each primer in a total volume of 100 µl containing 1×taq polymerase buffer (BCL) and 250 µM dNTP's. Samples were heated at 97°C for 5 mins, following which 2 units of taq polymerase were added (BCL). Reactions were cycled 30 times at 94°C for 1 min, 60°C for 1 min and 74°C for 2.5 mins. Restrictions were analysed by electrophoresis through a 2% agarose gel.

Polymorphism: HinfI restriction reveals a two allele system with constant bands at 382 bp, 292 bp and 40 bp. Allele 1 is 238 bp and allele 2 is 133 bp and 105 bp.

No Polymorphism: Restriction with AluI, CfoI, HpaII, HaeIII, RsaI and TaqI, of a panel of 10 unrelated individuals does not reveal any polymorphism.

Frequency: Determined from 67 unrelated individuals.

Allele 1 (A1) — 0.66

Allele 2 (A2) — 0.34

Chromosomal Location: 11p13.

Mendelian Inheritance: Codominant segregation was observed in three families (19 individuals).

Acknowledgement: This work was supported by the Leukaemia Research Fund, RMCH, Pendlebury.

Reference: Call, K.M. *et al.* (1990) *Cell* **60**, 509–520.

* To whom correspondence should be addressed at CRC Department of Cancer Genetics, Paterson Institute for Cancer Research, Christie Hospital, Manchester M20 9BX, UK