Published in final edited form as:

J Cell Biochem. 2012 April ; 113(4): 1426–1436. doi:10.1002/jcb.24017.

Continuously Generated H₂O₂ Stimulates the Proliferation and Osteoblastic Differentiation of Human Periodontal Ligament Fibroblasts

Youngji Choe^{1,†}, Ji-Yeon Yu^{1,†}, Young-Ok Son², Seung-Moon Park³, Jong-Ghee Kim¹, Xianglin Shi², and Jeong-Chae Lee^{1,2,*}

¹Institute of Oral Biosciences (BK21 program) and School of Dentistry, Chonbuk National University, Jeonju 561-756, South Korea

²Graduate Center for Toxicology, School of Medicine, University of Kentucky, Lexington, KY 40536-0305, USA

³Division of Biotechnology, Chonbuk National University, Iksan 570-752, South Korea

Abstract

Numerous studies have shown that hydrogen peroxide (H₂O₂) inhibits proliferation and osteoblastic differentiation in bone-like cells. Human periodontal ligament fibroblasts (PLF) are capable of differentiating into osteoblasts and are exposed to oxidative stress during periodontal inflammation. However, the cellular responses of PLF to H₂O₂ have not been identified. In this study, we examined how H₂O₂ affects the viability and proliferation of PLF by exposing the cells to glucose oxidase (GO) or direct addition of H₂O₂. We also explored the effects of GO on the osteoblastic differentiation of PLF and the mechanisms involved. The viability and proliferation in PLF were increased with the addition of 10 mU/ml GO but not by volumes greater than 15 mU/ml or by H₂O₂ itself. GO-stimulated DNA synthesis was correlated with the increase in cyclin E protein levels in the cells. Osteoblastic differentiation of PLF was also augmented by combined treatment with GO, as evidenced by the increases in alkaline phosphatase activity, mineralization, collagen synthesis, and osteocalcin content in the cells. The inductions of runt-related transcription factor 2 and osterix mRNA and proteins were further increased in PLF incubated in combination with GO compared to those in untreated cells. These results demonstrate that the continuous presence of H₂O₂ stimulates the proliferation of PLF and augments their potential to differentiate into osteoblasts through the up-regulation of bone-specific transcription factors. Collectively, we suggest that H₂O₂ may elicit the functions of PLF in maintaining the dimensions of the periodontal ligament and in mediating a balanced metabolism in alveolar bone.

Keywords

PERIODONTAL LIGAMENT FIBROBLASTS; GLUCOSE OXIDASE; HYDROGEN PEROXIDE; PROLIFERATION; OSTEOBLASTIC DIFFERENTIATION

INTRODUCTION

The human periodontal ligament (PDL) is a band of fibrous connective tissue located between the tooth-root cementum and the alveolar bone. The PDL consists of a

*Correspondence to: Prof. Jeong-Chae Lee, Department of Orthodontics and Institute of Oral Biosciences, Chonbuk National University, Jeonju 561-756, South Korea, leejc88@jbnu.ac.kr.

[†]They are equally contributed to this work.

heterogeneous cell population, where fibroblasts are the predominant cell type (Bordin et al., 1984; McCulloch and Bordin, 1991). PDL plays important roles in maintaining the stability and function of teeth. It has been suggested that PDL fibroblasts (PLF) are a source of osteoblasts required for alveolar bone remodeling by physiological or mechanical stimuli (Roberts et al., 1982). PLFs are naturally osteogenic and are able to differentiate into osteoblasts (Heo et al., 2010). Numerous studies also demonstrate the potential of PLFs in regulating both osteoblastic and osteoclastic differentiation within periodontal tissue [Kook et al., 2009 and 2011; Wattanaroonwong et al., 2011].

Cellular oxidative stress occurs if reactive oxygen species (ROS) are overproduced or if their removal is reduced. Persistent and prolonged oxidative stress causes various pathological disorders such as stroke, heart attack, aging, and several degenerative diseases [Buttke and Sandstrom, 1994; Chandra et al., 2000]. ROS can be also produced continuously by periodontal-pathogenic bacteria or their by-products. ROS causes oxidative damage of periodontal tissue and cells and eventually mediates periodontitis [Ara et al., 2009]. There are also reports supporting the relationship between the antioxidant status in plasma and the risk of periodontitis [Chapple et al., 2007; Kokeguchi and Yamamoto, 2009]. Chaves Neto et al. (2011) demonstrated that oxidative stress inhibited the proliferation of bone cells, and this result was facilitated when the cells were incubated with osteoblast-inducing media. These reports indicate that the processes required for alveolar bone remodeling are affected by intracellular redox states. Furthermore, the modes of action of PLF in response to ROS are quite important because of their potentials to regulate osteoblastic and osteoclastic activations.

Hydrogen peroxide (H_2O_2) is generated by almost all sources of oxidative stress and can penetrate cellular membranes [Forman and Torres, 2001; Nordberg and Arner, 2001]. H_2O_2 is also widely used in dental clinics to bleach teeth, enhance gingival healing, and diminish bacterial populations in dental plaque [Tipton et al., 1995]. Many studies have shown the susceptibility of bone cells to H_2O_2 , where both the proliferation and differentiation of osteoblasts were sensitively suppressed by exposure to this agent [Choi et al., 2009; Chaves Neto et al., 2011; Kim et al., 2010]. However, it is important to note that cellular responses to H_2O_2 can differ depending on the type of cells and the concentration of H_2O_2 . In lymphoma cells, H_2O_2 treatment induced growth inhibition and apoptosis [Son et al., 2009], whereas this same treatment stimulated proliferation in several other cell types [Herbert et al., 1996; Timblin et al., 1995]. Exposure of BHK-21 fibroblasts to a low H_2O_2 level (1 μM) stimulated proliferation, but treatment with 100 μM H_2O_2 induced growth inhibition and apoptotic cell death in the cells [Burdon et al., 1996]. We previously reported on the dual modes of action of H_2O_2 according to the methods of exposure, i.e., temporal versus continuous presence of this agent in the cultures [Son et al., 2009]. Furthermore, glucose oxidase (GO) that continuously generates H_2O_2 at relatively low concentrations significantly increased the proliferation rate in bovine aortic endothelial cells, but a direct addition of H_2O_2 inhibited the proliferation of the cells [Ruiz-Ginés et al., 2000].

Despite the critical roles of PLF in the process of alveolar bone remodeling and in the maintenance of tooth function and stability, little information on the mode of action of PLF in response to H_2O_2 is available. Therefore, we examined the viability and proliferation of PLF after exposure to GO or to H_2O_2 itself. We also explored the effects of H_2O_2 on osteoblastic differentiation of the cells. In addition, we aimed to determine the mechanisms by which H_2O_2 affects the proliferation and differentiation of PLF. In the present study, we demonstrated for the first time that continuously generated H_2O_2 stimulates both the proliferation rates and osteoblastic differentiation in PLFs depending on the concentrations exposed.

MATERIALS AND METHODS

CHEMICALS AND LABORATORY WARES

Fetal bovine serum (FBS) was purchased from Gibco-BRL (Gaithersburg, MD, USA). The primary antibodies specific for actin, runt-related transcription factor-2 (Runx2), osterix, osteopontin, cyclins, and cyclin-dependent kinase-2 (CDK2), and the secondary goat-anti rabbit antibody were supplied by Santa Cruz Biotechnology (Santa Cruz, CA, USA). Unless otherwise specified, the other chemicals and laboratory items were purchased from Sigma Chemical Co. (St. Louis, MO, USA) and Falcon Labware (Becton-Dickinson, Franklin Lakes, NJ, USA), respectively.

CELL CULTURES AND H₂O₂ EXPOSURE

PLFs were obtained from healthy male volunteers aged 20 to 30 years and cultured according to methods described elsewhere with slight modifications [Kook et al., 2009]. All the donors gave written informed consent for use of their tissues. This study was approved by the Ethical Committee of Chonbuk National University Hospital. Here, single cell suspensions of PLFs were incubated in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS and antibiotics (100 IU/ml penicillin G and 100 µg/ml streptomycin) in 100-mm culture dishes. The cultures were switched to a fresh batch of medium every 3 days. When PLFs reached > 90% confluence, the cells were suspended in the culture medium at a density of 5×10^5 /ml and then seeded in 6-well and 96-well flat-bottomed plates with 2 ml and 200 µl per well, respectively. When the cells reached 70–80% confluence, the medium was changed to a fresh culture medium to proliferate or to differentiate into osteoblasts and exposed to various concentrations of GO or H₂O₂. In this study, 10 nM dexamethasone, 50 µM ascorbic acid, and 20 mM β-glycerophosphate, named DAG, were used to induce osteoblastic differentiation of PLFs. All the experiments were performed using fibromodulin-positive PLFs (> 95%) at passages 4 through 8. In the present study, PLFs were exposed to GO ranging from 0 to 50 mU/ml or H₂O₂ at various doses (10 µM to 2 mM) through its direct addition into the cultures.

H₂O₂ DETERMINATION

The levels of H₂O₂ in the culture medium were determined using Amplex[®] Red Hydrogen Peroxide Assay Kit (Molecular Probes, Inc., Eugene, OR, USA) according to the manufacturer's instruction. In brief, PLFs suspended in DMEM without phenol red were divided to culture plates in the presence and absence of DAG. The culture supernatants (50 µl/sample) were collected at various times after GO exposure and mixed with the same volume of the Red reagent in 96-multiwell plates. After incubation for 30 min at room temperature, horseradish peroxidase (0.1 U/ml) was added into the mixtures. Finally, the absorbance was measured at 560 nm using a microplate reader (Molecular Devices, Sunnyvale, CA, USA).

MEASUREMENT OF CELL VIABILITY AND DNA SYNTHESIS

Cell viability was determined using water-soluble tetrazolium salt (WST)-8 reagent. Briefly, GO- or H₂O₂-exposed cultures were treated with WST-8 reagent at various times after exposure. Following incubation for an additional 4 h, absorbance was measured at 450 nm using a microplate reader (Molecular Devices). The level of DNA synthesis in the cells was measured by adding 0.5 µCi of ³H-thymidine deoxyribose (TdR; Amersham Pharmacia Biotech Inc., Piscataway, NJ, USA) to each well of the 96-multiwell plates during the 12 h prior to cell harvesting. After collection of the cells using a harvester (Inotech Inc., Dietikon, Switzerland), beta emission from the ³H-TdR-incorporated cells was measured for 1 min using a liquid scintillation counter (Packard Instrument Co., Downers Grove, IL, USA).

ASSAY OF CELLULAR ANTIOXIDANT ACTIVITIES

PLFs were exposed to increasing GO concentrations for 24 h and then adjusted to the analyses of intracellular antioxidant activities such as superoxide dismutase (SOD), catalase, and reduced glutathione (GSH). In this study, the assay kits specific for SOD (CAT No. 706002, Cayman Chemical Company, Ann Arbor, MI, USA), catalase (CAT No. A22180, Invitrogen, Carlsbad, CA, USA), and GSH (CAT No. ADI-900-160, Enzo Life Sciences, Plymouth Meeting, PA, USA) were used to determine the respective activities. All the experiments were carried out according to the manufacturer's instructions.

PROPIDIUM IODIDE (PI) STAINING AND CELL CYCLE ANALYSIS

PLFs were exposed to various GO concentrations. At various times, the cells (2×10^6 cells) were fixed with 70% ethanol at 4 °C for 24 h and then incubated at room temperature for 1 h in a staining mixture (500 μ l) containing 125 μ l of phosphate-buffered saline (PBS), 125 μ l of 1 mg/ml RNase, and 250 μ l of 50 μ g/ml PI. Ten thousand cells per experiment were counted to measure the PI intensity using a FACS Vantage[®] system (Becton Dickinson), and cell cycle progression was analyzed using the WinMDI 2.9 program.

WESTERN BLOT ANALYSIS

Cell lysates were prepared in NP-40 lysis buffer (30 mM Tris-Cl, pH 7.5, 1 mM EDTA, 150 mM NaCl, 1% NP-40, 1 mM PMSF, and protease inhibitor mixture containing 1 μ g/ml each of aprotinin and leupeptin). Protein contents were quantified using the Bradford method (1976). Samples of extracts containing equal amounts of protein (30 μ g/sample) were analyzed by SDS-PAGE (10–15% gels) and blotted onto polyvinyl difluoride membranes. The blots were probed with primary antibodies and incubated with horseradish peroxidase-conjugated anti-IgG in blocking buffer for 1 h. After washing, the blots were developed with enhanced chemiluminescence (Santa Cruz Biotechnology) and exposed to X-ray film (Eastman-Kodak, Rochester, NY, USA).

ALKALINE PHOSPHATASE (ALP) ACTIVITY ASSAY

PLFs were incubated in osteoblast-inducing medium with and without various GO concentrations (0–10 mU/ml), 500 U/ml SOD, or 500 U/ml catalase. At various times, the cells were washed several times with PBS (pH 7.4), sonicated for 20 s at 4 °C, and then centrifuged at 12,000g for 15 min. Protein levels were determined by the Bradford method (1976), and the normalized protein lysates were adjusted for the determination of ALP activity. ALP activity was determined at 37 °C in a buffer (10 mM MgCl₂ and 0.1 M alkaline buffer, pH 10.3) supplemented with 10 mM *p*-nitrophenylphosphate as the substrate. The reaction was stopped by adding 0.5 N NaOH, and absorbance was measured at 405 nm. ALP activity was expressed as nmol/min/mg of protein.

ALIZARIN RED STAINING

The degree of mineralization in PLFs was determined in 6-well plates using Alizarin red staining. In brief, PLFs were exposed to GO in osteoblast-inducing medium with and without antioxidant enzymes. At various times, they were fixed with ice-cold 70% (vol/vol) ethanol for 1 h. The cells were stained with 0.2% Alizarin red S in distilled water for 30 min at room temperature. After destaining and air-drying, the cell culture plates were evaluated under light microscopy. In order to quantify the red dye, the stains were solubilized with 10% acetylpyridinium chloride by shaking for 20 min, and the absorbance was measured using a microplate reader.

MEASUREMENT OF COLLAGEN AND OSTEOCALCIN

Collagen contents in PLF were determined by the Sirius Red-based colorimetric assay. In brief, PLFs were treated with various GO concentrations (0–10 mU/ml) and/or antioxidant enzymes in DMEM containing dexamethasone, ascorbic acid, and glycerophosphate. The medium was changed every 2 days. Ten days after treatment, the cells were fixed with Bouin's fluid for 1 h and washed several times with distilled water. The culture plates were stored at room temperature for drying before being stained with Sirius Red dye reagent for 1 h. After two washes with 10 mM HCl, the cells were treated with 100 mM NaOH, and the absorbance at 550 nm was measured. In addition, the osteocalcin contents in PLF were measured at 10 days after GO treatment using a sandwich ELISA assay kit (Biomedical Technologies Inc., USA). All the experiments were performed according to the manufacturer's instructions. The contents of collagen and osteocalcin were expressed as μg or ng per 10^6 cells.

RNA ISOLATION AND REAL TIME RT-PCR

The total RNA was extracted from PLFs using STAT-60 (Tel-Test, Inc., Friendwood, Tex., USA) and the real-time quantification of RNA targets was performed in the Rotor-Gene 2000 real-time thermal cycling system (Corbett Research, NSW, Australia) using a QuantiTect SYBR Green RT-PCR kit (QIAGEN, CA, USA), as described elsewhere (Heo and Lee, 2011). The primers used were 5'-GAGGGACTATGGCGTCAAACA-3' (sense), 5'-GGATCCCAAAAGAAGCTTTGC-3' (antisense) for Runx2, and 5'-TCAGCCGCCCGATCTTCCA-3' (sense), 5'-AATGGGTCCACCGCGCCAAG-3' (antisense) for osterix. The temperature of the PCR products was increased from 65 to 99°C at a rate of 1°C/5 sec, and the resulting data was analyzed using the software provided by the manufacturer.

STATISTICAL ANALYSIS

Unless specified otherwise, all the data are expressed as the mean $\pm$ standard deviation (SD) from three or more independent experiments. A one-way analysis of variance (ANOVA) followed by a Scheffe's test was used for multiple comparisons using the SPSS program (version 18.0). A value of $p < 0.05$ was considered statistically significant.

RESULTS

CONTINUOUS GENERATION OF H_2O_2 IN GO-EXPOSED PLFS

The concentrations of H_2O_2 in the culture supernatants were augmented proportionally to the amounts of GO added (Fig. 1A). The H_2O_2 levels produced were increased continuously after the addition of GO, peaked at around 4 h, and gradually decreased after 6 h. This dynamic pattern was more apparent in the culture supernatants exposed to higher doses of GO, such that 20 mU/ml GO treatment produced 28.9 μM at 4 h as a maximum concentration and this was reduced up to 13.5 μM at 24 h. These results were quite similar to the previous reports showing that GO treatment generates H_2O_2 continuously at low concentrations with a complex dynamic [Lee et al., 2006; Son et al., 2009; Kaczara et al., 2010]. The patterns of GO-mediated production were not affected by DAG at a significant level (Fig. 1B). In parallel with the previous report (Kaczara et al., 2010), the amounts of H_2O_2 generated by GO were also not influenced by the presence of 10% FBS (data not shown).

GO TREATMENT AFFECTS THE VIABILITY AND PROLIFERATION OF PLF BASED ON CONCENTRATION

The PLFs showed multiple responses to GO depending on the concentration. The viability of PLF was increased when exposed to 10 mU/ml GO for 24 h, whereas the same viability was markedly reduced after exposure to 25 mU/ml GO for the same time period (Fig. 2A). GO treatment (10 mU/ml) also resulted in the increase in TdR uptake in PLFs (Fig. 2B). This was accompanied by a significant increase in cells in the G₂/M phase of cell cycle progression (Fig. 2C). In contrast, the direct addition of H₂O₂ suppressed the viability of PLFs in a dose-dependent manner (Fig. 2D). The viability of the cells was reduced to approximately 30% of that of the untreated control cells when they were exposed to 1 mM H₂O₂. Pretreatment of PLFs with 500 U/ml catalase almost prevented a complete GO-mediated reduction in viability (Fig. 2E).

GO EXPOSURE AT LOW CONCENTRATIONS ENHANCES SOD ACTIVITY IN PLFs

SOD activity in PLFs was affected based on the concentration of GO added. Treatment with GO ranging from 1 to 10 mU/ml augmented SOD activity in the cells, but the addition of 20 mU/ml GO significantly reduced the activity (Fig. 3A). GO exposure did not increase catalase activity in the cells but reduced it in a dose-dependent manner (Fig. 3B). However, the reduction in GSH in PLFs was not changed with the addition of GO at the concentrations examined.

GO TREATMENT INCREASES THE PROLIFERATION RATE IN PLFS INCUBATED IN OSTEOGENIC MEDIUM

We subsequently explored whether GO treatment affects the proliferation rate of PLFs incubated in osteogenic medium supplemented with DAG. Incubation of PLFs for 7 days in the medium resulted in a decrease in DNA synthesis in the cells compared to that in untreated control cells (Fig. 4A). This decrease was not inhibited by the addition of 1 mU/ml GO, but the combination with 10 mU/ml GO for the same amount of time blocked the DAG-mediated decrease in TdR uptake level in the cells. Flow cytometric analysis after PI staining revealed a significant reduction of the cell population in the S phase in DAG-treated cells and its suppression by the combined treatment with 10 mU/ml GO (Figs. 4B and C). Treatment of PLFs with more than 20 mU/ml GO caused cytotoxic effects with the attendant migration into the sub-G₁ phase of cell cycle progression (data not shown). The levels of cyclin D1 and CDK2 proteins in PLFs were not affected by incubation in osteoblast-inducing medium regardless of the presence of GO ranging from 1 to 10 mU/ml (Fig. 5A). In contrast, a significant reduction in cyclin E protein level was found in cells incubated in DAG-containing medium for 7 days (Figs. 5A and B). In addition, combined treatment with GO prevented the DAG-mediated decrease in cyclin E level in a dose-dependent manner. When the DAG-treated PLFs were co-incubated with 10 mU/ml GO for 7 days, an approximately 5-fold increase in cyclin E level was observed compared to that of the untreated control cells (Fig. 5B).

GO TREATMENT AT LOW DOSES STIMULATES OSTEOBLASTIC DIFFERENTIATION OF PLF

Since it is known that the differentiation of cells into osteoblasts is accompanied by the expression and activity of ALP, we determined the effects of GO on ALP activity in PLFs during osteoblastic differentiation. Three days after incubation, DAG treatment itself increased ALP activity in the cells, whereas the increase was significantly attenuated by co-incubation with GO (Fig. 6A). Such GO-mediated suppression in ALP activity was observed when the DAG-treated cells were cultured for 7 days in the presence of 1 or 2.5 mU/ml GO (Fig. 6B). However, a combined treatment with 10 mU/ml GO for the same time period

augmented the ALP activity that had increased in DAG-treated PLFs. This augmentation was further apparent when the cells were incubated in combination with 10 mU/ml GO for 10 days (Fig. 6C). Figure 6D shows a time-dependent increase and acceleration in ALP activity in DAG-treated PLFs after combined treatment with 10 mU/ml GO during osteoblastic induction. We next evaluated the effects of GO on the mineralization of DAG-treated PLFs using Alizarin red staining. Figure 7A represents the transparent appearance of Alizarin red stained-PLFs and the increase in the number of stained cells due to GO in a dose-dependent manner when the cells were analyzed after 14 days of incubation. Results from colorimetric analysis showed that combined treatment with 5 and 10 mU/ml GO led to a significant increase in the optical density up to 175% and 195%, respectively, compared to that of the control (Fig. 7B). In parallel with these findings, GO-mediated augmentation of collagen content was observed in the cells co-incubated with DAG for 10 days (Fig. 8A). The combination with 10 mU/ml GO also augmented the osteocalcin levels that had increased in the DAG-treated PLFs (Fig. 8B).

GO-MEDIATED FACILITATION OF OSTEOGENIC DIFFERENTIATION IS INHIBITED BY CATALASE

In order to determine whether GO-mediated stimulation of osteoblastic differentiation in PLFs is related to the effects of H₂O₂, the DAG-treated PLFs were incubated in the presence and absence of SOD and catalase. GO-mediated acceleration of ALP activity in DAG-treated PLFs was significantly prevented with the addition of 500 U/ml catalase (Fig. 9A). Catalase also inhibited GO-mediated augmentations of collagen content (Fig. 9B), mineralization (Fig. 9C), and osteocalcin level (Fig. 9D). However, the addition of 500 U/ml SOD failed to suppress GO-mediated facilitation on the induction of osteogenic markers in the cells. None of these antioxidants alone significantly influenced cell viability or DNA synthesis (data not shown).

GO-MEDIATED FACILITATION OF OSTEOBLASTIC DIFFERENTIATION IN PLFS IS ACCOMPANIED BY THE INCREASED INDUCTIONS OF OSTERIX AND RUNX2

Transcription factors such as osterix and Runx2 play a key role during osteoblastic differentiation by affecting a diverse array of down-stream effectors required for bone formation. We investigated the effects of GO in the induction of these factors and osteopontin during osteoblastic differentiation using Western blot and real time RT-PCR analyses. The protein levels of osterix and Runx2 in DAG-treated PLFs barely increased after 3 days of addition of 10 mU/ml GO (Figs. 10A and B). This increase became prominent when the cells were incubated for 7 days in the presence of GO (Figs. 10C and D). Combined incubation with GO for 7 days also augmented the levels of osteopontin in DAG-treated cells. In parallel with this, mRNA levels of Runx2 and osterix were significantly increased in the cells exposed to 10 mU/ml GO, compared to the untreated controls (Figs. 10E and F). The GO-mediated increase of Runx2 and osterix at mRNA levels was significantly diminished by treating the cells with 500 U/ml catalase.

DISCUSSION

The present findings demonstrate for the first time that GO at low concentrations increases the viability of PLFs and their ability to synthesize DNA in cultures supplemented with and without DAG. We also showed that DAG-stimulated osteoblastic differentiation in cells was augmented by combined treatment with GO. In addition, the PLF DNA synthesis elicited by GO was not induced by direct addition of H₂O₂ into the cultures. These results are in part consistent with previous reports that showed that the proliferation rate in bovine aortic endothelial cells was facilitated by GO but not by H₂O₂ [Ruiz-Ginés et al., 2000]. Burdon et al. (1996) also reported that exposure to 1 μ M H₂O₂ stimulated the proliferation of BHK-21

fibroblasts, but this agent at 0.5 and 1 mM caused growth inhibition and apoptotic cell death. However, there have been many reports demonstrating the opposite action of H₂O₂ on cells. The direct addition of H₂O₂ inhibited viability as well as osteoblastic differentiation in primary bone marrow stromal cells [Bai et al., 2004; Liu et al., 2004], MC3T3-E1 preosteoblastic cells [Choi et al., 2009; Kim et al., 2010; Xu et al., 2011], and primary rabbit calvarial osteoblasts [Bai et al., 2004]. Many studies have shown that active components with antioxidant potential protect cells against the H₂O₂-induced inhibition of proliferation and osteoblastic differentiation [Choi et al., 2009; Kim et al., 2010; Liu et al., 2004]. These reports suggest that H₂O₂-mediated oxidative stress acts predominantly as an inhibitory mediator on cell proliferation and differentiation. Although we cannot explain the exact reasons involved in the opposing roles of H₂O₂ on cells, accumulating evidence supports three main explanations. One is that the cellular responses to H₂O₂ are affected according to the origins of cells examined. Second, the concentration of H₂O₂ might affect cellular responses to the agent, regardless of the method of exposure, i.e., GO versus H₂O₂ itself. Third, the cellular responses to H₂O₂ could depend on the time at which the cells are exposed to the agent and whether it is in a continuous or temporary manner. When cells are incubated with GO, H₂O₂ is generated continuously in small amounts, whereas the direct addition of H₂O₂ to the cultures leads to a high but temporary stimulus.

Studies have shown that differentiated osteoblastic cells, not proliferating cells, are sensitively affected by H₂O₂. This is believed to be due to the subsequent decrease in the intracellular antioxidant defense system according to the differentiation. For example, MC3T3-E1 cells co-incubated with ascorbic acid and β -glycerophosphate exhibited lower expressions and activities of catalase, GSH peroxidase, total SOD, and Cu/Zn SOD compared to those of the untreated control cells [Chaves Neto et al., 2011]. It was also reported that MnSOD activity regulated cellular proliferation and quiescence. Thus, its decrease facilitated a superoxide signal-mediated proliferation, but its increase induced quiescence [Sarsour et al., 2008]. Similarly, the expression of MnSOD in MC3T3-E1 cells was higher in cells grown in osteoblast-inducing medium than it was in cells grown in growth medium [Chaves Neto et al., 2011]. These findings support a correlation between the changes in intracellular antioxidant activities and the susceptibility of bone cells to H₂O₂. However, our present data showed that the activity of total SOD increased in PLFs exposed to GO ranging from 1 to 10 mU/ml, although volumes greater than 5 mU/ml GO decreased the activity of catalase. The levels of reduced GSH were not changed in PLFs even when the cells were treated with 20 mU/ml GO. Moreover, exogenous SOD did not affect the differentiation of PLFs into osteoblasts. This suggests that the cellular antioxidant defense system is not directly related to the proliferation elicited in GO-exposed PLFs. Rather, the induction of cyclin E protein in PLFs is thought to be related to GO-stimulated proliferation. This is supported by the roles of cyclin E on cell cycle progression, the concentration of which increases in the late G₁ phase, decreases in the early S phase and induces the initial processes of DNA replication.

Osteogenic transcription factors including Runx2 and osterix are essential for bone formation and osteoblast differentiation [Ichida et al., 2004; Komori, 2005]. Runx2, known as core-binding factor 1 (Cbfa1), is located in the promoter regions of all the osteoblast-specific genes and controls their expressions. The results from this study show that osteoblastic differentiation in DAG-treated PLFs is augmented by combined treatment with GO, as demonstrated by the increases in ALP activity, mineralization, and collagen and osteocalcin contents. GO-mediated increases in these osteoblastic markers are correlated with the inductions of Runx2 and osterix. We previously found that stimulating PLFs with lithium chloride significantly elicited mineralized nodule formation and ALP activation, which is accompanied by the up-regulation of Runx2 and osterix [Heo et al., 2010]. Therefore, it is suggested that the activations of the transcription factors Runx2 and osterix

are required for the osteogenic differentiation of PLFs. Combined treatment with GO at low concentrations facilitates the osteoblastogenesis of the cells by stimulating these transcription factors.

As it has been suggested that H₂O₂ sensitively down-regulates osteoblastic differentiation rather than proliferation, our current results demonstrated a contradictory action of H₂O₂ on the proliferation and osteoblastic differentiation of PLFs. One of the possible mechanisms that can explain these conflicting results is increased protein phosphorylation according to H₂O₂ exposure [Ruiz-Ginés et al., 2000; Takada et al., 2003]. It has been documented that signaling pathways including bone morphogenetic protein (BMP)-Smad signaling, mitogen-activated protein kinases (MAPKs), and phosphatidylinositol 3 kinase are involved in osteoblast differentiation [Bai et al., 2004; Ghosh-Choudhury et al., 2002]. Accumulating evidence also suggests that the transcriptional activity of Runx2 is controlled through phosphorylation by cellular kinases such as MAPKs and protein kinase A [Ge et al., 2009; Greenblatt et al., 2010]. This indicates that the osteoblastic differentiation of cells is affected by the statuses of cellular protein kinases. There is a report showing that fibroblast growth factor-2 induces the osteoblastic differentiation of vascular smooth muscle cells by activating Runx2 through MAPK-dependent and oxidative stress-sensitive signaling pathways [Nkahara et al., 2010]. It is also known that H₂O₂ induces the tyrosine phosphorylation of various growth factor receptors and oncogens, eventually activating MAPKs [Guyton et al., 1996; Rao, 1996]. More detailed experiments will be needed to elucidate the mechanisms by which GO accelerates the osteoblastic differentiation of PLFs. Investigation to verify the up-stream and down-stream effectors of Runx2 and osterix are also required in future studies.

In conclusion, PLFs are capable of differentiating into osteoblasts and regulating the balanced activation of osteoblasts and osteoclasts during orthodontic tooth movement. Although the precise mechanisms involved in the GO-mediated stimulation of proliferation and osteoblastic differentiation in PLFs are not clearly defined, our present findings suggest that H₂O₂ at relatively low concentrations elicits the functions of PLFs in maintaining the dimensions between the tooth root and alveolar bone, as well as in mediating a balanced metabolism in alveolar bone remodeling. It should also be noted that ROS are generated in a small amount around and/or in the tension or compression side of the PDL after mechanical stimuli. Collectively, this study demonstrates that GO treatment stimulates the proliferation of PLFs through the up-regulation of cyclin E and augments their potential to differentiate into osteoblasts through the induction of bone-specific transcription factors.

Acknowledgments

This research was supported by Basic Science Research Program through the National Research Foundation (NRF) funded by the Ministry of Education, Science and Technology (No. 2009-0067712). Part of this research was supported by grants from the National Institutes of Health (NIH) (R01ES015518 and R01CA116697).

References

- Ara T, Kurata K, Hirai K, Uchihashi T, Uematsu T, Imamura Y, Furusawa K, Kurihara S, Wang PL. Human gingival fibroblasts are critical in sustaining inflammation in periodontal disease. *J Periodont Res*. 2009; 44:21–27. [PubMed: 19515019]
- Bai XC, Lu D, Bai J, Zheng H, Ke ZY, Li XM, Luo SQ. Oxidative stress inhibits osteoblastic differentiation of bone cells by ERK and NF-kappaB. *Biochem Biophys Res Commun*. 2004; 314:197–207. [PubMed: 14715266]
- Bordin S, Page RC, Narayanan AS. Heterogeneity of normal human diploid fibroblasts: Isolation and characterization of one phenotype. *Science*. 1984; 223:171–173. [PubMed: 6691142]

- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976; 72:248–254. [PubMed: 942051]
- Burdon RH, Gill V, Alliangana D. Hydrogen peroxide in relation to proliferation and apoptosis in BHK-21 hamster fibroblasts. *Free Radic Res.* 1996; 24:81–93. [PubMed: 8845916]
- Buttke TM, Sandstrom PA. Oxidative stress as a mediator of apoptosis. *Immunol Today.* 1994; 15:7–10. [PubMed: 8136014]
- Chandra J, Samali A, Orrenius S. Triggering and modulation of apoptosis by oxidative stress. *Free Radic Biol Med.* 2000; 29:323–333. [PubMed: 11035261]
- Chapple IL, Milward MR, Dietrich T. The prevalence of inflammatory periodontitis is negatively associated with serum antioxidant concentrations. *J Nutr.* 2007; 137:657–664. [PubMed: 17311956]
- Chaves Neto AH, Machado D, Yano CL, Ferreira CV. Antioxidant defense and apoptotic effectors in ascorbic acid and β -glycerophosphate-induced osteoblastic differentiation. *Dev Growth Differ.* 2011; 53:88–96. [PubMed: 21261614]
- Choi EM, Kim GH, Lee YS. *Atractylodes japonica* root extract protects osteoblastic MC3T3-E1 cells against hydrogen peroxide-induced inhibition of osteoblastic differentiation. *Phytother Res.* 2009; 23:1537–1542. [PubMed: 19441015]
- Ekuni D, Tomofuji T, Sange T, Irie K, Azuma T, Maruyama T, Tamaki N, Murakami J, Kokeguchi S, Yamamoto T. Vitamin C intake attenuates the degree of experimental atherosclerosis induced by periodontitis in the rat by decreasing oxidative stress. *Arch Oral Biol.* 2009; 54:495–502. [PubMed: 19296928]
- Forman HJ, Torres M. Redox signaling in macrophages. *Mol Aspects Med.* 2001; 22:189–216. [PubMed: 11679166]
- Ge C, Xiao G, Jiang D, Yang Q, Hatch NE, Roca H, Franceschi RT. Identification and functional characterization of ERK/MAPK phosphorylation sites in the Runx2 transcription factor. *J Biol Chem.* 2009; 284:32533–32543. [PubMed: 19801668]
- Ghosh-Choudhury N, Abboud SL, Nishimura R, Celeste A, Mahimainathan L, Choudhury GG. Requirement of BMP-2-induced phosphatidylinositol 3-kinase and Akt serine/threonine kinase in osteoblast differentiation and Smad-dependent BMP-2 gene transcription. *J Biol Chem.* 2002; 277:33361–33368. [PubMed: 12084724]
- Greenblatt MB, Shim JH, Zou W, Sitara D, Schweitzer M, Hu D, Lotinun S, Sano Y, Baron R, Park JM, Arthur S, Xie M, Schneider MD, Zhai B, Gygi S, Davis R, Glimcher LH. The p38 MAPK pathway is essential for skeletogenesis and bone homeostasis in mice. *J Clin Invest.* 2010; 120:2457–2473. [PubMed: 20551513]
- Guyton KZ, Liu Y, Gorospe M, Xu Q, Holbrook NJ. Activation of mitogen-activated protein kinase by H_2O_2 : role in cell survival following oxidant injury. *J Biol Chem.* 1996; 271:4138–4142. [PubMed: 8626753]
- Heo JS, Lee JC. β -Catenin mediates cyclic strain-stimulated cardiomyogenesis in mouse embryonic stem cells through ROS-dependent and integrin-mediated PI3K/Akt pathways. *J Cell Biochem.* 2011; 112:1880–1889. [PubMed: 21433060]
- Heo JS, Lee SY, Lee JC. Wnt/ β -catenin signaling enhances osteoblastogenic differentiation from human periodontal ligament fibroblasts. *Mol Cells.* 2010; 30:449–454. [PubMed: 20848229]
- Herbert JM, Bono F, Savi P. The mitogenic effect of H_2O_2 for vascular smooth muscle cells is mediated by an increase of the affinity of basic fibroblast growth factor for its receptor. *FEBS Lett.* 1996; 395:43–47. [PubMed: 8849686]
- Ichida F, Nishimura R, Hata K, Matsubara T, Ikeda F, Hisada K, Yatani H, Cao X, Komori T, Yamaguchi A, Yoneda T. Reciprocal roles of MSX2 in regulation of osteoblast and adipocyte differentiation. *J Biol Chem.* 2004; 279:34015–34022. [PubMed: 15175325]
- Kaczara P, Sarna T, Burke JM. Dynamics of H_2O_2 availability to ARPE-19 cultures in models of oxidative stress. *Free Radic Biol Med.* 2010; 48:1064–1070. [PubMed: 20100568]
- Kim YH, Lee YS, Choi EM. Chrysoeriol isolated from *Eurya cillata* leaves protects MC3T3-E1 cells against hydrogen peroxide-induced inhibition of osteoblastic differentiation. *J Appl Toxicol.* 2010; 30:666–673. [PubMed: 20981859]
- Komori T. Regulation of skeletal development by the Runx family of transcription factors. *J Cell Biochem.* 2005; 95:445–453. [PubMed: 15786491]

- Kook SH, Hwang JM, Park JS, Kim EM, Heo JS, Jeon YM, Lee JC. Mechanical force induces type I collagen expression in human periodontal ligament fibroblasts through activation of ERK/JNK and AP-1. *J Cell Biochem.* 2009; 106:1060–1067. [PubMed: 19206162]
- Kook SH, Jang YS, Lee JC. Human periodontal ligament fibroblasts stimulate osteoclastogenesis in response to compression force through TNF- α -mediated activation of CD4+ T cells. *J Cell Biochem.* 2011; 112:2891–2901. [PubMed: 21618593]
- Lee JC, Son YO, Choi KC, Jang YS. Hydrogen peroxide induces apoptosis of BJAB cells due to formation of hydroxyl radicals via intracellular iron-mediated Fenton chemistry in glucose oxidase-mediated oxidative stress. *Mol Cells.* 2006; 22:21–29. [PubMed: 16951546]
- Liu AL, Zhang ZM, Zhu BF, Liao ZH, Liu Z. Metallothionein protects bone marrow stromal cells against hydrogen peroxide-induced inhibition of osteoblastic differentiation. *Cell Biol Int.* 2004; 28:905–911. [PubMed: 15566960]
- McCulloch CA, Bordin S. Role of fibroblast subpopulations in periodontal physiology and pathology. *J Periodontol Res.* 1991; 26:144–154. [PubMed: 1830616]
- Nakahara T, Sato H, Shimizu T, Tanaka T, Matsui H, Kawai-Kowase K, Sato M, Iso T, Arai M, Kurabayashi M. Fibroblast growth factor-2 induces osteogenic differentiation through a Runx2 activation in vascular smooth muscle cells. *Biochem Biophys Res Commun.* 2010; 394:243–248. [PubMed: 19903460]
- Nordberg J, Arner ES. Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free Radic Biol Med.* 2001; 31:1287–1312. [PubMed: 11728801]
- Rao GN. Hydrogen peroxide induces complex formation of SHC-Grb2-SOS with receptor tyrosine kinase and activates Ras and extracellular signal-regulated protein kinases group of mitogen-activated protein kinases. *Oncogene.* 1996; 13:713–719. [PubMed: 8761292]
- Roberts WE, Mozsary PG, Klingler E. Nuclear size as a cell-kinetic marker for osteoblast differentiation. *Am J Anat.* 1982; 165:373–384. [PubMed: 7158610]
- Ruiz-Ginés JA, López-Ongil S, González-Rubio M, González-Santiago L, Rodríguez-Puyol M, Rodríguez-Puyol D. Reactive oxygen species induce proliferation of bovine aortic endothelial cells. *J Cardiovasc Pharmacol.* 2000; 35:109–113. [PubMed: 10630740]
- Sarsour EH, Venkataraman S, Kalen AL, Oberley LW, Goswami PC. Manganese superoxide dismutase activity regulates transitions between quiescent and proliferative growth. *Aging Cell.* 2008; 7:405–417. [PubMed: 18331617]
- Son YO, Jang YS, Heo JS, Chung WT, Choi KC, Lee JC. Apoptosis-inducing factor plays a critical role in caspase-independent, pyknotic cell death in hydrogen peroxide-exposed cells. *Apoptosis.* 2009; 14:796–808. [PubMed: 19418225]
- Takada Y, Mukhopadhyay A, Kundu GC, Mahabeleshwar GH, Singh S, Aggarwal BB. Hydrogen peroxide activates NF-kappa B through tyrosine phosphorylation of I kappa B alpha and serine phosphorylation of p65: evidence for the involvement of I kappa B alpha kinase and Syk protein-tyrosine kinase. *J Biol Chem.* 2003; 278:24233–24241. [PubMed: 12711606]
- Timblin CR, Janssen YW, Mossman BT. Transcriptional activation of the proto-oncogene c-jun by asbestos and H₂O₂ is directly related to increased proliferation and transformation of tracheal epithelial cells. *Cancer Res.* 1995; 55:2723–2726. [PubMed: 7796393]
- Tipton DA, Braxton SD, Dabbous MK. Effects of a bleaching agent on human gingival fibroblasts. *J Periodontol.* 1995; 66:7–13. [PubMed: 7891254]
- Wattanaroonwong N, Schoenmaker T, de Vries TJ, Everts V. Oestrogen inhibits osteoclast formation induced by periodontal ligament fibroblasts. *Arch Oral Biol.* 2011; 56:212–219. [PubMed: 21035111]
- Xu ZS, Wang XY, Xiao DM, Hu LF, Lu M, Wu ZY, Bian JS. Hydrogen sulfide protects MC3T3-E1 osteoblastic cells against H₂O₂-induced oxidative damage-implications for the treatment of osteoporosis. *Free Radic Biol Med.* 2011; 50:1314–1323. [PubMed: 21354302]

Fig. 1. Continuous and dynamic production of H_2O_2 in the cultures of PLFs exposed to GO. PLFs were exposed to 5, 10, and 20 mU/ml GO (left panel) or 10 mU/ml GO with DAG (10 nM dexamethasone, 50 μM ascorbic acid, and 20 mM β -glycerophosphate) (right panel) and then the culture supernatants were collected at various times intervals after GO exposure during 24 h incubation. The results indicate the mean $\pm$ SD from triplicate experiments.

Fig. 2.

Viability and proliferation of PLFs are increased by the addition of GO but not by H₂O₂. PLFs were exposed to the indicated GO concentrations (0–50 mU/ml) for 24 h and then processed for the WST-8 assay (A), tritium uptake assay (B), and flow cytometric analysis after PI staining (C). (D) The cells were also exposed to H₂O₂ itself with increasing concentrations (0–2 mM) for 24 h and then processed for the cell viability assay. (E) PLFs were exposed to 10 mU/ml GO in the presence and absence of 500 U/ml catalase for 24 h prior to the WST-8 assay. In the experiment of C, cell cycle progression for each experiment was analyzed using the WinMDI 2.9 program. * $p < 0.05$ and *** $p < 0.001$ vs. the untreated control values. # $p < 0.05$ vs. the experimental values.

Fig. 3.

The activities of intracellular antioxidant systems depend on the concentration of GO. PLFs were exposed to various GO concentrations (0–20 mU/ml) for 24 h. The activities of SOD and catalase and the levels of reduced GSH were then determined. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ vs. the untreated control values.

Fig. 4.

Effects of GO on the proliferation and cell cycle progression of DAG-treated PLFs. PLFs were exposed to the indicated GO concentrations (0–10 mU/ml) in the presence of DAG. After 7 days of incubation, the levels of DNA synthesis and the cell populations in each stage of cell cycle progression were determined by tritium uptake assay (A) and flow cytometric analysis after PI staining (B and C), respectively. The data in B show representative results from three independent experiments. * $p < 0.05$ and ** $p < 0.01$ vs. the untreated control values. # $p < 0.05$ vs. DAG treatment alone.

Fig. 5. Effects of GO on the induction of cell cycle regulatory proteins in DAG-treated PLFs. (A) The cells were incubated in the DAG-containing osteogenic medium with and without the indicated doses (0–10 mU/ml) of GO for 7 days and then processed for Western blot analysis using total protein lysates. (B) The values represented are the mean $\pm$ SD of three independent experiments, where actin was used as the control protein. * $p < 0.05$ and *** $p < 0.001$ vs. the untreated control values. ## $p < 0.01$ and ### $p < 0.001$ vs. DAG treatment alone.

Fig. 6.

GO augmentation of the DAG-mediated increase in ALP activity in PLFs.

Cells were exposed to the indicated GO concentrations (0–10 mU/ml) in the presence of DAG and processed for the analysis of ALP activity after 3 (A), 7 (B), and 10 days (C) of incubation. (D) PLFs were also incubated in DAG-containing medium with and without 10 mU/ml GO. At various times, ALP activity was determined. * $p < 0.05$ and ** $p < 0.01$ vs. the untreated control values.

Fig. 7.

Stimulating effect of GO on mineralization in DAG-treated PLFs.

(A) Cells were cultured with the DAG-containing osteogenic medium in the presence and absence of GO for 14 days. The resulting mineralization was assessed by Alizarin red staining. Each microscopic image shown is a representative of five separate experiments.

(B) Absorbance specific for Alizarin red was measured, and ** $p < 0.01$ and *** $p < 0.001$ represent significant differences between the cells treated with DAG only and those treated in combination with GO.

Fig. 8.

GO treatment increases the contents of collagen and osteocalcin in DAG-treated PLFs. Cells were incubated in the osteogenic medium with and without the indicated GO concentrations (0–10 mU/ml). The cellular levels of collagen (A) and osteocalcin (B) were determined after 10 days of incubation. * $p < 0.05$ vs. the untreated control values.

Fig. 9.

Inhibitory effect of catalase on the GO-mediated facilitation of osteogenic differentiation in PLFs.

Cells were incubated in DAG-containing osteogenic medium in the presence of 10 mU/ml GO, 500 U/ml SOD, and/or 500 U/ml catalase for 10 days and then processed for the ALP activity (A), collagen synthesis (B), mineralization (C), and osteocalcin content. * $p < 0.05$ and ** $p < 0.01$ vs. DAG treatment alone. # $p < 0.05$ and ### $p < 0.001$ vs. GO treatment.

Fig. 10.

Combined treatment with GO accelerates the expressions of osteogenic transcription factors in DAG-treated PLFs.

Cells were incubated in DAG-containing medium with and without the increasing GO concentrations (0–10 mU/ml). The levels of Osterix, Runx2, and osteopontin proteins were determined by Western blot analysis using total protein lysates after 3 (A) and 7 days (C) of incubation. The expression patterns of these proteins were analyzed using a densitometer from three independent experiments after normalizing the bands to the level of actin. The values represented in B and D are the mean $\pm$ SD corresponding to A and C, respectively. The mRNA levels of Runx2 (E) and Osterix (F) in the cells exposed to 10 mU/ml GO and/or 500 U/ml catalase were also analyzed after 7 days of osteoblastic differentiation. * p < 0.05, ** p < 0.01, and *** p < 0.001 vs. DAG treatment only. # p < 0.05 vs. GO + DAG treatment.