Published in final edited form as:

Atherosclerosis. 2010 May ; 210(1): 177–182. doi:10.1016/j.atherosclerosis.2009.11.029.

A synonymous variant in scavenger receptor, class b, type i gene is associated with lower sr-bi protein expression and function

Jason Constantineau*, Erin Greason*, Michael West*, Megan Filbin**, Jeffrey S. Kieft**, Martha Z. Carletti⁺, Lane K. Christenson⁺, and Annabelle Rodriguez*

*Department of Medicine, Division of Endocrinology and Metabolism, The Johns Hopkins University School of Medicine, Baltimore MD

⁺Department of Molecular & Integrative Physiology, University of Kansas Medical Center, Kansas City KS

**Department of Biochemistry and Molecular Genetics, University of Colorado Denver School of Medicine, Aurora, CO

Abstract

Objective—A synonymous variant within scavenger receptor class B type I gene (*SCARB1*), exon 8 rs5888, has been associated with altered lipid levels and cardiovascular risk in humans. The objective was to determine if rs5888 decreased SR-BI protein expression and function *in vitro*.

Methods—SR-BI RNA secondary structure, turnover, polysomal distribution and protein expression were examined in COS cells transfected with wild-type or rs5888-SR-BI plasmids by selective 2'-hydroxyl acylation and primer extension assays, actinomycin D inhibition, polysomal profiling, and western blotting. SR-BI function in murine macrophages stably expressing wild-type or rs5888-SR-BI was assessed by measuring the specific cell association of ¹²⁵I,³H-cholesteryl ester (CE) radiolabeled HDL.

Results—Rs5888 changed RNA secondary structure and led to marked differences in the polysomal profiles compared with wild-type transcript ($p < 0.02$). As compared to wild-type cells, COS cells expressing rs5888 had significantly lower SR-BI protein expression ($p < 0.04$), but no difference in total RNA transcript levels. There were no differences in SR-BI RNA turnover in murine macrophages, whereas specific cell association of ¹²⁵I ($p < 0.0001$) or ³H-CE ($p < 0.00001$) was significantly lower in rs5888 cells.

Conclusions—The rs5888 variant affected SR-BI RNA secondary structure, protein translation, and was significantly associated with reduced SR-BI protein expression and function *in vitro*.

Keywords

SCARB1; SR-BI; HDL; atherosclerosis; synonymous; SNPs

© 2009 Elsevier Ireland Ltd. All rights reserved.

Corresponding author: Dr. Annabelle Rodriguez, Division of Endocrinology and Metabolism, Johns Hopkins University School of Medicine, Johns Hopkins Bayview Medical Center, Mason F. Lord Building, Center Tower, Room 639, 5200 Eastern Avenue, Baltimore, MD 21224, Telephone: 410.550.8487, Facsimile: 410.550.3219, arodrig5@jhmi.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

While much is known regarding the role of scavenger receptor class B type I (SR-BI) in cardiovascular and reproductive physiology in mice (1), much less is known about its role in humans. This receptor was first isolated and characterized as a physiologically relevant lipoprotein receptor in murine tissues by Acton et al. (2). The receptor is highly expressed in the liver, and in steroidogenic tissues such as the adrenal glands and gonads (3). SR-BI participates in the selective uptake of cholesteryl esters (CE) from HDL, low-density lipoprotein (LDL), and very low density lipoprotein (VLDL); it is regulated by a number of factors including corticotrophin, estrogens, cyclic AMP, gonadotropins, testosterone and peroxisomal proliferator activated receptors (4–8).

Acton et al. (9) were the first to identify single nucleotide polymorphisms (SNPs) of the *SCARB1* gene in a white European population and associated some of these common variants with plasma lipid levels and body mass index (BMI). The nonsynonymous exon 1 SNP (rs4238001 [G2S], which encodes for an amino acid change) was significantly associated with higher HDL cholesterol (HDL-C) and lower LDL cholesterol (LDL-C) levels. The synonymous or silent exon 8 SNP (rs5888 [A350], which does not encode for an amino acid change) was associated with lower LDL-C levels, and the intron 5 SNP (C>T) showed an association with BMI. These authors were the first to suggest that *SCARB1* SNPs might affect lipid metabolism in humans.

We recently reported a significant association between the rs5888 SNP and lipid levels in a cohort of the Amish Family Diabetes Study (10). We found that HDL-C levels were significantly higher in women carriers of the exon 8 SNP, suggesting an association of this synonymous SNP with lower SR-BI protein levels. Our findings were consistent with Richard et al. (11), wherein these investigators had also previously reported an interaction with the exon 8 SNP and increased HDL-C levels.

Until recently, synonymous or silent SNPs were thought to be only markers for unidentified causal SNPs; however, accumulating evidence now suggests that synonymous SNPs might directly affect protein expression and/or function of their gene products (12–14). For example, Kimchi-Sarfaty et al. (15) showed that a synonymous SNP of the Multidrug Resistance 1 (*MDR1*) gene affected the conformation of its gene product, P-glycoprotein, and consequently the functionality of this protein.

Given the results of our previous work showing a significant association between the rs5888 SNP and HDL-C levels in humans (10), the goal of the present study was to determine if the synonymous *SCARB1* rs5888 SNP might directly affect SR-BI protein expression and function. Using several *in vitro* models, we indeed found that the rs5888 SNP affected SR-BI RNA secondary structure, which changed its ability to undergo productive protein translation leading ultimately to significantly lower SR-BI protein expression. Moreover, SR-BI function was significantly lower in macrophages expressing the rs5888 variant.

METHODS

Prediction of RNA Secondary Structure

Secondary structures of both wild-type and rs5888 SR-BI full-length cDNA were predicted using the *Mfold* program (version 3.2). Transcript sequences were folded in an automated manner by the *Mfold* program (<http://www.bioinfo.rpi.edu/applications/mfold/rna/form1.cgi>) to generate secondary structures.

Generation of human SR-BI expression Vectors

See online supplement.

Transient transfection of COS-7 Cells

See online supplement.

Stably expressing wild-type and rs5888 murine RAW macrophage cell lines

See online supplement.

Cell association uptake assays

See online supplement.

mRNA turnover assays

See online supplement.

Western blotting

See online supplement.

Polysomal profiling experiments

COS cells (2×10^7 cells) were transfected with either 200 ng/ml of DNA containing pSG5-wild-type or rs5888 using FuGene6 according to the manufacturer's protocol. Polysomal RNA was isolated as previously described (19). Briefly, immediately prior to sample collection the cells were exposed to 100 μ g/ml cycloheximide for 15 minutes at 37°C. Cells were washed in 4°C Dulbecco's PBS containing cycloheximide. Inclusion of cycloheximide fixed ribosomes along a strand of RNA, thus eliminating ribosomal run-off and stabilizing the polysomal RNA. Cells were lysed in low salt buffer (LSB, 20 mM Tris (pH 7.4), 10 mM NaCl and 3 mM MgCl₂) containing 1.2% Triton N-101. Cells were pelleted and resuspended in ~ 400 μ l of polysomal lysis buffer supplemented with Rnasin (Promega) and a cocktail of proteinase inhibitors. The cells were homogenized in an ice-cold glass dounce homogenizer and 3/4 of the supernatant was transferred into a tube containing 100 μ l of LSB containing heparin (10 μ g/ μ l). The remaining 1/4 of lysate sample representing total RNA was placed in Tri Reagent (Sigma). The supernatant was layered onto a continuous sucrose gradient (15–40%) and centrifuged at $36,000 \times g$ for 4 h. Continuous gradients were made using the Thermo Gradient Maker. Following centrifugation, each sucrose/RNA gradient was pushed through the Thermo UA6 optical detector (254nm) with integrated chart recorder and fraction collector in order to identify the fractions corresponding to the non-polysomal, 80S ribosomal, and polysomal RNA as previously described (19). RNA from individual fractions were then isolated and analyzed by quantitative RT-PCR. Each experiment was performed three times.

Selective 2'-hydroxyl acylation and primer extension (SHAPE) assays (20,21)

DNA encoding exons 7–9 of wild-type and rs5888 SR-BI RNA were generated by PCR from plasmid templates and this product was used in *in vitro* transcriptions, which were purified by denaturing gel electrophoresis, passively eluted into RNase-free water, and concentrated using Amicon concentrators. Four pmol of pure RNA was renatured by heating to 85°C for 45 seconds and cooled on the bench top, and then 3 μ L of folding buffer [333 mM Hepes (pH8.0), 333 mM NaCl, 20 mM MgCl₂] was added to the RNA (final volume of 30 μ L) and incubated at 37°C for 10 minutes. To probe the RNA, either 1 μ L of 130 mM *N*-methylisatoic anhydride (NMIA) in anhydrous dimethylsulfoxide (DMSO), or DMSO alone was added to the RNA and reactions were incubated 37°C for 60 minutes, then passed

through a Micro Bio-Spin® P-30 column. Modified positions were mapped by annealing ~2 pmol of modified RNA to a 5'-[³²P]-labeled cDNA primer and conducting reverse transcription (RT) using SuperScript™ Reverse Transcriptase III (Invitrogen). RT reactions were resolved on 10% denaturing polyacrylamide sequencing gels (alongside dideoxy-sequencing lanes), dried, visualized using a phosphorimager, and analyzed using the programs ImageQuant, Excel, and KaliedaGraph. Each experiment was performed twice.

Native Gel Electrophoresis

Two µg of wild type and rs5888 RNAs (exons 7–9) were incubated with either 2.5 mM MgCl₂ or 2.5 mM EDTA in 10% glycerol + 1X TH Buffer (66 mM Tris, 34 mM HEPES, pH 7.4) at 37°C for 5 min (20 µL final volume). Samples were run on either 8% polyacrylamide gels (1mm thick) in 1XTH Buffer + either 2.5 mM MgCl₂ or 2.5 mM EDTA at 5W each, 4°C for 16 hours. Gels were removed from the apparatus and visualized with ethidium bromide.

Statistical analysis

Student's t-test was used for the analyses. Two-way ANOVA was used to determine statistical differences in the polysomal experiments. Probability values less than 0.05 were considered statistically significant.

RESULTS

Given that rs5888 is a common (minor allele frequency ~40%) synonymous or silent SNP residing within exon 8 of the *SCARB1* gene, our first step was to examine whether the minor allele might predict changes in the SR-BI mRNA secondary structure. As shown in online supplement Figure I, using *Mfold* software (22), substitution of the major C allele with the minor T allele predicted changes in the secondary structure of the SR-BI mRNA.

Thermodynamic-based models predicted that the rs5888 variant would lead to an alteration of the mRNA secondary structure in the vicinity of the SNP. To directly test this prediction, we next used selective 2' acetylation monitored by primer extension (SHAPE) to probe the secondary structure of *in vitro* transcribed and purified RNA comprising exons 7–9 of both wild-type and rs5888-SR-BI. In the SHAPE secondary structure probing assay, modification of the 2' hydroxyl by NMIA depends on whether the nucleotide is base paired (2'-OH protected) or single-stranded (2'-OH modified), and hence changes in the modification pattern indicate changes in secondary structure. When the modification patterns of wild-type and rs5888 exon 7–9 RNAs are compared (Figure 1A and B), specific differences are observed, indicating a change in secondary structure. The most clearly observed differences are in the vicinity of the variant and these are clearly visible on the gel (Figure 1A), but other regions of subtle difference are found in other parts of exons 7–9 (Figure 1B). To further explore our observation that rs5888 possessed an altered structure, we subjected the exon 7–9 RNA to native gel electrophoresis. Native gel electrophoresis is very sensitive to changes in the overall global structure of RNA; these changes are reflected in a change in the migration rate of the altered RNA relative to wild-type RNA. When *in vitro* transcribed and purified wild-type and rs5888 RNA were electrophoresed through a gel containing 2.5 mM EDTA and a gel containing 2.5 mM MgCl₂, differences in the migration patterns were observed (Figure 1C). In the EDTA gel, the RNA is not folded into its full native structure, but residual secondary structure can form. In EDTA, the wild-type RNA runs as a single band, whereas the rs5888 RNA has a second, lower mobility band visible. This pattern likely indicates that alternate secondary structures can form in the variant RNA that are not formed in the wild-type RNA. In the Mg²⁺-containing gel, RNA forms its native structure. In Mg²⁺ wild-type RNA runs as two bands, with the upper band being more intense. However, the

rs5888 RNA runs as a smeared band, suggesting a change in the conformation of the RNA that may be interpreted as a structurally heterogeneous population with exchange between different conformational states. Note that these differences are not due to differences in the purities of the two RNA samples, as both run as a single tight band on a denaturing gel (Figure 1C, uppermost urea band).

To test whether the rs5888 SNP was associated with problems with mRNA translation, we performed polysomal profiling experiments to probe for alterations in SR-BI protein translation. Polysomal RNA was isolated from transiently transfected COS cells expressing the full-length wild-type or rs5888-SR-BI transcripts. As shown in Figure 2A, total SR-BI RNA levels were not significantly different between wild-type and rs5888 expressing cells. However, there was a significant difference between the overall distribution of SR-BI RNA transcripts in wild-type expressing cells compared with rs5888 cells (Figure 2B, $p < 0.02$, two-way ANOVA). Specifically, in the wild-type expressing cells, the majority of the SR-BI transcripts migrated with the monosomal and disomic fractions (fractions 5–6), with the remainder associated with almost equal levels of the heavier polysomal fractions (fractions 7–12). Conversely, the results for the rs5888 expressing cells showed a shift in SR-BI RNA transcripts to the lighter fractions (i.e., non-translating fractions; fractions 1–3 contain the ribonucleoprotein associated, 40S, and 60S associated mRNAs). Additionally, the rs5888 SR-BI transcripts were associated with polysomal fractions (fractions 7–9). The mRNA transcripts in these fractions contain 4 to 6 polysomes, respectively. Moreover, as shown in Figure 2C, SR-BI protein levels were significantly lower in rs5888 expressing cells compared with wild-type cells (67% lower, $p < 0.04$).

Given that total RNA transcripts were similar between wild-type and rs5888-SR-BI expressing cells but not the distribution of the two transcripts in the polysomal fractions, we next determined whether these effects might be due to alterations in SR-BI RNA turnover. We examined SR-BI RNA turnover in murine macrophages stably expressing wild-type and rs5888-SR-BI in the presence of actinomycin D (20 $\mu\text{g/ml}$) for varying periods of time (0–24 h). As shown in Figure 3, baseline total SR-BI RNA levels were similar in wild-type and rs5888 expressing macrophages. More importantly, in the presence of actinomycin D, 18S and SR-BI RNA turnover were similar between the wild-type and rs5888-SR-BI macrophages, suggesting that differences in SR-BI protein expression were not due to an effect of the variant on RNA transcription.

We then examined whether expression of rs5888 would affect SR-BI function. Nontransfected, wild-type and rs5888-SR-BI expressing macrophages were exposed to ^{125}I , ^3H -CE labeled HDL (5 μg protein/ml) in the presence and absence of excess, unlabeled HDL (250 μg protein/ml) for 16 h at 37°C. As shown in Figure 4, specific cell association of the radiolabeled apoprotein moiety (^{125}I) ($p < 0.0001$) and of the lipid labeled moiety (^3H -CE) ($p < 0.00001$) were significantly lower in rs5888 expressing cells as compared with wild-type cells. Selective CE uptake (ng/mg cell protein) was also calculated as per Kraemer et al. (23) and the results showed negative selective CE uptake for nontransfected cells (or more holoparticle uptake), 1.2 ng/mg cell protein for rs5888 cells, and 214 ng/mg cell protein for wildtype SR-BI expressing cells.

DISCUSSION

That a synonymous SNP altered the prediction of the mRNA secondary structure was consistent with observations reported by Nackley et al. (24) and others (25). Nackley et al. (24) found that synonymous haplotypes of the catechol-O-methyltransferase gene (*COMT*) also predicted changes in the mRNA secondary structure, based on predictions generated by the *Mfold* software. These synonymous SNPs predicted structures with lower free energies

and with configurations that were more stem-like, similar to what we have observed for the synonymous rs5888 SNP.

The mechanism by which synonymous SNPs change the secondary structure is unclear. It is possible that changes in the secondary structure interfered with initiation of protein synthesis at the AUG start site. It is possible that the altered secondary structure interferes with loading of the ribosomal 40S and 60S subunits on to the mRNA, as this is suggested by the results of the polysomal profiling, wherein SR-BI RNA levels are elevated in the untranslated fractions isolated from cells transfected with the rs5888 plasmid.

The lack of change in mRNA expression showed that the reduced SR-BI protein expression by the rs5888 SNP was due to a post-transcriptional effect. The results of the polysomal profiling showed an enrichment of the rs5888 RNA in untranslated fractions. To the best of our knowledge, there have not been other publications that specifically examined the effect of synonymous SNPs on protein translation using polysomal profiling. However, Signori et al. (26) did utilize polysomal profiling to examine the effects of a somatic mutation in the 5'-UTR of *BRCA1* gene on translation efficiency. These investigators transiently transfected HEK293 cells with plasmids containing either the wild-type or mutant plasmids and then performed polysomal profiling. They found that cells transfected with the mutant allele had mRNA that was more associated with the lighter, monosomic fractions, suggesting a defect in protein translation. We also do not believe that the post-transcriptional effects of the rs5888 variant on SR-BI protein expression is Dicer miRNA/siRNA dependent, as Dicer has been shown to be undetectable in monocytes and virtually undetectable or absolutely undetectable in terminally differentiated macrophages (27). We also confirmed this observation by Klase et al. (27) in that we were unable to detect Dicer protein expression in either wildtype or rs5888 macrophages transfected with scrambled or Dicer-siRNA oligonucleotides (data not shown).

We are the first to show that a synonymous *SCARB1* SNP significantly reduced SR-BI function *in vitro* as evidenced by the significant reduction in specific cell association of dual-labeled HDL. Our findings are consistent with those of Acton et al. (2) and many subsequent investigators (28–30) who have shown strong positive correlations between SR-BI protein expression and activity. Nonetheless, more work is needed to fully understand the intracellular processing and cell surface functionality of rs5888-SR-BI alone, or as part of haplotypes, associated with and without cardiovascular disease. The results from the study by Duan et al. (25) suggested that the biological effect of a particular SNP might be greatly influenced by its association with other SNPs in a haplotype. Our studies concentrated solely on the effect of the rs5888 SNP alone, and not on its effect within a haplotype.

While SR-BI knockout mice, in the background of either apoE or LDL receptor deficiency, have been shown to have accelerated atherosclerosis (1), little is known regarding complete SR-BI deficiency in humans. We were the first to show that deficiency of SR-BI in primary human macrophages is inversely correlated with HDL levels (31), but we have yet to identify subjects with complete SR-BI deficiency. While it is tempting to speculate that human carriers of the rs5888 variant might be SR-BI deficient (especially since we (10) and others (9,11) have shown associations of rs5888 with higher HDL levels and lower SR-BI protein levels), our study (31) lacked sufficient power to determine a significant association between SR-BI protein levels in human carriers of the rs5888 variant and measures of atherosclerotic disease. Nonetheless, we are continuing to recruit subjects with the aim of elucidating whether this variant or others, either alone or as part of a haplotype, significantly affect SR-BI protein expression, and in turn, affect risk for cardiovascular disease.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by a NIH NHLBI grant (HL075646) to Dr. Annabelle Rodriguez.

REFERENCES

1. Trigatti B, Rayburn H, Vinals M, Braun A, Miettinen H, Penman M, Hertz M, Schrenzel M, Amigo L, Rigotti A, Krieger M. Influence of the high density lipoprotein receptor SR-BI on reproductive and cardiovascular pathophysiology. *Proc Natl Acad Sci USA*. 1999; 96:9322–9327. [PubMed: 10430941]
2. Acton S, Rigotti A, Landschulz KT, Xu S, Hobbs HH, Krieger M. Identification of scavenger receptor SR-BI as a high density lipoprotein receptor. *Science*. 1996; 271:518–520. [PubMed: 8560269]
3. Cao G, Zhao L, Stangl H, Hasegawa T, Richardson JA, Parker KL, Hobbs HH. Developmental and hormonal regulation of murine scavenger receptor, class B, type I. *Mol Endo*. 1999; 13:1460–1473.
4. Rigotti A, Edelman ER, Seifert P, Iqbal SN, DeMattos RB, Temel RE, Krieger M, Williams DL. Regulation by adrenocorticotropic hormone of the in vivo expression of scavenger receptor class B type I (SR-BI), a high density lipoprotein receptor, in steroidogenic cells of the murine adrenal gland. *J Biol Chem*. 1996; 271:33545–33549. [PubMed: 8969220]
5. Landschulz KT, Pathak RK, Rigotti A, Krieger M, Hobbs HH. Regulation of scavenger receptor, class B, type I, a high density lipoprotein receptor, in liver and steroidogenic tissues of the rat. *J Clin Invest*. 1996; 98:984–995. [PubMed: 8770871]
6. Langer C, Gansz B, Goepfert C, Engel T, Uehara Y, von Dehn G, Jansen H, Assmann G, von Eckardstein A. Testosterone up-regulates scavenger receptor BI and stimulates cholesterol efflux from macrophages. *Biochem Biophys Res Commun*. 2002; 296:1051–1057. [PubMed: 12207878]
7. Azhar S, Nomoto A, Leers-Sucheta S, Reaven E. Simultaneous induction of an HDL receptor protein (SR-BI) and the selective uptake of HDL-cholesteryl esters in a physiologically relevant steroidogenic cell model. *J Lipid Res*. 1998; 39:1616–16280. [PubMed: 9717722]
8. Chinetti G, Gbaguidi FG, Griglio S, Mallat Z, Antonucci M, Poulain P, Chapman J, Fruchart J-C, Tedgui A, Najib-Fruchart J, Staels B. CLA-1/SR-BI is expressed in atherosclerotic lesion macrophages and regulated by activators of peroxisome proliferator-activated receptors. *Circulation*. 2000; 101:2411–2417. [PubMed: 10821819]
9. Acton S, Osgood D, Donoghue M, Corella D, Pocovi M, Cenarro A, Mozas P, Keilty J, Squazzo S, Woolf EA, Ordovas JM. Association of polymorphisms at the SR-BI gene locus with plasma lipid levels and body mass index in a white population. *Arterioscler Thromb Vasc Biol*. 1999; 19:1734–1743. [PubMed: 10397692]
10. Roberts CGP, Shen H, Mitchell BD, Damcott CM, Shuldiner AR, Rodriguez A. Variants in scavenger receptor class B type I gene are associated with HDL cholesterol levels in younger women. *Hum Hered*. 2007; 64:107–113. [PubMed: 17476110]
11. Richard E, von Muhlen D, Barrett-Connor E, Alcaraz J, Davis R, McCarthy JJ. Modification of the effects of estrogen therapy on HDL cholesterol levels by polymorphisms of the HDL-C receptor, SR-BI: the Rancho Bernardo Study. *Atherosclerosis*. 2005; 180:255–262. [PubMed: 15910850]
12. Komar AA. Genetics. SNPs, silent but not invisible. *Science*. 2007; 315:466–467. [PubMed: 17185559]
13. Nielsen KB, Sorensen S, Cartegni L, Corydon TJ, Doktor TK, Schroeder LD, Reinert LS, Elpeleg O, Krainer AR, Gregersen N, Kiems J, Andresen BS. Seemingly neutral polymorphic variants may confer immunity to splicing-inactivating mutations: a synonymous SNP in exon 5 of MCAD protects from deleterious mutations in a flanking exonic splicing enhancer. *Am J Hum Genet*. 2007; 80:416–432. [PubMed: 17273963]

14. Sauna ZE, Kimchi-Sarfaty C, Ambudkar SV, Gottesman MM. Silent polymorphisms speak: how they affect pharmacogenomics and the treatment of cancer. *Cancer Res.* 2007; 67:9609–9612. [PubMed: 17942888]
15. Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar SV, Gottesman MM. A "silent" polymorphism in the MDR1 gene changes substrate specificity. *Science.* 2007; 315:525–528. [PubMed: 17185560]
16. McFarlane AS, Koj A. Short-term measurement of catabolic rates using iodine-labeled plasma proteins. *J Clin Invest.* 1970; 49:1903–1911. [PubMed: 5460289]
17. Rodriguez A, Usher D. Anti-atherogenic effects of the acyl-CoA:cholesterol acyltransferase inhibitor, avasimibe (CI-1011), in cultured primary human macrophages. *Atherosclerosis.* 2002; 161:45–54. [PubMed: 11882316]
18. Schriewer H, Jabs HU, Schultze J, Assmann G. Preparation of high-density lipoprotein labeled exclusively at the cholesteryl ester moiety. *Clin Chim Acta.* 1982; 123:139–144. [PubMed: 7116635]
19. Holets LM, Carletti MZ, Kshirsagar SK, Christenson LK, Petroff MG. Differentiation-induced posttranscriptional control of B7-H1 in Human trophoblast cells. *Placenta.* 2008 In Press.
20. Merino EJ, Wilkinson KA, Coughlan JL, Weeks KM. RNA structure analysis at single nucleotide resolution by selective 2'-hydroxyl acylation and primer extension (SHAPE). *J Am Chem Soc.* 2005; 127:4223–4231. [PubMed: 15783204]
21. Wilkinson KA, Merino EJ, Weeks KM. Selective 2'-hydroxyl acylation analyzed by primer extension (SHAPE): quantitative RNA structure analysis at single nucleotide resolution. *Nat Protoc.* 2006; 1:1610–1616. [PubMed: 17406453]
22. Zuker, M.; Mathews, DH.; Turner, DH. Algorithms and thermodynamics for RNA secondary structure prediction: a practical guide. In: Barciszewski, J.; Clark, BFC., editors. *RNA Biochemistry and Biotechnology.* New York, NY: Kluwer Academic Publishers; p. 11–43.
23. Kraemer FB, Shen W-J, Harada K, Patel S, Osuga J-I, Ishibashi S, Azhar S. Hormone-sensitive lipase is required for high-density lipoprotein cholesteryl ester-supported adrenal steroidogenesis. *Mol Endo.* 2004; 18:549–557.
24. Nackley AG, Shabalina SA, Tchivileva IE, Satterfield K, Korchynskiy O, Makarov SS, Maixner W, Diatchenko L. Human catechol-O-methyltransferase haplotypes modulate protein expression by altering mRNA secondary structure. *Science.* 2006; 314:1930–1933. [PubMed: 17185601]
25. Duan J, Wainwright MS, Comeron JM, Naruya Saitou, Sanders AR, Gelernter J, Gejman PV. Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability and synthesis of the receptor. *Hum Mol Genet.* 2003; 12:205–216. [PubMed: 12554675]
26. Signori E, Bagni C, Papa S, Primerano B, Rinaldi M, Amaldi F, Fazio VM. A somatic mutation in the 5'-UTR of BRCA1 gene in sporadic breast cancer causes down-modulation of translation efficiency. *Oncogene.* 2001; 20:4596–4600. [PubMed: 11494157]
27. Klase Z, Kale P, Winograd R, Gupta MV, Heydarian M, Berro R, McCaffrey T, Kashanchi F. HIV-I TAR element is processed by Dicer to yield a viral micro-RNA involved in chromatin remodeling of the viral LTR. *BMC Molecular Biology.* 2007; 8:63–82. [PubMed: 17663774]
28. Pagler TA, Rhode S, Neuhofer A, Laggner H, Strobl W, Hinterndorfer C, Volf I, Pavelka M, Eckhardt ERM, van der Westhuyzen DR, Xhutz GJ, Stangl H. SR-BI-mediated high density lipoprotein (HDL) endocytosis leads to HDL resecrection facilitating cholesterol efflux. *J Biol Chem.* 2006; 281:11193–11204. [PubMed: 16488891]
29. Sun B, Eckhardt ERM, Shetty S, van der Westhuyzen DR, Webb NR. Quantitative analysis of SR-BI-dependent HDL retroendocytosis in hepatocytes and fibroblasts. *J Lipid Res.* 2006; 47:1700–1713. [PubMed: 16705213]
30. Brundert M, Heeren J, Bahar-Bayansar M, Ewert A, Moore KJ, Rinninger F. Selective uptake of HDL cholesteryl esters and cholesterol efflux from mouse peritoneal macrophages independent of SR-BI. *J Lipid Res.* 2006 epub.
31. West M, Greason E, Kolmakova A, Jahangiri A, Asztalos B, Pollin TI, Rodriguez A. Scavenger receptor class B type I protein as an independent predictor of HDL cholesterol levels in subjects with hyperalphalipoproteinemia. *J Clin Endocrinol Metab.* 2009; 94:1451–1457. [PubMed: 19158204]

Figure 1. RNA Structural Changes due to SNP

A. Gel of wild-type and rs5888 RNAs probed using SHAPE, oriented 5' at the top and 3' at the bottom. Exons 7–8 are indicated by vertical grey line, and the C→T SNP indicated by an arrow. Lanes 1–4 are dideoxy sequencing ladders, lanes 5 and 7 contain control reactions for dimethyl sulfoxide (DMSO) background cleavage, and lanes 6 and 8 contain the NMIA modified wild-type and rs5888 RNAs. Dashed boxes and enlarged gel region highlights a region of interest where rs5888 is becoming more single stranded (green) or more double stranded (red) as compared to wild-type. **B.** A quantitated trace of the gel shown in panel A, and turned on its side. Black is WT, Orange is rs5888. The two are overlaid. The dashed box indicates the portion of the gel that is boxed in panel A. The two traces are very similar, showing that overall, the secondary structures between the two are quite similar. However, distinct areas of local changes are seen. These are indicated with red or green bars above the trace. There is a cluster of changes right around the site of the SNP. **C.** Native gel analysis of both wild-type and rs5888 RNAs in 2.5 mM EDTA (top panel, unfolded) or 2.5 mM MgCl₂ (bottom panel, folded).

Figure 2. Polysomal profiling of SR-BI mRNA in transfected COS cells

(A) SR-BI RNA levels were similar in cells transfected with either wild-type and rs5888-SR-BI plasmids. (B) Polysomal profiling showed that the rs5888 SR-BI does not properly assemble ribosomal units resulting in increased amounts of SR-BI mRNA associated with the RNP, 40S and 60S pools (i.e., fractions 1–5) when compared to the wild-type ($p < 0.02$, two-way ANOVA). (C) SR-BI protein levels were significantly lower in COS cells transfected with rs5888-SR-BI compared with cells transfected with wild-type SR-BI, $p < 0.04$ compared to wild-type cells. The western blot image is representative of three independent experiments, while the bar graphs are the mean $\pm$ standard error of three independent experiments.

Figure 3. SR-BI mRNA turnover is similar between macrophages stably expressing wild-type and rs5888-SR-BI

SR-BI and 18S RNA levels were similar in wild-type and rs5888-SR-BI expressing macrophages in the presence of actinomycin D (20 µg/ml) for varying periods of time (0–24 h).

Specific Cell Association of ^{125}I , ^3H CE radiolabeled HDL

Figure 4. Specific cell association of ^{125}I , ^3H -CE labeled HDL is significantly lower in murine macrophages stably expressing rs5888-SR-BI

Nontransfected, wildtype, and rs5888 stably expressing RAW macrophages were incubated with ^{125}I , ^3H -CE labeled HDL (5 μg protein/ml) in the presence and absence of excess, unlabeled HDL (250 μg protein/ml) for 16 h at 37°C. Specific cell association of the ^{125}I -labeled apoprotein moiety ($p < 0.0001$) and of the ^3H -CE labeled moiety ($p < 0.00001$) were significantly lower in cells expressing the rs5888-SR-BI compared to wild-type cells. The results are the mean $\pm$ standard error of two independent experiments.