Transcriptional Response of the Cellobiose Dehydrogenase Gene to Cello- and Xylooligosaccharides in the Basidiomycete *Phanerochaete chrysosporium*

Chiaki Hori, Hitoshi Suzuki,* Kiyohiko Igarashi, and Masahiro Samejima

Department of Biomaterial Sciences, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan

Cellobiose dehydrogenase (CDH) gene transcripts were quantified by reverse transcription-PCR (RT-PCR) in cultures of *Phanerochaete chrysosporium* supplemented with various cello- and xylooligosaccharides in order to elucidate the mechanism of enhanced CDH production in xylan/cellulose culture. Cellotriose and cellotetraose induced *cdh* expression, while xylobiose and xylotriase induced expression of cellobiohydrolase genes, especially *cel7C*.

The basidiomycete *Phanerochaete chrysosporium* produces various extracellular carbohydrate-degrading enzymes (14, 24) and oxidative enzymes (13) that degrade wood, which contains cellulose, hemicellulose, and lignin as major components. Cellulose is a β -1,4-linked homopolymer of glucose, and the cellulolytic system of this fungus involves multiple endoglucanases (EGs) and cellobiohydrolases (CBHs) (20, 21), which hydrolyze cellulose into cellobiose and cellobiohydrolases. In addition to hydrolytic enzymes, redox enzymes such as cellobiose dehydrogenase (CDH) are involved in cellulose degradation (1, 3, 4, 23). CDH oxidizes cellobiose and cellobiohydrolases into the corresponding 1,5- δ -lactones (5, 15). The role of CDH in cellulose degradation has been widely discussed (7, 8, 10, 25), and its relationship with polysaccharides monooxygenase was recently pointed out (12, 16, 22). A study by Dumonceaux and coworkers on a CDH-deficient mutant of the wood-rotting basidiomycete *Trametes versicolor* indicated that CDH is also involved in degradation of other plant cell wall components (2). We have recently reported that arabinoxylan, the major hemicellulose of monocots, enhances fungal growth and CDH production when it is supplemented into cellulose-degrading cultures, although it could not serve as a carbon source for *P. chrysosporium* by itself (6). In the present study, a similar response of the fungus was observed when glucuronoxylan (from beech wood) was added to cellulolytic culture, suggesting that the main-chain structure of xylan affects fungal growth, extracellular protein production and CDH activity. To elucidate

the mechanism of these effects, we used reverse transcription-PCR (RT-PCR) to quantify CDH gene transcripts in cultures supplemented with xylose or xylooligosaccharides as well as glucose or cellobiohydrolases.

Phanerochaete chrysosporium strain K-3 (9) was cultivated in Kremer and Wood medium (11) for 4 days with cellulose (CF11; Whatman, Fairfield, NJ), with or without xylan (from oat spelt, Serva Electrophoresis, Heidelberg, Germany; from beech, Sigma-Aldrich, St. Louis, MO). Briefly, xylan from beech wood has side chains of 4-O-methyl- β -D-glucuronic acid whereas that from oat spelt has additional side chains of α -L-arabinose. The packed volume of fungal mycelium and protein concentration of the culture filtrate were estimated as described previously (6). Cellobiose dehydrogenase (CDH) activity of the culture filtrates was assayed using cellobiose and cytochrome *c* as described previously (17). For transcriptional analysis in cultures with purified xylooligosac-

Received 18 January 2012 Accepted 28 February 2012

Published ahead of print 9 March 2012

Address correspondence to Masahiro Samejima, amsam@mail.ecc.u-tokyo.ac.jp.

* Present address: Department of Chemical Engineering & Applied Chemistry, University of Toronto, Toronto, ON, Canada.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.00150-12

FIG 1 Time courses of fungal volume (A), concentration of extracellular protein (B), and CDH activity (C) of *P. chrysosporium* grown in culture medium containing cellulose (filled square), cellulose and oat spelt xylan (open circle), and cellulose and beech xylan (filled triangle). Error bars show the standard deviations for the results from three experiments.

FIG 2 Time course of *cdh* transcripts of *P. chrysosporium* grown in culture medium supplemented with xylooligosaccharides and cellobiosaccharides. (A) X, xylose; X₂, xylobiose; X₃, xylotriose; X₄, xylotetraose. (B) G, glucose; C₂, cellobiose; C₃, cellotriose; C₄, cellotetraose; C₅, cellopentose. Control means culture without saccharides. Error bars show standard deviations for results from three experiments. Note that the scale on the vertical axis is not the same in panels A and B.

charides, *P. chrysosporium* was precultivated for 3 days and the mycelium was transferred into resting medium as previously described (18). After resting cultivation for 6 h, 100 μ M glucose, cellobiose, cellotriose, cellotetraose, cellopentose, xylose, xylobiose, xylotriose, or xylotetraose was added to the medium. Five

milliliters of culture was collected every hour until 5 or 6 h, and the mycelium was immediately frozen in liquid nitrogen for the preparation of total RNA and corresponding cDNA. The gene transcripts of *cdh*, *cel6A*, *cel7C*, and *cel7D* were quantified using quantitative PCR (qPCR) since the corresponding proteins are the

FIG 3 Time courses of *cel6A* (A), *cel7C* (B), and *cel7D* (C) transcripts of *P. chrysosporium* grown in culture medium supplemented with xylose (X), xylobiose (X₂), xylotriose (X₃), and xylotetraose (X₄). Control means culture without saccharides. Error bars show standard deviations for results from three experiments. Note that the scale on the vertical axis is not the same in panels A, B, and C.

dominant cellulases in medium containing cellulose. The primer sets were listed in the previous papers (18, 19). The actin gene was used as an internal standard.

P. chrysosporium was cultivated for 4 days in synthetic media containing cellulose only (C), cellulose and oat spelt xylan (COX), and cellulose and beech xylan (CBX) as carbon sources. As shown in Fig. 1A, the fungal volumes in COX and CBX cultures were larger than that in C culture during cultivation. As shown in Fig. 1B, production of extracellular proteins increased markedly in COX and CBX cultures but only weakly in C culture. The concentrations of extracellular protein in COX and CBX cultures were 0.37 g/liter and 0.34 g/liter, respectively, while that in culture C was 0.14 g/liter. As shown in Fig. 1C, the CDH activity in culture C was 2.2 U/liter on day 4, whereas the corresponding values in COX and CBX cultures were 42 and 41 U/liter, respectively. These results indicate that the effect of xylan is due to the β -1,4-xylan main chain structure of xyans but not the side chains.

P. chrysosporium was cultivated for 5 or 6 h in media containing xylose (X), xylooligosaccharide (DP = 2 to 4, xylobiose [X_2], xylotriose [X_3], or xylootetraose [X_4]), glucose (G), or cellobiosaccharide (degree of polymerization [DP] = 2 to 5, cellobiose [C_2], cellobiose [C_3], cellobiose [C_4], or cellobiose [C_5]), and the level of *cdh* gene transcripts was quantified by real-time RT-PCR. As shown in Fig. 2A and B, there was no change of *cdh* expression in the cultures supplemented with X and G compared with the corresponding controls. However, *cdh* was quickly upregulated (within 1 h) in the presence of X_2 , X_3 , and X_4 , reaching maximum values of 7.0×10^2 , 7.4×10^2 , and 7.4×10^2 , respectively. However, in cultures supplemented with C_3 and C_4 , the numbers of *cdh* transcripts reached 1.0×10^4 and 1.6×10^4 , respectively, at 2 h after supplementation. Thus, *cdh* gene expression is induced more strongly by cellobiosaccharides than by xylooligosaccharides. We previously found that production of glycoside hydrolase (GH) family 10 xylanase was enhanced in xylan-containing culture (6), so we speculated that xylooligosaccharides produced by the xylanase promote CDH production in the culture. However, the results of the present study indicate that upregulation of *cdh* transcription by short and unsubstituted xylooligosaccharides is not sufficient to account directly for the enhanced production of secreted protein. We next examined the effect of xylooligosaccharides on the expression of cellobiohydrolase (*cel*) genes. *cel6A*, *cel7C*, and *cel7D* transcripts were all increased by xylooligosaccharides; in particular, *cel7C* transcripts were increased up to 20-fold by xylobiose and xylotriose (Fig. 3).

In conclusion, the induction of cellulolytic enzymes by xylan in cellulose-degrading culture is not simply due to xylooligosaccharides but is more complex, involving induction by cellobiose and cellobiose. One possible interpretation is that xylooligosaccharides promote the production of cellobiohydrolases, which produce cellobiosaccharides that in turn enhance the expression of other cellulolytic genes, including *cdh*. We are currently addressing the complex gene regulation system in *P. chrysosporium* by analysis of large-scale transcriptomic data; the results will be reported shortly.

ACKNOWLEDGMENTS

This research was supported by Grants-in-Aid for Scientific Research to M.S. (no. 20380100 and 23248025) from the Japanese Ministry of Education, Culture, Sports, and Technology and a Research Fellowship to C.H. (no. 233288) from the Japan Society for the Promotion of Science (JSPS).

REFERENCES

1. Ayers AR, Ayers SB, Eriksson KE. 1978. Cellobiose oxidase, purification and partial characterization of a hemoprotein from *Sporotrichum pulverulentum*. Eur. J. Biochem. 90:171–181.
2. Dumonceaux T, Bartholomew K, Valeanu L, Charles T, Archibald F. 2001. Cellobiose dehydrogenase is essential for wood invasion and non-essential for kraft pulp delignification by *Trametes versicolor*. Enzym. Microb. Technol. 29:478–489.
3. Eriksson K. 1978. Enzyme mechanisms involved in cellulose hydrolysis by the rot fungus *Sporotrichum pulverulentum*. Biotechnol. Bioeng. 20:317–332.
4. Eriksson KE, Pettersson B, Westermark U. 1974. Oxidation: an important enzyme reaction in fungal degradation of cellulose. FEBS Lett. 49:282–285.
5. Henriksson G, Pettersson G, Johansson G, Ruiz A, Uzcategui E. 1991. Cellobiose oxidase from *Phanerochaete chrysosporium* can be cleaved by papain into two domains. Eur. J. Biochem. 196:101–106.
6. Hori C, Igarashi K, Katayama A, Samejima M. 2011. Effects of xylan and starch on secretome of the basidiomycete *Phanerochaete chrysosporium* grown on cellulose. FEMS Microbiol. Lett. 321:14–23.
7. Igarashi K, Samejima M, Eriksson K. 1998. Cellobiose dehydrogenase enhances *Phanerochaete chrysosporium* cellobiohydrolase I activity by relieving product inhibition. Eur. J. Biochem. 253:101–106.
8. Igarashi K, Tani T, Rie K, Masahiro S. 2003. Family 3 beta-glucosidase from cellulose-degrading culture of the white-rot fungus *Phanerochaete chrysosporium* is a glucan 1,3-beta-glucosidase. J. Biosci. Bioeng. 95:572–576.
9. Johnsrud SC, Eriksson KE. 1985. Cross-breeding of selected and mutated homokaryotic strains of *Phanerochaete chrysosporium* K-3—new cellulase deficient strains with increased ability to degrade lignin. Appl. Microbiol. Biotechnol. 21:320–327.
10. Kawai R, Igarashi K, Kitaoka M, Ishii T, Samejima M. 2004. Kinetics of substrate transglycosylation by glycoside hydrolase family 3 glucan (1 $\rightarrow$ 3)-beta-glucosidase from the white-rot fungus *Phanerochaete chrysosporium*. Carbohydr. Res. 339:2851–2857.
11. Kremer SM, Wood PM. 1992. Evidence that cellobiose oxidase from *Phanerochaete chrysosporium* is primarily an Fe(III) reductase. Kinetic comparison with neutrophil NADPH oxidase and yeast flavocytochrome b₂. Eur. J. Biochem. 205:133–138.
12. Langston J, et al. 2011. Oxidoreductive cellulose depolymerization by the enzymes cellobiose dehydrogenase and glycoside hydrolase 61. Appl. Environ. Microbiol. 77:7007.
13. Levasseur A, et al. 2008. FOLY: an integrated database for the classification and functional annotation of fungal oxidoreductases potentially involved in the degradation of lignin and related aromatic compounds. Fungal Genet. Biol. 45:638–645.
14. Martinez D, et al. 2004. Genome sequence of the lignocellulose degrading fungus *Phanerochaete chrysosporium* strain RP78. Nat. Biotechnol. 22:695–700.
15. Morpeth FF. 1985. Some properties of cellobiose oxidase from the white-rot fungus *Sporotrichum pulverulentum*. Biochem. J. 228:557.
16. Phillips CM, Beeson WT, Cate JH, Marletta MA. 2011. Cellobiose dehydrogenase and a copper-dependent polysaccharide monooxygenase potentiate cellulose degradation by *Neurospora crassa*. ACS Chem. Biol. 6:1399–1406.
17. Samejima M, Eriksson KE. 1992. A comparison of the catalytic properties of cellobiose:quinone oxidoreductase and cellobiose oxidase from *Phanerochaete chrysosporium*. Eur. J. Biochem. 207:103–107.
18. Suzuki H, Igarashi K, Samejima M. 2010. Cellobiose and cellobiose as inducers of the genes encoding cellobiohydrolases in the basidiomycete *Phanerochaete chrysosporium*. Appl. Environ. Microbiol. 76:6164.
19. Suzuki H, Igarashi K, Samejima M. 2008. Real-time quantitative analysis of carbon catabolite derepression of cellulolytic genes expressed in the basidiomycete *Phanerochaete chrysosporium*. Appl. Microbiol. Biotechnol. 80:99–106.
20. Uzcategui E, Johansson G, Ek B, Pettersson G. 1991. The 1,4-beta-D-glucan glucanohydrolases from *Phanerochaete chrysosporium*. Re-assessment of their significance in cellulose degradation mechanisms. J. Biotechnol. 21:143–159.
21. Uzcategui E, Ruiz A, Montesino R, Johansson G, Pettersson G. 1991. The 1,4-beta-D-glucan cellobiohydrolases from *Phanerochaete chrysosporium*.

- rium*. I. A system of synergistically acting enzymes homologous to *Trichoderma reesei*. J. Biotechnol. 19:271–285.
22. Westereng B, et al. 2011. The putative endoglucanase PcGH61D from *Phanerochaete chrysosporium* is a metal-dependent oxidative enzyme that cleaves cellulose. PLoS One 6:e27807.
 23. Westermarck U, Eriksson K. 1974. Cellobiose:quinone oxidoreductase, a new wood-degrading enzyme from white-rot fungi. Acta Chem. Scand. B 28:209–214.
 24. Wymelenberg AV, et al. 2005. The *Phanerochaete chrysosporium* secretome: database predictions and initial mass spectrometry peptide identifications in cellulose-grown medium. J. Biotechnol. 118:17–34.
 25. Yoshida M, Igarashi K, Kawai R, Aida K, Samejima M. 2004. Differential transcription of beta-glucosidase and cellobiose dehydrogenase genes in cellulose degradation by the basidiomycete *Phanerochaete chrysosporium*. FEMS Microbiol. Lett. 235:177–182.