Published in final edited form as:

Biochem Biophys Res Commun. 2012 May 4; 421(2): 361–366. doi:10.1016/j.bbrc.2012.04.018.

Live free or die: Stretch-induced apoptosis occurs when adaptive reorientation of annulus fibrosus cells is restricted

Rosalyn D Abbott¹, Alan K Howe², Helene M Langevin³, and James C Iatridis^{4,*}

¹School of Engineering, University of Vermont, Burlington, VT 05405

²Department of Pharmacology, University of Vermont, Burlington, VT 05405

³Department of Neurology, University of Vermont, Burlington, VT 05405

⁴Department of Orthopaedics, Mount Sinai School of Medicine, New York, NY 10029

Abstract

High matrix strains in the intervertebral disc occur during physiological motions and are amplified around structural defects in the annulus fibrosus (AF). It remains unknown if large matrix strains in the human AF result in localized cell death. This study investigated strain amplitudes and substrate conditions where AF cells were vulnerable to stretch-induced apoptosis. Human degenerated AF cells were subjected to 1Hz-cyclic tensile strains for 24 hours on uniformly collagen coated substrates and on substrates with 40 μ m stripes of collagen that restricted cellular reorientation. AF cells were capable of responding to stretch (stress fibers and focal adhesions aligned perpendicular to the direction of stretch), but were vulnerable to stretch-induced apoptosis when cytoskeletal reorientation was restricted, as could occur in degenerated states due to fibrosis and crosslink accumulation and at areas where high strains occur (around structural defects, delaminations, and herniations).

Keywords

Intervertebral disc; annulus fibrosus; cyclic tensile stretch; cytoskeleton; actin filaments; apoptosis; caspase-3

Introduction

The loss of cellularity in human intervertebral (IVD) degeneration is generally associated with reduced glucose and accumulation of lactic acid due to anaerobic metabolism and limited waste removal in IVDs with calcified endplates [1,2]. However, some cell death likely occurs due to structural changes in the matrix [3,4] that affect the cell microenvironment. Matrix strains are concentrated adjacent to large defects, such as clefts, fissures, and herniations that occur with degeneration [5,6], near nucleotomy sites [7], as well as near small defects such as needle puncture injuries [8]. Notably cell death is also known

© 2012 Elsevier Inc. All rights reserved

*Corresponding author: James C. Iatridis, PhD Mount Sinai School of Medicine Leni and Peter W. May Department of Orthopaedics One Gustave L. Levy Place, Box 1188 New York, NY 10029-6574 212-241-1517 (office) 212-876-3168 (fax) james.iatridis@mssm.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

to occur adjacent to injury sites, as near needle puncture injuries in the IVDs [9]. In this study, we explore the concept that some of the reduced cellularity in the IVD with degeneration is associated with stretch-induced apoptosis caused by high strain environments that occur due to structural changes and defects.

Under healthy conditions AF surface strains up to 6% were measured along the fiber direction [10] and AF fiber strains were calculated to be ~15% with a structural model [11]. AF strains calculated from MRI support the presence of large axial, radial and shear strains under physiological loading and further demonstrate strain amplification during injury and degeneration [5,7]. High magnitudes of cyclic equibiaxial strain induced apoptosis at 15% stretch in rabbit AF cells [12] and at 20% stretch in rat AF cells [13,14], however equibiaxial strain suppresses reorientation effects observed in pure uniaxial stretch of other cell types [15,16,17]. The hypothesis of this study was that human degenerated AF cells obtained from surgical samples reorient perpendicular to excessive uniaxial strains (10%, 15%, and 20% stretch), as observed in other cell types, but will undergo apoptosis when their adaptive reorientation is restricted as might occur around defects or crosslinking found with degeneration.

Materials and Methods

Isolation and culture of human AF cells from surgical samples

AF tissue was obtained from patients undergoing anterior interbody fusions for low back pain secondary to degenerative disc disease with institutional review board approval and graded by the surgeon as either Grade 4 (n=2) or Grade 5 (n=2) on the Thompson scale [18]. Cells were released from surgical tissue by sequential enzymatic digestion (0.2% protease, 0.2% collagenase, Sigma-Aldrich, St. Louis, MO). Isolated cells were filtered (70 μ m mesh, Thermo Fisher Scientific, Pittsburgh, PA), washed twice with PBS, then expanded in culture in high glucose DMEM (Invitrogen Lifesciences- Gibco, Carlsbad, CA), containing 10% fetal bovine serum (Atlanta Biologicals, Lawrenceville, GA), 1% penicillin/streptomycin (Gibco), 0.5% fungizone (Gibco), and 50 μ g/mL ascorbate (Sigma-Aldrich). All experiments were performed on passage II or III cells.

Experiments on a uniform collagen coated surface

Cells were seeded at a density of 100,000 cells/cm² in pre-coated silicone elastic membranes (4 cm² molds from STREX, www.b-bridge.com, coated with type I bovine collagen, Sigma-Aldrich C4361, 40 μ g/mL in PBS, overnight at 4°C). The cells were cultured in hypoxic conditions (5% O₂) for 24 hours to allow for attachment. Cells were assigned to either an unstretched control group (placed in the same incubator for 24hrs), or stretched 10%, 15%, or 20% on a uniaxial cell stretch device (STREX) loaded at 1Hz for 24hrs in hypoxic conditions. Control and stretched cells were imaged following loading, the supernatant was saved, and the remaining cells were either trypsinized (Gibco, no EDTA) for flow cytometry to evaluate stretch-induced apoptosis, lysed (Qiagen, Valencia, CA) for qRT-PCR, or fixed with 10% formalin (Fisher) for immunohistochemistry to evaluate reorientation effects.

For flow cytometry, the supernatant was centrifuged to obtain detached cells. Ethanol induced apoptotic positive controls cells were always used to set up compensations and quadrants. Detached cells, attached cells, and apoptotic controls were washed twice with cold PBS and were resuspended in 100 μ l of 1 $\times$ Binding Buffer (BD Biosciences, San Diego, CA). They were stained with 10 μ l of a 50 μ g/ml stock of propidium iodine and 5 μ l of FITC annexin V (Annexin V-FITC apoptosis detection kit I, BD Bioscience), and were added to TRUcount tubes (BD Bioscience). The tubes were gently vortexed and incubated for 15 min at 25°C in the dark. 400 μ l of 1 $\times$ Binding Buffer was then added to each tube and

analyzed by flow cytometry within 1 hour. Phosphatidylserine is externalized in apoptosis [19], therefore, Annexin V staining of phosphatidylserine on propidium iodide (PI) negative cells was used to quantify apoptosis. Dual staining of Annexin V and PI identified late stage apoptotic or necrotic cells, as both dyes are membrane impermeable and can only enter and stain cells with compromised membrane integrity. PI staining in the absence of annexin V staining identifies necrotic material. Absolute numbers of cells were calculated for detached and attached conditions, and results were expressed as percentage difference from control.

For qRT-PCR, RNA was isolated from the lysed cells, cDNA synthesized, and 9 genes were investigated: Caspase-3, Collagen types I & II, Elastin, MMP1, Proliferating Cell Nuclear Antigen (PCNA), Aggrecan, and SOX9 (Taqman, Applied Biosystems, Bedford, MA). Relative gene expression was calculated using the comparative Ct method normalized to cells in the control group and 18S.

For immunohistochemistry, cells were stained with a primary antibody against paxillin (BD Bioscience, Clone 349, 1/500) and an Alexa 594-conjugated secondary antibody (Invitrogen, 1/500) to indicate attachment to the ECM through focal adhesions. Cells were counterstained with DAPI (Invitrogen, 1:1000) and Alexa 488-phalloidin (Invitrogen, 1/100) to visualize the nucleus and F-actin, respectively.

The 2D discrete Fourier Transform (DFT) was used to calculate orientation distributions of control and stretched cells [20] using a custom Matlab script. An autocorrelation function was used to determine if there was non-randomness in the resulting Fourier transform. It was determined that all three stretch conditions displayed a Gaussian distribution, while the control cells were random. A Gaussian distribution was then fit to the stretch data using a least square routine to characterize symmetric distributions [21]. The mean of the distribution was used to characterize the maximum angle of alignment within an image, while the standard deviation was used as an index to characterize the overall variance within an image.

Experiments with micropatterned collagen to restrict reorientation adaptation

A technique akin to micromolding in capillaries [22] was used to coat stripes of collagen type I to restrict the adaptive reorientation mechanism of the cells in response to high strains (Figure 1). A master mold was created using standard soft lithography techniques [22]. Briefly, a photomask was created with AutoCad (width of features = 40 μm , length = 20 mm, Autodesk, San Rafael, CA) and printed (Advance Reproductions Corporation, North Andover, MA). A silicon wafer was spin coated with a uniform, 50–80 μm -thick layer of SU-8 2035 negative photoresist (MicroChem, Newton, MA). The photoresist was exposed to ultraviolet light through the photomask to create a positive relief of the pattern using the following protocol: soft bake at 75°C for 2 minutes and 105°C for 7 minutes, expose for 15 seconds 100% power (IntelliRay 400, UviTron) and post exposure bake at 75°C for 1 minute and 105°C for 5 minutes. Non-exposed photoresist was removed with SU8 developer (MicroChem). PDMS stamps (Sylgard 184 elastomer base with the curing agent 10:1, Dow-Corning, Midland, MI) were cured against the master and then removed, trimmed to size, and compressed against the STREX silicone elastic membranes. A corona was used to ionize the air to create a plasma around the electrode increasing the wettability of the substrate and PDMS, allowing a collagen type I fitc-conjugated solution (at a higher concentration 100 $\mu\text{g}/\text{mL}$ overnight at 4°C) solution to coat the surface through the microfabricated channels. Regions in contact with the PDMS did not have collagen type I and therefore did not promote cell attachment. The collagen was removed and the surface was rinsed and incubated with 3% BSA (in PBS) at 37°C for 30 minutes. Cells were seeded on the membrane as described earlier, with the modification that after 30 minutes the cells remaining in the supernatant were removed and the surface was washed twice to ensure that

only cells with good adhesion to the collagen remained. After 24 hours the cells were stretched at 1 Hz, 20% strain for 24 hours.

The small number of cells on the micropatterned membranes required apoptotic cells to be quantified using immunohistochemistry for cleaved caspase-3, a marker of apoptosis [23]. Cells were incubated with 10% neutral buffered formalin (30 minutes) and a 10% normal goat block (Vector Laboratories, Burlingame, CA, in 1% BSA/PBS, 30 minutes) and incubated overnight with a cleaved caspase-3 primary antibody (AB3623, Millipore, Temecula, CA, 1:250 in 1% BSA/PBS). The cells were stained with a goat anti-rabbit Alexa 546-conjugated secondary antibody (Invitrogen, 1/800), DAPI (1:1000), and Alexa 488-phalloidin (1/100). At least 20 cells were imaged for each chamber condition and percent apoptotic cells was calculated [positive caspase-3 staining/ total cells $\times$ 100].

Statistics

All data was assessed for normality using the Ryan–Joiner test. A one-way ANOVA was done for flow cytometry, orientation results, and cleaved caspase-3 counts with a Fisher's post-hoc test. A one sample t-test was done on the $\Delta\Delta C_T$ values to assess if there were any differences from control for each strain amplitude ($\Delta\Delta C_T=0$). All statistics were done with Minitab software (Version 16, State College, PA) and significance was always defined as $p < 0.05$.

Results

Cell stretch on uniform collagen substrates resulted in a small trend of increased early stage apoptosis (Annexin V positive cells) in the detached cell population (affecting $<1\%$ of the cells) that was not significant (Supplement 1A&B). No differences were observed for late stage apoptosis/necrosis (dual Annexin V and PI positive cells). CTS had variable effects on mRNA expression, with no significant changes (Supplement 1C). The largest effect of stretching on uniformly coated collagen substrates was cellular reorientation (Figure 2). Unstretched cells exhibited a random pattern of cellular orientations as quantified with DFT and autocorrelation (cross-correlation of a signal with itself), while all of the stretched cells had a preferential orientation that exhibited a Gaussian pattern. The mean of the Gaussian curve fit quantified maximum angle of alignment (where angles ranged from 0–180 degrees from the direction of stretch) and standard deviation indicated relative variance within the image. No significant differences in orientation were detected between stretch groups for both the mean (10%: $86.4^\circ \pm 5.8^\circ$, 15%: $102.0^\circ \pm 12.4^\circ$, 20%: $86.0^\circ \pm 15.9^\circ$, $p = 0.274$) and standard deviation of the Gaussian curve fit (10%: $67.6^\circ \pm 19.4^\circ$, 15%: $47.2^\circ \pm 19.0^\circ$, 20%: $128.4^\circ \pm 129.3^\circ$, one way ANOVA, $p = 0.273$). Cellular reorientation in response to stretch was associated with a clear redistribution of actin stress fibers (Phalloidin), and focal adhesion sites (Paxillin) perpendicular to the direction of stretch and some nuclear (DAPI) reorientation. Therefore, when cells were allowed to reorient little apoptosis and changes in gene expression occurred.

Cleaved caspase-3 staining on uniformly coated chambers was minimal for control and stretched cells ($< 8\%$, Figure 3), supporting the flow cytometry results. There was also little observed cleaved caspase-3 staining ($< 5\%$) when the cells were grown without stretch on stripes of collagen that prevented reorientation. Stretching the cells on the striped substrate, however, resulted in a significant increase in percentage of cells positive for cleaved caspase-3 (at least 20 cells for each group, one way ANOVA, $p < 0.001$, average of 90%). The restricted stretched cells also had an observed rounded, or 'blebbed' morphology.

Discussion

AF cells from surgical samples underwent cyclic uniaxial stretch to test the hypothesis that AF cells are capable of reorienting in response to high stretch magnitudes but undergo apoptosis when their adaptive reorientation is restricted. Results indicated that AF cells responded to 24 hrs of cyclic stretch on uniformly coated collagen surfaces by redistribution of actin and focal adhesions perpendicular to the stretch direction without notable apoptosis or matrix gene expression. However, AF cells were highly susceptible to stretch-induced apoptosis when adaptive reorientation was inhibited by restricting cell adhesion surfaces to 40 μm stripes parallel to the direction of stretch. Results support the concept that cells must maintain their ability to reorient in order to adapt to high strain environments. Specifically, degenerated AF cells have the capacity to reorient, but stretch-induced apoptosis may be a contributor to cell death during degeneration due to high strains concentrated around delaminations, herniations and other structural defects, since cells would have a reduced capacity to adapt in the more highly crosslinked, fibrotic, and compacted degenerated AF tissue.

Uniaxial cyclic stretching of 2D cell cultures causes the actin cytoskeleton and focal adhesions of many cell types to align perpendicular to the direction of stretch [15,16,17]. Different cell types orient at different rates, for instance, fibroblasts orient within 3 hours [24] and vascular endothelial cells orient within 6 hours [25], while some cell types, including neutrophils, do not reorient [26]. Reorientation perpendicular to direction of stretch is a mechanism for both remodeling and intracellular signaling by a structural and biochemical adaption to mechanical stretch [27] related to Rho activity and the extent of cell stretching [17] that can be predicted by minimizing the potential energy of the stress fiber-focal adhesion complex [28]. All three strain amplitudes (10, 15, and 20%) in this study had similar effects on human AF cells and resulted in the same reorientation patterns. It is likely that minimizing the potential energy by cytoskeletal remodeling is an important adaptation that allows cells to adapt to high strains observed when collagen fibers undergo tension as occurs in bending and torsion in the IVD [29]. However, age and degeneration related changes in the collagen network, including increases in crosslink density [30,31], AF matrix remodeling [3], tissue compaction [32], and compromised lamellar structure [33] may limit the capacity for cells to reorient in response to high strains.

When cells were restricted from reorienting cleaved caspase-3 staining and a rounded cell morphology, often called “blebs” was observed. Cleavage of the effector protein, caspase-3 is critical for dismantling of the cell and the development of apoptotic bodies [23], while the “bleb” morphology is a result of the cell detaching from the underlying substrate as it undergoes apoptosis [34]. Therefore the restricted orientation model is consistent with the stretch-induced apoptosis observed in other studies on AF cells [12,13,14]. We highlight that differences in applied mechanical stimuli (uniaxial vs. biaxial) strongly influence cellular adaptive mechanisms. Excessive biaxial stretch leads to apoptosis [12,13,14] and in this study excessive uniaxial stretch leads to reorientation. However, when cells are unable to reorient to minimize strain energy (i.e., equibiaxial strain and/or uniaxial strain with restricted adhesion area) they are susceptible to apoptosis at high stretch magnitudes.

This *in vitro* model has the advantage of precise boundary conditions, to analyze the effect of stretch without confounding factors present in the more complex *in situ* environment, however, direct interpolation to the native environment is speculative. For example, pure tensile uniaxial strain will not always occur *in situ* [35]. Under healthy conditions AF surface strains up to 6% were measured along the fiber direction [10], therefore the strains chosen are high and are likely to only be present in degenerated states near structural defects, and injury sites. While these differences exist, we believe the inferences remain that

cells with restricted motility are at greater risk of apoptosis. Finally, the cells used were obtained from degenerated human IVDs, which have an altered gene expression pattern compared to cells obtained from healthy IVDs [36]. However, it is likely that healthy AF cells are able to reorient in response to stretch since degenerated cells did.

In conclusions, high uniaxial strains on a uniformly coated collagen surface promoted cytoskeletal reorientation without apoptosis and with minimal alterations in gene expression. However, AF cells were vulnerable to stretch-induced apoptosis when cellular reorientation was restricted. Results indicate that degenerated AF cells have the capacity to adapt to stretch but that high strains and inhibited cellular reorientation, as might occur during degeneration, puts cells at risk of stretch-induced apoptosis. These AF cells exemplify, on a cellular-level, the famous quote of John Stark “Live free or die!”

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work made possible by funding from the NIH (R01AR051146, R01AR057397) and by the AO Research Fund (project F-09-10I) of the AO Foundation.

References

- [1]. Bibby SR, Urban JP. Effect of nutrient deprivation on the viability of intervertebral disc cells. *Eur Spine J.* 2004; 13:695–701. [PubMed: 15048560]
- [2]. Bibby SR, Jones DA, Ripley RM, Urban JP. Metabolism of the intervertebral disc: effects of low levels of oxygen, glucose, and pH on rates of energy metabolism of bovine nucleus pulposus cells. *Spine (Phila Pa 1976).* 2005; 30:487–496. [PubMed: 15738779]
- [3]. Roughley PJ. Biology of intervertebral disc aging and degeneration: involvement of the extracellular matrix. *Spine (Phila Pa 1976).* 2004; 29:2691–2699. [PubMed: 15564918]
- [4]. Adams MA, Roughley PJ. What is intervertebral disc degeneration, and what causes it? *Spine (Phila Pa 1976).* 2006; 31:2151–2161. [PubMed: 16915105]
- [5]. O'Connell GD, Vresilovic EJ, Elliott DM. Human intervertebral disc internal strain in compression: the effect of disc region, loading position, and degeneration. *J Orthop Res.* 2011; 29:547–555. [PubMed: 21337394]
- [6]. Tsantrizos A, Ito K, Aebi M, Steffen T. Internal strains in healthy and degenerated lumbar intervertebral discs. *Spine (Phila Pa 1976).* 2005; 30:2129–2137. [PubMed: 16205337]
- [7]. O'Connell GD, Malhotra NR, Vresilovic EJ, Elliott DM. The Effect of Discectomy and the Dependence on Degeneration of Human Intervertebral Disc Strain in Axial Compression. *Spine (Phila Pa 1976).* 2011
- [8]. Michalek AJ, Buckley MR, Bonassar LJ, Cohen I, Iatridis JC. The effects of needle puncture injury on microscale shear strain in the intervertebral disc annulus fibrosus. *Spine J.* 2010; 10:1098–1105. [PubMed: 20971041]
- [9]. Korecki CL, Costi JJ, Iatridis JC. Needle puncture injury affects intervertebral disc mechanics and biology in an organ culture model. *Spine.* 2008; 33:235–241. [PubMed: 18303454]
- [10]. Stokes IA. Surface strain on human intervertebral discs. *J Orthop Res.* 1987; 5:348–355. [PubMed: 3625358]
- [11]. Broberg KB. On the mechanical behaviour of intervertebral discs. *Spine (Phila Pa 1976).* 1983; 8:151–165. [PubMed: 6857386]
- [12]. Rannou F, Lee TS, Zhou RH, Chin J, Lotz JC, Mayoux-Benhamou MA, Barbet JP, Chevrot A, Shyy JY. Intervertebral disc degeneration: the role of the mitochondrial pathway in annulus fibrosus cell apoptosis induced by overload. *Am J Pathol.* 2004; 164:915–924. [PubMed: 14982845]

- [13]. Zhang YH, Zhao CQ, Jiang LS, Dai LY. Cyclic stretch-induced apoptosis in rat annulus fibrosus cells is mediated in part by endoplasmic reticulum stress through nitric oxide production. *Eur Spine J.* 2011; 20:1233–1243. [PubMed: 21336971]
- [14]. Zhang YH, Zhao CQ, Jiang LS, Dai LY. Lentiviral shRNA silencing of CHOP inhibits apoptosis induced by cyclic stretch in rat annular cells and attenuates disc degeneration in the rats. *Apoptosis.* 2011; 16:594–605. [PubMed: 21479580]
- [15]. Ritchie AC, Wijaya S, Ong WF, Zhong SP, Chian KS. Dependence of alignment direction on magnitude of strain in esophageal smooth muscle cells. *Biotechnol Bioeng.* 2009; 102:1703–1711. [PubMed: 19170241]
- [16]. Boccafroschi F, Bosetti M, Gatti S, Cannas M. Dynamic fibroblast cultures: response to mechanical stretching. *Cell Adh Migr.* 2007; 1:124–128. [PubMed: 19262127]
- [17]. Kaunas R, Nguyen P, Usami S, Chien S. Cooperative effects of Rho and mechanical stretch on stress fiber organization. *Proc Natl Acad Sci U S A.* 2005; 102:15895–15900. [PubMed: 16247009]
- [18]. Thompson JP, Pearce RH, Schechter MT, Adams ME, Tsang IK, Bishop PB. Preliminary evaluation of a scheme for grading the gross morphology of the human intervertebral disc. *Spine (Phila Pa 1976).* 1990; 15:411–415. [PubMed: 2363069]
- [19]. Waring P, Lambert D, Sjaarda A, Hurne A, Beaver J. Increased cell surface exposure of phosphatidylserine on propidium iodide negative thymocytes undergoing death by necrosis. *Cell Death Differ.* 1999; 6:624–637. [PubMed: 10453073]
- [20]. Marquez JP. Fourier analysis and automated measurement of cell and fiber angular orientation distributions. *International Journal of Solids and Structures.* 2006; 43:6413–6423.
- [21]. Swindale NV. Orientation tuning curves: empirical description and estimation of parameters. *Biol Cybern.* 1998; 78:45–56. [PubMed: 9518026]
- [22]. Kim E, Xia Y, Whitesides GM. Micromolding in Capillaries: Applications in Materials Science. *Journal of the American Chemical Society.* 1996; 118:5722–5731.
- [23]. Porter AG, Janicke RU. Emerging roles of caspase-3 in apoptosis. *Cell Death Differ.* 1999; 6:99–104. [PubMed: 10200555]
- [24]. Neidlinger-Wilke C, Grood E, Claes L, Brand R. Fibroblast orientation to stretch begins within three hours. *J Orthop Res.* 2002; 20:953–956. [PubMed: 12382959]
- [25]. Iba T, Sumpio BE. Morphological response of human endothelial cells subjected to cyclic strain in vitro. *Microvasc Res.* 1991; 42:245–254. [PubMed: 1779881]
- [26]. Haston WS, Shields JM, Wilkinson PC. The orientation of fibroblasts and neutrophils on elastic substrata. *Exp Cell Res.* 1983; 146:117–126. [PubMed: 6861903]
- [27]. Kaunas R, Usami S, Chien S. Regulation of stretch-induced JNK activation by stress fiber orientation. *Cell Signal.* 2006; 18:1924–1931. [PubMed: 16581230]
- [28]. Stamenovic D, Lazopoulos KA, Pirentis A, Suki B. Mechanical Stability Determines Stress Fiber and Focal Adhesion Orientation. *Cell Mol Bioeng.* 2009; 2:475–485. [PubMed: 20376295]
- [29]. Krismer M, Haid C, Rabl W. The contribution of anulus fibers to torque resistance. *Spine.* 1996; 21:2551–2557. [PubMed: 8961442]
- [30]. Sivan SS, Tsitron E, Wachtel E, Roughley P, Sakkee N, van der Ham F, Degroot J, Maroudas A. Age-related accumulation of pentosidine in aggrecan and collagen from normal and degenerate human intervertebral discs. *Biochem J.* 2006; 399:29–35. [PubMed: 16787390]
- [31]. Pokharna HK, Phillips FM. Collagen crosslinks in human lumbar intervertebral disc aging. *Spine (Phila Pa 1976).* 1998; 23:1645–1648. [PubMed: 9704370]
- [32]. Iatridis JC, Setton LA, Foster RJ, Rawlins BA, Weidenbaum M, Mow VC. Degeneration affects the anisotropic and nonlinear behaviors of human anulus fibrosus in compression. *J Biomech.* 1998; 31:535–544. [PubMed: 9755038]
- [33]. Bernick S, Walker JM, Paule WJ. Age changes to the anulus fibrosus in human intervertebral discs. *Spine (Phila Pa 1976).* 1991; 16:520–524. [PubMed: 1711242]
- [34]. Lu Q, Rounds S. Focal adhesion kinase and endothelial cell apoptosis. *Microvasc Res.* 2011

- [35]. Bruehlmann SB, Matyas JR, Duncan NA. ISSLS prize winner: Collagen fibril sliding governs cell mechanics in the annulus fibrosus: an in situ confocal microscopy study of bovine discs. *Spine (Phila Pa 1976)*. 2004; 29:2612–2620. [PubMed: 15564909]
- [36]. Gruber HE, Hoelscher GL, Hanley EN Jr. Annulus cells from more degenerated human discs show modified gene expression in 3D culture compared with expression in cells from healthier discs. *The Spine Journal*. 2010; 10:721–727. [PubMed: 20650410]

Highlights

- Human annulus fibrosus cells reorient perpendicular to cyclic uniaxial stretch.
- By reorienting cells do not undergo apoptosis in response to high strains.
- Restricting cells from reorienting in response to stretch results in apoptosis.

Figure 1. Fabrication of 40 μm collagen stripes to restrict cellular reorientation

A silicon wafer was spin coated with a uniform layer of negative photoresist. UV radiation and a photomask was used to selectively expose the photoresist. Areas that were not exposed to the UV radiation become soluble in developer leaving only the desired pattern. PDMS was then cast against the photoresist master to create a relief of the 3D topography. The PDMS containing the microchannel features was then brought in contact with the silicone elastic substrate. A FITC labeled collagen type I solution submerged the PDMS overnight, confirming the collagen stripe pattern (scale bar= 100 μm).

Figure 2. Cytoskeletal remodeling in response to stretch involved reorientation perpendicular to the direction of stretch (90°) through the redistribution of actin and focal adhesions

The discrete fourier transform (DFT) was used to quantify angular distribution of cells. Representative images of unstretched (A) and 20% stretched (B) cells with the corresponding DFT (C and D, respectively), where the location of peaks in the amplitude corresponds to preferential orientation of cells in that direction. Autocorrelation demonstrated that control images were random (E) while the stretch groups displayed a Gaussian pattern (F). Gaussian curves were fit to the DFT of stretched cells with the mean indicating angle of alignment, and standard deviation the amount of variance within an image. Cellular orientation was clear on imaging of control (G) and stretched (H) cells stained with DAPI (blue), Phalloidin (green), and anti-paxillin (red). Scale bar= $100\mu\text{m}$ in all images, the stretch direction (B,H) was horizontal, represented by the black arrow, and white arrows (G,H) indicate focal adhesion sites randomly around the control cells and redistributed to the poles of stretched cells perpendicular to stretch.

Figure 3. Restricting the adaptive reorientation mechanism, and applying 20% cyclic stretch results in positive staining of cleaved caspase-3, indicating the initiation of an apoptotic pathway Images of control and stretched cells on either a uniformly collagen coated substrate or restricted to 40 μm collagen stripes stained with DAPI (blue), cleaved caspase-3 (red), and Phalloidin (green) (A). The stretch direction was horizontal and is represented by a white arrow on the images. Scale bars are 50 μm . There was significantly more cleaved caspase-3 staining in the restricted orientation group subjected to 20% cyclic stretch (B, one way ANOVA, $p < 0.001$, Fisher's post hoc test).