Isolation and mapping of a polymorphic DNA sequence (pCJ52.94T1) on chromosome 16 [D16S159]

C.Julier*, Y.Nakamura, G.M.Lathrop*, J.-M.Lalouel and R.White

Howard Hughes Medical Institute, University of Utah Medical School, Salt Lake City, UT 84132, USA

SOURCE/DESCRIPTION: A 3.8 kb TaqI fragment from cosmid CJ52.94, isolated from a library constructed from a mouse/human somatic cell hybrid that contains a chromosomellq-16q derivative as the only human chromosome (1), was subcloned into the AccI site of pUC18.

POLYMORPHISM: TaqI identifies a 5 allele insertion/deletion type polymorphism, with bands at 3.7 kb (T1), 3.6 kb (T2), 3.3 kb (T3), 3.2 kb (T4) and 2.9 kb (T5). MspI and PstI, and probably other restriction enzymes, also detect this polymorphism.

FREQUENCY: Estimated from 113 unrelated Caucasians.

T1 : 0.15
T2 : 0.06
T3 : 0.07
T4 : 0.04
T5 : 0.68

NOT POLYMORPHIC FOR: unknown.

CHROMOSOMAL LOCALIZATION: pCJ52.94T1 has been assigned to chromosome 16 by multipoint linkage analysis (2) with loci known to map on this chromosome (1).

MENDELIAN INHERITANCE: Codominant segregation of the TaqI RFLPs was observed in 57 three generation families.

PROBE AVAILABILITY: Contact C.J.

OTHER COMMENTS: This probe may contain repetitive DNA sequences, and should be used in the presence of an excess of total human DNA in the prehybridization and hybridization mixtures.

REFERENCES:

1. C. Julier et al., in preparation
2. G.M. Lathrop et al., Am. J. Hum. Genet. 37:482-498 (1985)

*Present address: CEPH, 27 rue Juliette Dodu, 75010 Paris, France