Isolation and mapping of a polymorphic DNA sequence (CJ52.112) on chromosome 16 [D16S158]

C.Julier*, Y.Nakamura, G.M.Lathrop*, J.-M.Lalouel and R.White

Howard Hughes Medical Institute, University of Utah Medical School, Salt Lake City, UT 84132, USA

SOURCE/DESCRIPTION: Cosmid CJ52.112 was isolated from a library constructed from a mouse/human somatic cell hybrid that contains a chromosomellq-16q derivative as the only human chromosome (1). The Sau3A partial digest was inserted into pWe vector.

POLYMORPHISM: TaqI identifies a two-allele RFLP, with polymorphic fragments at 5.9 kb (T1) and 3.6 and 2.3 kb (T2)

FREQUENCY: Estimated from 87 unrelated Caucasians.

T1 : 0.92

T2 : 0.08

NOT POLYMORPHIC FOR: MspI and PstI.

CHROMOSOMAL LOCALIZATION: CJ52.112 has been assigned to chromosome 16 by multipoint linkage analysis (2) with loci known to map on this chromosome (1).

MENDELIAN INHERITANCE: Codominant segregation of the TaqI RFLPs was observed in 44 three generation families.

PROBE AVAILABILITY: Contact C.J.

OTHER COMMENTS: This probe contains repetitive DNA sequences, and needs to be used in the presence of an excess of total human DNA in the prehybridization and hybridization mixtures.

REFERENCES:

1. C. Julier et al., in preparation
2. G.M. Lathrop et al., Am. J. Hum. Genet. 37:482-498 (1985)

*Present address: CEPH, 27 rue Juliette Dodu, 75010 Paris, France