Nucleotide sequence of a late developmentally regulated prespore-enriched *Dictyostelium discoideum* gene

David C.C. Widdowson, Pippa S. Jagger and B. David Hames

Department of Biochemistry, University of Leeds, Leeds LS2 9JT, UK
Submitted September 11, 1989

EMBL accession no. X16124

Expression of the 3F gene is restricted to prespore cells during late stages of *D. discoideum* development and is partially cyclic AMP responsive (1). We have sequenced a 3F cDNA clone constructed using mRNA from the 18-20h stage of development. Comparison between the molecular weight of the predicted amino acid sequence (23.5kD) and that estimated previously (1) by hybrid rescue/SDS-PAGE (27kD) indicates that sequence encoding no more than 33 amino acids is missing from the 3' end of the cDNA clone. The cDNA sequence predicts a putative leader sequence (overlined) and cleavage site, as well as an N-linked glycosylation site (*). Discounting the leader, the protein is highly hydrophilic (71 charged residues) and is characterised towards its C-terminus by three near-perfect repeats of the 9 amino acid sequence SPK(S/T)DAKEA and the beginning of a fourth repeat (overlined). The 5' untranslated region contains 8 clustered copies of the sequence (AAC) followed immediately by 6 copies of the sequence (AAT) (underlined). The repetitive sequence (AAC)_n occurs in several developmentally regulated genes of *D. discoideum* expressed maximally during aggregate formation and has been proposed as a potential regulatory element in the coordination of gene expression (2,3). The (AAT) repeat has not been reported previously but may play an analogous role either alone or in conjunction with the (AAC)_n element.

```

TTTTTTTTTATAAATACTACAAAAACAACACACAACAACAACAATAATAATAATAATACTTTTAATA
                                     M K L L S K L I L T L
ATATTACAAAAATAAAATTATAATTAATTAATAAAAAAAAAAATGAATTATTATCAAAATTAATTTAACATTG
A L A T Y A S A S E N F R Y I N W D N P P H D V T
GCATTAGCAACATATGCTAGTGAAGTGAATTTTAGATATATTAATTGGGATAATCCACCACATGATGTAACA
F Y E G D V L Q F T T N E G R N * S T I T L I S D T
TTTTTAGAAGGTGATGTTTACAATTACACCAATGAAGGTAGAAATTCACCATTAACATCATTAGTGATACT
E N G D K S F D G V L N E D Q R S F V Q K A L P P
GAAAATGGTGATAAATCATTCGATGGTGTCTCAACGAAGATCAAAGATCATTTGTTCAAAAAGCATTACCACCT
G R Y T F K D L N S G S K S I I R V K E S K E L A
GGCAGATACACTTTCAAAGATTAAATTCGGTAGTAAATCAATTATTAGAGTTAAAGAATCTAAAGAATTAGCT
K E V R P I D R L K D N A D A A N T E N A Q K S P
AAAGAAGTCAGACCAATTGATCGTTTAAAGGATAATGCTGATGCTGCCAATACTGAAAATGCTCAAAAATCACCA
N T Q S T Q K G S P K S D A K E A S P K T D A K E
AATACTCAATCCACTCAAAAAGGTTCTCCAAAATCCGATGCAAAAGAAGCATCACCAAAACCTGATGCAAAAGAA
A S P K S D A K E A S P K T D T K Q G S S P K T D
GCATACCAAAATCAGATGCTAAAGAAGCATCACCAAAACCTGATACCAACAAGGCTCCTCTCCAAAACCTGAT
T K S S T Q K P S S S S D S S K A K A E A N T A A
ACTAAATCATCAACTCAAAAACCATCTTCATCTCCGATTCTTCAAAAGCAAAGCTGAAGCCAATACCGCAGCC
N N E E
AACAAATGAAGAA

```

References

1. Corney, A. J. *et al.* (1989) Mol. Microbiol. (submitted).
2. Kimmel, A. R. and Firtel, R. A. (1985) Mol. Cell. Biol., **5**, 2123-2130.
3. Saxe, S.A. and Kimmel, A.R. (1988) Dev. Genet., **9**, 259-265.