Published in final edited form as:

Bioorg Med Chem Lett. 2011 November 15; 21(22): 6788–6792. doi:10.1016/j.bmcl.2011.09.034.

Synthesis of Purine Modified 2'-C-Methyl Nucleosides as Potential Anti-HCV Agents

Hong-wang Zhang^{a,b}, Longhu Zhou^{a,b}, Steven J. Coats^c, Tamara R. McBrayer^{b,c}, Phillip M. Tharnish^{b,c}, Lavanya Bondada^{a,b}, Mervi Detorio^{a,b}, Sarah A. Amichai^{a,b}, Melissa D. Johns^{a,b}, Tony Whitaker^c, and Raymond F. Schinazi^{a,b}

^aCenter for AIDS Research, Laboratory of Biochemical Pharmacology, Department of Pediatrics, Emory University School of Medicine, Atlanta, GA 30322, USA

^bVeterans Affairs Medical Center, Decatur, GA 30033, USA

^cRFS Pharma, LLC, 1860 Montreal Road, Tucker, GA 30084, USA

Abstract

Based on the anti-hepatitis C activity of 2'-C-methyl-adenosine and 2'-C-methyl-guanosine, a series of new modified purine 2'-C-methyl nucleosides was prepared as potential anti-hepatitis C virus agents. Herein, we report the synthesis of both 6-modified and 2-modified purine 2'-C-methyl-nucleosides along with their anti-HCV replication activity and cytotoxicity in different cells.

Hepatitis C virus (HCV) infects more than 200 million individuals worldwide according to the World Health Organization. In addition, an estimated 3 to 4 million people contract HCV each year. Long-term infection can lead to chronic liver disease, such as cirrhosis of the liver or hepatocellular carcinoma. Until recently the standard of care therapy employed interferon- α (IFN) often in combination with the nucleoside analog ribavirin. The impact on standard of care by approval of the two HCV protease inhibitors Incivek and Victrelis remains unclear as both drugs require response-guided therapy regimens that can shorten the duration of IFN therapy in infected persons with an early viral response from 48 weeks to as few as 24 weeks. In addition, serious side effects and limited efficacy emphasize the urgent need for improved therapeutic agents.¹ Moreover, there is no established vaccine for HCV. As a result, there is an urgent need for safe and effective therapeutic agents that combat HCV infection and that have high genetic barrier to resistance.

Among the most successful preclinical and clinical nucleosides with potent HCV NS5B polymerase inhibition are 2'-C-methyl analogs.² 2'-C-Methyl adenosine, **1** and 2'-C-methyl guanosine, **2** (Fig. 1) display selective *in vitro* anti-HCV activity as non-obligatory chain terminators toward RNA elongation. However, *in vivo* these analogs display poor bioavailability due to deamination of the adenine base and purine nucleoside phosphorylase (PNP) promoted glycosidic bond cleavage in the case of **1** and poor cellular uptake and/or inefficient phosphorylation for **2**.³

Correspondence to: Raymond F. Schinazi.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Base-modification of nucleoside analogs has been successful in the development of potent antiviral agents.⁴ Within the 2'-C-methyl nucleoside series, the 7-deaza modification of the adenosine analog results in a significant improvement of *in vivo* anti-HCV activity.⁵ In this study, novel purine modified 2'-C-methyl nucleoside analogs were prepared which explored the steric, electronic, and hydrogen bonding effects on anti-HCV activity as measured by a cell-based replication assay.

The synthesis of 2'-C-methyl-2-amino-6-methyl purine analogs is outlined in Scheme 1. Compound **3**, as a single isomer, was readily prepared in three sugar modification steps⁵ from commercially available starting materials. Condensation of **3** with 2-amino-6-chloropurine in the presence of DBU and TMSOTf at 80 °C for 5 h gave the 6-chloro nucleoside **4** as the pure β -anomer in 92% yield. A modification to the standard Vorbrüggen conditions involving the addition of TMSOTf at -78 °C improved the reaction yield from around 60%^{3a} to over 90%. Cross-coupling reactions have been used successfully for the introduction of carbon-linked substituents into purine moieties.⁶ Hence compound **4** was allowed to react with trimethylaluminum under Pd(PPh₃)₄ catalyzed conditions in THF to give the corresponding 6-methyl-purine derivative **5** in 54% yield. Subsequently, the benzoyl groups were removed to afford the 6-methyl purine nucleoside analog **6**.⁷ Monophosphate prodrug, (aryloxy)phosphoramidate **7**, was prepared by following Uchiyama's procedure in the presence of *tert*-butyl magnesium chloride in THF.^{8,9}

The 6-phosphonate substituted purine is an attractive target because the phosphonate group may mimic the hydrogen bonding characteristics of a 6-amino or 6-hydroxyl group of endogenous nucleoside bases and thus may result in nucleoside analogs that are inhibitors of adenosine deaminase (ADA).¹⁰ Application of the Arbuzov reaction to 6-chloro purine **4** by treatment with triethyl phosphite afforded 6-phosphonate **8** in 78% yield (Scheme 2).¹¹ Removal of the benzoyl groups with saturated NH₃ in ethanol or NaOEt in ethanol provided **9**¹² and **10** respectively.

A series of 1,2,4-triazolo [5,1-*i*] adenine derivatives were identified as potent adenosine A_{2a} receptor antagonists.¹³ Application of these interesting base modifications to a 2'-Me sugar nucleoside provides potential HCV inhibitors. Among many synthetic approaches for preparation of these purine base modifications,¹⁴ an efficient cyclization method utilizing N, O-bis-(trimethylsilyl)acetamide (BSA) to undergo a dehydrative rearrangement was chosen as shown in Scheme 3.

The ipso displacement of 6-chloro group of compound **4** by acetohydrazide or 2-furoic hydrazide afforded compounds **11a** and **11b**, which are used for the next step without purification. Heating with BSA for 5 h furnished the desired tricycles **12** *via* a dehydrative rearrangement process.¹³ After removal of the benzoyl protecting groups, **13a** and **13b** were isolated in 50% and 90% yield, respectively.

The 6-H nucleoside **15** was isolated in 21% yield during the preparation of **12a** (Fig. 2). Formation of **15** can be explained by a base promoted Wolff-Kishner reduction.¹⁵ This postulate is supported by the detection of the 6-hydrazo nucleoside **14** by LC/MS during the preparation of **11a**. During the BSA promoted dehydrative rearrangement reaction, intermediate **14** was converted to **15** while the intermediate **11a** proceeded to **12a**. In a confirmation reaction, 6 equivalents of hydrazine were allowed to react with compound **4** at 110 °C for 2 days. The reduced nucleoside **15** was cleanly formed without removal of the benzoyl groups.

Previous studies have found 2-position modified purines with potent antiviral activity and/or a reduction of toxicity.¹⁶ The 2-hydrazine-, 2-azido- and 2-triazole-purine nucleosides were

targeted as they potentially offer a variety of steric, electronic, and hydrogen bonding interactions which may enhance recognition by HCV NS5B polymerase. Treatment of the 2,6-dichloro purine nucleoside **16** with methanolic ammonia removed the benzoyl protecting groups with concomitant amination and methoxylation to afford 6-aminopurine **17** and 6-methoxypurine **18** respectively (Scheme 4). Nucleophilic substitution of the 2-chloro group of **17** with hydrazine hydrate gave the 2-hydrazine substituted purine nucleoside **19**.¹⁷ Treatment of **19** with sodium nitrite in acetic acid provided a 63% yield of the 2'-azidopurine **20**. Based on analysis of the ¹H-NMR, compound **20** exists in both an azido and an N1 tetrazole tautomeric form.¹⁸ 1,3-Dipolar cycloaddition of azide **20** with ethynyltrimethylsilane through a Cu (I)-catalyzed 1,3-cycloaddition reaction generated the triazole analogs **21**¹⁹ and compound **22**.²⁰ Compound **21** is formed by desilylation during the cycloaddition reaction. Compound **21** may also be prepared from **22** by treatment with aqueous HF in THF and formation of **21** during the cycloaddition step may be suppressed by addition of BSA to the reaction mixture.²¹

The 2-hydroxylamino substitution was selected for evaluation due to its structural and electronic similarity to hydrazine. As shown in Scheme 5 the 6-chloropurine **23** was nitrated with tetrabutylammonium nitrate (TBAN) and trifluoroacetic anhydride (TFAA) to afford the 2-nitro analog **24** in 79% yield.²² Nucleophilic substitution with sodium azide produced compound **25** in 99% yield. Side reactions and lower yields occurred when more than 1 equivalent of sodium azide was utilized in this reaction. Careful hydrogenation under mild heating afforded 2-hydroxypurines **26** and **27** in modest yields.²³ However, removal of the benzoyl groups of **26** or **27** with NH₃/CH₃OH or sodium methoxide under a variety of conditions failed to give the desired compounds, but instead lead to complex mixtures.

Based on the biological activity of the 4-amino-1*H*-imidazo[4,5-*d*]pyridazin-7(6*H*)-one ring system as a purine isostere,²⁴ 4-amino-1-(2-methyl-β-D-ribofuranosyl)-3*H*-imidazo[4,5-*d*]pyriazin-7-(6*H*)-one, **32** and 7-amino-1-(2-methyl-δ-D-ribofuranosyl)-1*H*-imidazo [4,5-*d*]pyridazin-4-(5*H*)-one, **35** were synthesized (Scheme 6). The Vorbrüggen glycosylation reaction of **4** with ethyl 5(4)-cyano-1*H*-imidazole 4(5)carboxylate **28** gave two regioisomeric nucleosides **29** and **30** in 64% and 36%, respectively. The regioisomeric structures were assigned based on analysis of their ¹H-NMR data and comparison to their ribofuranosyl analogs.²⁵ Compounds **29** and **30** were independently treated with excess *t*-butylamine in ethanol to obtain the deprotected and transesterified **31** and **34** in 91% and 78% yield, respectively. The two compounds were separately reacted with hydrazine, followed by heating with sodium ethoxide in anhydrous ethanol at reflux to yield the respective target nucleoside analogs **32** and **35** in 71% and 62% yield. The unexpected bis-adduct products **33** and **36** were also obtained albeit in low yields of 5% and 12%, respectively.

All of the modified purine 2'-*C*-methyl nucleosides were evaluated for inhibition of HCV RNA replication at 10 μM in Huh7 cells using a subgenomic HCV replicon system.²⁶ Cytotoxicity in Huh7 cells was determined simultaneously with anti-HCV activity by extraction and amplification of both HCV RNA and ribosomal RNA (rRNA).²⁷ To determine the spectrum of activity of the compounds, anti-HIV activity was evaluated *versus* HIV-1_{LAI} in primary human peripheral blood mononuclear (PBM) cells and AZT was used as a positive standard. Cytotoxicity was determined in PBM, human lymphoblastoid CEM, and African Green monkey Vero cells.²⁸ A subset of the compounds were tested for their ability to inhibit and/or act as substrates of adenosine deaminase.²⁹ The antiviral and cytotoxicity results are summarized in Table 1.

In general, nucleosides with the 6-position modifications of the purine base did not display any marked anti-HCV activity (up to 10 μM) or toxicity to Huh7 PBM, CEM, or Vero cell

lines. The lack of a nti-HCV activity or cytotoxicity of these nucleosides may be due to inefficient uptake and/or their inability to be intracellularly metabolized to the corresponding nucleoside triphosphates. However, the monophosphate prodrug **7**, which would bypass the initial phosphorylation step, also did not display any inhibition of HCV replicon RNA replication.

Modification in the 2-position of the purine base proved to be somewhat interesting. The substituted purines 2-Cl,6-NH₂, **17**, 2-Cl,6-OMe, **18**, 2-azido,6-NH₂, **20**, 2-(1*H*-triazole),6-NH₂, **21**, 2-(4-(trimethylsilyl)-1*H*-triazole),6-NH₂, **22** along with the purine isosteres **32–36** all displayed no antiviral activity or cytotoxicity. In contrast, hydrazine **19** possessed anti-HCV activity (85% inhibition at 10 μM; EC₅₀ = 6 μM) with no observed cytotoxicity which encouraged us to prepare its (2*S*)-ethyl(phenoxy)phosphorylamino–propanoate prodrug (Fig. 3). This phosphoramidate prodrug was more active *versus* HCV replication with an EC₅₀ of 0.9 μM. However stability studies of **19** in pH 7.4 phosphate buffer at 23 °C showed complete conversion of **19** to a complex mixture of products after 3 days. One of decomposition products was identified as 2'-*C*-methyladenosine, a known inhibitor of HCV NS5B polymerase.³⁰

The two 2-hydroxyamino nucleosides **26** and **27** displayed broad toxicity against most cell lines tested. Compounds **9**, **10**, **13b**, **21**, **32**, and **35** were tested for their ability to inhibit the ADA catalyzed conversion of adenosine to inosine. These compounds were chosen due to their structural similarity to known ADA inhibitors.^{10,13} Only compound **21** was found effective with an average IC₅₀ = 3.7 ± 0.4 μM. Compounds **9**, **10**, **21**, and **32** were evaluated as potential substrates of adenosine deaminase; however, none underwent a nitrogen to oxygen transformation.

In conclusion, a variety of purine-modified 2'-*C*-methyl nucleosides were synthesized and evaluated as potential anti-HCV agents. Among these synthesized nucleoside analogs, none displayed potent and selective anti-HCV or anti-HIV activity. Interestingly, some of these purine analogs warrant further study as potential inhibitors of adenosine deaminase.³¹

Acknowledgments

This work is supported in part by NIH grant 2P30-AI-050409 (CFAR), 5R37-AI-041980, and by the Department of Veterans Affairs. Dr. R. F. Schinazi is the principal founder of RFS Pharma, LLC. His laboratory received no funding for this work from the company and *vice versa*.

References and notes

1. Zeuzem S, Feinman SV, Rasenack J, Heathcote EJ, Lai MY, Gane E, O'Grady J, Reichen J, Diago M, Lin A, Hoffman J, Brunda M. *N Engl J Med*. 2000; 343:1666. [PubMed: 11106715]
2. (a) Carroll SS, Tomassini JE, Bossernan M, Getty K, Stahlhut MW, Eldrup AB, Bhat B, Hall D, Simcoe AL, Lafemina R, Rutkowski CA, Wolanski B, Yang Z, Migliaccio G, Francesco RD, Kuo LC, MacCoss M, Olsen DB. *J Biol Chem*. 2003; 278:11979. [PubMed: 12554735] (b) Clark JL, Hollecker L, Mason JC, Stuyver LJ, Tharnish PM, Lostia S, McBrayer TR, Schinazi RF, Watanabe KA, Otto MJ, Furman PA, Stec WJ, Patterson SE, Pankiewicz KW. *J Med Chem*. 2005; 48:5504. [PubMed: 16107149]
3. (a) Eldrup AB, Allerson CR, Bennett CF, Bera S, Bhat B, Bhat N, Bosserman MR, Brooks J, Burlein C, Carroll SS, Cook PD, Getty KL, MacCoss M, McMasters DR, Olsen DB, Prakash TP, Prhavc M, Song Q-L, Tomassini JE, Xia J. *J Med Chem*. 2004; 47:2283. [PubMed: 15084127] (b) Eldrup AB, Prhavc M, Brooks J, Bhat B, Prakash TP, Song QL, Bera S, Bhat N, Dande P, Cook PD, Bennett CF, Carroll SS, Ball RG, Bosserman M, Burlein C, Colwell LF, Fay JF, Flores OA, Getty K, LaFemina RL, Leone J, MacCoss M, McMasters DR, Tomassini JE, Langen DV, Wolanski B, Olsen DB. *J Med Chem*. 2004; 47:5284. [PubMed: 15456273]

4. Kumar R, Sharma N, Nath M, Saffran HA, Tyrrell DL. *J Med Chem.* 2001; 44:4225. [PubMed: 11708924]
5. (a) Harry-O'kuru RE, Smith JM, Wolfe MS. *J Org Chem.* 1997; 62:1754.(b) Tamerlani, G.; Salsini, L.; Lombardi, I.; Bartalucci, D.; Cipolletti, G. *US Pat Appl Publ* 2004158059. 2004. p. 12(c) Storer, R.; Moussa, A.; Chaudhuri, N.; Waligora, F. *PCT Int Appl WO* 2004052899. 2004. p. 90
6. Hocek M, Pohl R, Cisarova I. *Eur J Org Chem.* 2005; 14:3026.(b) Hocek M, Dvorakova H. *J Org Chem.* 2003; 68:5773. [PubMed: 12839482]
7. Spectral data for compound **6**: $^1\text{H-NMR}$ (CD_3OD , 400 MHz): δ 8.46 (s, 1 H), 5.98 (s, 1 H), 4.17 (d, $J=9.2$ Hz, 1 H), 4.00 (m, 1 H), 3.97 (d, $J=2.0$ Hz, 1 H), 3.81 (dd, $J=2.8$ Hz, $J=12.4$ Hz, 1 H), 2.56 (s, 3 H), 0.92 (s, 3 H). *LC/MS* (m/z), calcd for $\text{C}_{12}\text{H}_{18}\text{N}_5\text{O}_4$ ($\text{M}^+ + \text{H}$): 296.13; found, 296.19.
8. Uchiyama M, Aso Y, Noyori R, Hayakawa Y. *J Org Chem.* 1993; 58:373.
9. Bobeck DR, Schinazi RF, Coats SJ. *Antiviral Therapy.* 2010; 15:935. [PubMed: 21041908]
10. (a) Gillerman I, Fischer B. *J Med Chem.* 2011; 54:107. [PubMed: 21138280] (b) Šilhár P, Pohl R, Votruba I, Hocek M. *Org Lett.* 2004; 6:3225. [PubMed: 15355018] (c) Šilhár P, Pohl R, Votruba I, Hocek M. *Org Biomol Chem.* 2005; 3:3001. [PubMed: 16186932]
11. Qu G-R, Xia R, Yang X-N, Li J-G, Wang D-C, Guo H-M. *J Org Chem.* 2008; 73:2416. [PubMed: 18307357]
12. Spectral data for compound **9**: $^1\text{H-NMR}$ (CD_3OD , 400 MHz): δ 8.63 (s, 1 H), 6.04 (s, 1 H), 4.27 (m, 4 H), 4.18 (d, $J=9.2$ Hz, 1 H), 3.97 (m, 2 H), 3.81 (dd, $J=3.2$ Hz, $J=12.8$ Hz, 1 H), 1.33 (t, $J=7.2$ Hz, 6 H), 0.94 (s, 3 H). $^{13}\text{C-NMR}$ (CD_3OD , 400 MHz): δ 158.97, 152.61, 149.80, 147.58, 141.32, 89.56, 81.47, 77.54, 70.59, 62.46, 58.15, 17.45, 13.91. *LC/MS* (m/z), calcd for $\text{C}_{15}\text{H}_{25}\text{N}_5\text{O}_7\text{P}$ ($\text{M}^+ + \text{H}$), 418.14; found, 418.33.
13. Silverman LS, Caldwell JP, Greenlee WJ, Kiselgof E, Matasi JJ, Tulshian DB, Arik L, Foster C, Bertorelli R, Monopoli A, Ongini E. *Bioorg & Med Chem Lett.* 2007; 17:1659. [PubMed: 17234414]
14. Baraldi PG, Tabrizi MA, Gessi S, Borea PA. *Chem Rev.* 2008; 108:238. [PubMed: 18181659]
15. (a) Unciti-Broceta A, Infantas MJ, Gallo MA, Espinosa A. *Chem Eur J.* 2007; 13:1754. [PubMed: 17146826] (b) Kos NJ, Jongejan H, Van der Plas HC. *Gazz Chim Ital.* 1987; 117:369.
16. (a) Gupta M, Nair V. *Tetrahedron Lett.* 2005; 46:1165.(b) Kohgo, S.; Ohnui, H.; Kodama, E.; Matsuoka, M.; Mitsuya, H. *US Pat Appl Publ* 20090234110. 2009. p. 25
17. (a) Xu Y, Ikeda R, Sugiyama H. *J Am Chem Soc.* 2003; 125:13519. [PubMed: 14583048] (b) Schaeffer, HJ. *US Pat* 19804199574. 1980. p. 18
18. (a) Lioux T, Gosselin G, Mathé C. *Eur J Org Chem.* 2003:3997.(b) Sodem RS, Fiala ES. *Chem Res Toxicol.* 1998; 11:1453. [PubMed: 9860487] (c) Elzein E, Kalla R, Li X-f, Perry T, Marquart T, Micklatcher M, Li Y, Wu Y-Z, Zeng D, Zablocki J. *Bioorg Med Chem Lett.* 2007; 17:161. [PubMed: 17045477]
19. Spectral data for compound **21**: $^1\text{H-NMR}$ (CD_3OD , 400 MHz): δ 8.79 (d, $J=1.2$ Hz, 1 H), 8.59 (s, 1 H), 7.87 (d, $J=1.2$ Hz, 1 H), 6.16 (s, 1 H), 4.18 (d, $J=9.2$ Hz, 1 H), 4.00 (m, 2 H), 3.86 (dd, $J=3.2$ Hz, $J=12.8$ Hz, 1 H), 0.96 (s, 3 H). $^{13}\text{C-NMR}$ (CD_3OD , 400 MHz): δ 156.96, 149.80, 149.56, 140.32, 133.57, 123.63, 118.39, 91.87, 83.21, 79.03, 72.28, 59.85, 19.07. *LC/MS* (m/z), calcd for $\text{C}_{13}\text{H}_{17}\text{N}_8\text{O}_4$ ($\text{M}^+ + \text{H}$), 349.13; found, 349.16.
20. Cosyn L, Palaniappan KK, Kim SK, Duong HT, Gao ZG, Jacobson KA, Calenbergh SV. *J Med Chem.* 2006; 49:7373. [PubMed: 17149867]
21. Coats SJ, Link JS, Gauthier D, Hlasta D. *Org Lett.* 2005; 7:1469. [PubMed: 15816729]
22. Wanner MJ, KuÈnzle JK, Jzerman IAP, Koomen G-J. *Bioorg Med Chem Lett.* 2000; 10:2141. [PubMed: 10999489]
23. Wanner MJ, Koomen G-J. *J Chem Soc, Perkin Trans.* 2001; 11908
24. Ujjinamatada RK, Paulman RL, Ptak RG, Hosmane RS. *Bioorg Med Chem.* 2006; 14:6359. [PubMed: 16765050]
25. (a) Ujjinamatada RK, Phatak P, Burger AM, Hosmane RS. *J Med Chem.* 2008; 51:694. [PubMed: 18173230] (b) Berry DA, Wotring LL, Drach JC, Townsend LB. *Nucleosides Nucleotides.* 1994; 13:2001.

26. Rondla R, Coats SJ, McBrayer TR, Grier J, Johns M, Tharnish PM, Whitaker T, Zhou L-H, Schinazi RF. *Antivir Chem Chemother*. 2009; 20:99. [PubMed: 19843980]
27. Stuyver LJ, Whitaker T, McBrayer TR, Hernandez-Santiago BI, Lostia S, Tharnish PM, Ramesh M, Chu CK, Jordan R, Shi J, Rachakonda S, Watanabe KA, Otto MJ, Schinazi RF. *Antimicrob Agents Chemother*. 2003; 47:244. [PubMed: 12499198]
28. (a) Schinazi RF, Sommadossi JP, Saalman V, Cannon DL, Xie MW, Hart GC, Smith GA, Hahn EF. *Antimicrob Agents Chemother*. 1990; 34:1061. [PubMed: 2393266] (b) Stuyver LJ, Lostia S, Adams M, Mathew J, Pai BS, Grier J, Tharnish P, Choi Y, Chong Y, Choo H, Chu CK, Otto MJ, Schinazi RF. *Antimicrob Agents Chemother*. 2002; 46:3854. [PubMed: 12435688]
29. The assay was performed by combining 140 μ M of adenosine (final concentration) and 0.01 units of adenosine deaminase (Sigma catalog # A5043) in 50 mM potassium phosphate (pH 7.4). Compounds were tested at 10 μ M. Absorbance @ 265 nm was measured over four minutes and the slope was observed. Changes in the slope by incubation with compound *versus* no compound indicate inhibition as the slope approaches zero.
30. Sodum RS, Fiala SE. *Chem Res Toxicol*. 1998; 11:1453. [PubMed: 9860487]
31. (a) Koscsó B, Csóka B, Pacher P, Haskó G. *Expert Opin Investig Drugs*. 2011; 20:757.(b) Antoniolli L, Fornai M, Colucci R, Tuccori M, Blandizzi C. *Expert Opin Investig Drugs*. 2011; 20:717.

Figure 1.
Important 2'-C-methyl purine analogs.

Figure 2.
Side products of **12a** formation.

Phosphoramidate of **19**

Figure 3.
(2*S*)-ethyl(phenoxy)phosphorylamino)-propanoate prodrug.

Scheme 1.

Reagents and conditions: a) 2-amino-6-chloropurine, DBU, TMSOTf, CH₃CN, 80 °C, 5 h, 92%; b) Pd(Ph₃P)₄, Al(CH₃)₃, THF, 75 °C, 8 h, 54%; c) NH₃/CH₃OH, rt, 2 d, 62%; d) *t*-BuMgCl, phenyl ethoxyalaninyl phosphorochloridate, THF, rt, 18 h, 9 %.

Scheme 2.

Reagents and conditions: a) $\text{P}(\text{OEt})_3$, 130°C , 18 h; 78%; b) NH_3/EtOH , rt, 4 d, 44%; c) NaOEt/EtOH , rt, 2 d, then 50°C for 30 min, 19%.

Scheme 3.

Reagents and conditions: a) H₂NNHCOR, 110 °C, 2 d; b) BSA, 130 °C, 5 h, **12a**: 39%; **12b**: 72%; c) NaOMe/MeOH, rt, 18 h, **13a**: 50%; **13b**: 90%.

Scheme 4.

Reagents and conditions: a) NH₃/CH₃OH, THF, rt, 3 d, **17**: 46%, **18**: 21%; b) hydrazine, EtOH, rt, 16 h, 92%; c) NaNO₂, HOAc, rt, 1 h, 78%. e) sodium L-ascorbate, CuSO₄•5 H₂O, TMSCH, *t*-BuOH/H₂O (v/v, 1:1), rt, 18h, **21**: 12%, **22**: 30%.

Scheme 5.

Reagents and conditions: a) TBAN, TFAA, CH₂Cl₂, 85°C, 3 h, 79%; b) NaN₃, DMF, -18 °C, 1 h, then 0 °C, 2 h, 99%; c) H₂ (1 atm), Pd/C, EtOAc, EtOH, 45–50 °C, **26**: 1.5 h, 43%; **27**: 3 h, 41%.

Scheme 6.

Reagents and conditions: a) DBU, TMSOTf, CH₃CN, 0 °C, 30 min, 50 °C, overnight, **29**: 64%, **30**: 36%; b) *t*-butylamine, MeOH, rt, 2 days, for **31**: 91%, for **34**: 78%; c) i) hydrazine monohydrate, MeOH, 0 °C; ii) NaOEt (cat), EtOH, reflux, **32**: 71%, **33**: 5%, **35**: 62%, **36**: 12%.

Table 1

Anti-HCV activity, anti-HIV activity, and cytotoxicity of synthesized compounds in cellular assays.

Cmpd	Anti-HCV activity: % inhib @ 10 μ M in Huh7 cells			Anti-HIV activity, μ M			Cytotoxicity in: CC ₅₀ , μ M		
	HCV	rRNA	EC ₅₀	EC ₅₀	PBM	CEM	Vero		
6	0	0	70.1	>100	>100	12.9	>100		
7	0	0	68.2	>100	>100	>100	>100		
9	18.7	14.8	>100	>100	>100	>100	>100		
10	19.3	14.9	>100	>100	>100	>100	>100		
13a	0	0	>100	>100	>100	>100	>100		
13b	0	0	>100	>100	>100	>100	>100		
17	0	5.8	>100	>100	>100	>100	>100		
18	14.3	9.8	>100	>100	>100	>100	>100		
19	85.2	7.0	>100	>100	>100	>100	>100		
20	5.2	0	>100	>100	>100	>100	>100		
21	0	0	>100	>100	>100	>100	>100		
22	0	0	>100	>100	>100	>100	>100		
26	65.9	76.1	8.7	42.1	>100	27.4	41.5		
27	99.5	99.9	16.6	65.4	43.2	7.3	>100		
32	0	0	>100	>100	>100	>100	>100		
33	0	0	>100	>100	>100	>100	>100		
35	0	0	>100	>100	>100	>100	>100		
36	0	9.0	>100	>100	>100	>100	>100		