Fatal thrombocytopenia associated with intracardiac mass

Platelets are anuclear blood cells whose function is to form blood clots when we injure ourselves, to prevent excessive blood loss. Thrombocytopenia is a relative low number of platelets in the blood. Thrombocytopenia has several possible causes, including the trapping of platelets in spleen, reduced production of platelets and increased breakdown of platelets. Some examples of an increase in platelet breakdown include pregnancy, idiopathic thrombocytopenic purpura (ITP), thrombotic thrombocytopenic purpura (TTP), hemolytic uremic syndrome, infection, autoimmune diseases and medications (e.g., heparin). Thrombocytopenia may often be asymptomatic and can only be revealed by full blood cell count. Alternatively, it may cause excessive bruises and/or bleeding. In some cases, platelet count is so low that dangerous internal bleeding occurs.

In this case, the author reports a 22-year-old male presented with progressive breathlessness on exertion, increasing jaundice, easy bruising and petechial rash over a period of 15 days. The patient had no history or family history of common disorders that may cause these symptoms. Importantly, the patient had severe thrombocytopenia, revealed by peripheral blood smear evaluation, along with an elevation in direct bilirubin level, total leukocyte count and neutrophil count. Accordingly, several possible causes of thrombocytopenia were examined. For instance, bone marrow study failed to detect any possible abnormality in hematopoiesis, excluding the possibility that there was a decrease in platelet production. Also, abdominal ultrasounds showed normal-sized spleen, therefore, ruling out that platelets were trapped in the spleen. Further, color Doppler excluded the possibility of deep vein thrombosis in abdomen or lower limbs. Also, the patient had no recent exposure to heparin. Anti-phospholipid antibody, anti-system lupus erythematosus (SLE) antibodies and anti-PF4 antibodies were negative, so thrombocytopenia of this patient was not likely to be caused by antibody-induced

autoimmune responses. Both ITP and TTP were ruled out in this case as there was no evidence of end organ damage and micro-angiopathic hemolytic anemia, and the patient showed no response to steroids. Since there were normal lactate dehydrogenase and haptoglobin levels, hemolysis was also excluded. Anti-infection and anti-inflammatory medications as well as platelet transfusion failed to improve the symptoms of this patient. He died of cardiac arrest just 1 day after an episode of melena.

As most common causes of thrombocytopenia were excluded, what was the possible explanation of this fatal thrombocytopenia? The patient had facial puffiness and soft tender hepatomegaly, which is a common sign of right ventricular inflow obstruction. Indeed, cardiac auscultation found long mid-diastolic murmur in tricuspid area and harsh phasic ejection systolic murmur, grade III/VI, without ejection click at pulmonary area. Further evidence came from trans-thoracic echocardiography, which showed a homogenous mobile mass obstructing right ventricular inlet and outlet, similar to large right atrial myxoma. The finding was confirmed by contrast-enhanced computerized tomography of chest and abdomen. Therefore, the fatal thrombocytopenia appears to be associated with intracardiac mass in the right ventricle, which may resemble large right atrial myxoma.

Thrombocytopenia has been reported to be associated with cardiac tumors.^[1-4] Large cardiac tumors often result in intracardiac blood flow obstruction, eventually leading to right heart failure. Cardiac tumor-associated thrombocytopenia is often associated with other hematologic disorders such as anemia or erythrocytosis as well. The mechanism by which intracardiac tumor leads to thrombocytopenia remains unclear, although it has been postulated that abnormal mechanical shear stress, caused by tumor-induced flow obstruction, may be responsible for the increase in the breakdown of platelets.

In conclusion, the current study presented a rare case with fatal thrombocytopenia associated with intracardiac mass. It provides useful insights into the clinical complications of thrombocytopenia associated with intra-cardiac tumors as well as how to diagnose such conditions. This report will have important implications for helping other patients with similar thrombocytopenia associated with intra-cardiac tumors.

Access this article online	
Quick Response Code: 	Website: www.jcdronline.com
	DOI: 10.4103/0975-3583.95356

Kan Chen

*Center for Excellence in Vascular Biology,
Department of Pathology,
Brigham and Women's Hospital, Boston,
Massachusetts, USA*

Address for correspondence:

*Dr. Kan Chen,
Center for Excellence in Vascular Biology,
Department of Pathology,
Brigham and Women's Hospital,
New Research Building 750,
Boston, MA 02115, USA.
E-mail: kchen7@rics.bwh.harvard.edu*

REFERENCES

1. Kucharski W, Kosmala W, Silber M, Poreba R, Zych B, Zembala M. Thrombocytopenia and disseminated intravascular coagulation in a patient with left atrial myxoma: A case report. *Kardiol Pol* 2003;59:421-4.
2. Tsuda H, Imazeki N, Fuse Y, Maruyama T, Kitani A, Mizuno K. Cardiac angiosarcoma with gastrointestinal bleeding, hypoxemia, thrombocytopenia and microangiopathic hemolytic anemia. *Gan No Rinsho* 1986;32:1035-40.
3. Fishman AD, Hoffman A, Volterra F, Frymus M, Gentilucci M. Intracaval and intracardiac metastatic nonseminomatous germ cell tumor: A rare cause of hemolytic anemia and thrombocytopenia. *Cancer Invest* 2002;20:996-1001.
4. Semino A, Danova M, Perlini S, Palladini G, Riccardi A, Perfetti V. Unusual manifestations of disseminated neoplasia at presentation: Right-sided heart failure due to a massive cardiac metastasis and autoimmune thrombocytopenia in pleomorphic rhabdomyosarcoma of the adult, *Am J Clin Oncol* 2006;29:102-3.

How to cite this article: Chen K. Fatal thrombocytopenia associated with intracardiac mass. *J Cardiovasc Dis Res* 2012;3:65-6.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) First Page File:

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) Article File:

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1024 kb. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) Images:

Submit good quality color images. Each image should be less than **4096 kb (4 MB)** in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) Legends:

Legends for the figures/images should be included at the end of the article file.