CORRESPONDENCE

Treatment Strategies for Resistant Arterial Hypertension

by Dr. med. Felix Mahfoud, Dr. med. Frank Himmel, Dr. med. Christian Ukena, Prof. Dr. med. Heribert Schunkert, Prof. Dr. med. Michael Böhm, Prof. Dr. med. Joachim Weil in volume 43/2011

A Pseudo-Quality Assurance Fig Leaf

The review article draws impressive conclusions on renal denervation and also suggests that this might soon become almost standard procedure.

However, equally impressive is the frenzied speed with which this procedure is becoming widespread after only one controlled trial in Germany (1), and one with a limited number of patients, despite the fact that this trial on high blood pressure, a disorder strongly affected by psychological factors (white-coat hypertension), involved no placebo control, and that the current standard diagnostic procedure for hypertension was never followed: instead, normal office blood pressure readings were used. The authors' conclusion is therefore incorrect: at the current time a register would be a pseudo-quality assurance fig leaf. Much more beneficial would be further controlled trials with placebo treatments in a comparator population and appropriate standards for reading blood pressure, before this procedure is used in the huge population of up to 22% of cases of treatment-refractory hypertension (2). Unfortunately, we cannot put the genie back in the bottle.

DOI: 10.3238/arztebl.2012.0311a

REFERENCES

1. Esler MD, Krum H, Sobotka PA, Schlaich MP, Schmieder RE, Böhm M: Renal sympathetic denervation in patients with treatment-resistant hypertension (the Symplicity HTN-2 Trial): a randomised controlled trial. *Lancet* 2010; 376: 1903–9.
2. Lowel H, Meisinger C, Heier M, Hoyer H, Alte D, Volzke H: Epidemiology of hypertension in Germany. Selected results of population-representative cross-sectional studies. *Dtsch Med Wochenschr* 2006; 131: 2586–91.
3. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. *Dtsch Arztebl Int* 2011; 108(43): 725–31.

Dr. med. Akhil Chandra, Lüneburg, AChandra@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Anxiety as an Important Factor

In the discussion of the secondary causes of hypertension, one important factor was not mentioned: anxiety. Anxiety disorders often present physically, even without manifest anxiety, as tachycardia, shortness of

breath, cardialgia, dizziness, multifocal pain, high temperature, and arterial hypertension (1). In somatized anxiety attacks, systolic blood pressure can reach 300 mmHg, and diastolic blood pressure 120 mmHg.

The explanation for this is as follows: latent feelings of anxiety put the autonomic nervous system, muscle tone (pains), and the thyroid (hyperthyroidism) into “high gear.” Anxiety should therefore be considered in every case of resistant high blood pressure, particularly when other equivalents of anxiety such as nocturnal waking with sweating and craving for food are identified. The authors even lump together high blood pressure and anxiety under the heading of “white-coat hypertension.”

In these patients, blood pressure values usually fall significantly after discussion of their anxiety, of which they were not previously conscious, and its biographical origin. Anger or grief can also be associated with anxiety. The decisive factor is to put a name to the emotion (“Your high blood pressure is anxiety/anger/etc.”) and to understand it.

Only after diagnosis and treatment have addressed the emotional component of hypertension, and receptive methods have been exhausted, should the invasive procedure of renal denervation be considered.

DOI: 10.3238/arztebl.2012.0311b

REFERENCES

1. Küttemeyer M: Das chronische Müdigkeitssyndrom: eine Form der somatisierten Angstneurose. In: Berg PA (ed.) *Chronisches Müdigkeits- und Fibromyalgiesyndrom*. 2nd revised edition; Berlin: Springer 2003: 77–87.
2. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. *Dtsch Arztebl Int* 2011; 108(43): 725–31.

Dr. med. Mechthilde Küttemeyer

Fichtenstraße 52, 68535 Neckarhausen, kuete@arcor.de

Conflict of interest statement

The author declares that no conflict of interest exists.

An Experimental Procedure Paid for by the General Public

There are two aspects of renal nerve ablation that ought to be addressed. Renal nerve ablation is a fascinating method that was extremely successful in animal experiments as long as 30 years ago and may prove be beneficial to patients.

- Ablation could potentially be used for millions of patients in Germany alone. For such widespread use, a single controlled trial (Symplicity 2) involving 102 patients and lasting six months is no basis for general recommendations to German physicians. Symplicity 2 did not include a placebo group, and the endpoint, blood pressure, was not

verified by independent researchers. This unreliable method may explain why ablation led to a decrease in systolic office blood pressure of 32mmHg but only 11mmHg in 24-hour blood pressure. The publication on baroreceptor stimulation which was also discussed by the authors shows unambiguously how striking the effects of even placebo treatment can be on blood pressure in patients with treatment-refractory hypertension. Because of these shortcomings of the Symplicity 2 trial, the US authorities, unlike their German counterparts, have to our knowledge not approved renal nerve ablation and have demanded a better-controlled trial.

- It is regrettable that in our health-care system an experimental procedure is being developed at the expense of the general public rather than manufacturers. This is quite unlike the situation for pharmaceutical products, which are only approved after much more intensive testing. A voluntary register of uncontrolled data is no solution for determining the clinical value of ablation. What is needed for this purpose are appropriate randomized clinical trials of sufficient duration: a couple of months is insufficient for the indication of hypertension. The extent to which uncontrolled data can overestimate the efficacy of interventions used to treat hypertension is exemplified by the history of dilation of atherosclerotic renal arterial stenosis.

DOI: 10.3238/arztebl.2012.0311c

REFERENCES

1. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. *Dtsch Arztebl Int* 2011; 108(43): 725–31.

Prof. Dr. med. Johannes Mann

Städt. Klinikum München-Schwabing, prof.j.mann@googlemail.com

Prof. Dr. med. Karl Hilgers, Prof. Dr. med. Roland Veelken

Universitätsklinik Erlangen-Nürnberg

Prof. Dr. med. Friedrich Luft, Universitätsklinik Charité, Berlin

Conflict of interest statement

Prof. Hilgers has received fees for consultancy on the Advisory Board of Daiichi Sankyo. He has also received reimbursement for participation fees, travel expenses, and accommodation costs and lecture fees from Novartis.

Prof. Veelken has received reimbursement of participation fees for conferences and travel and accommodation costs from Novartis. He has received lecture fees from Novartis and Roche.

Prof. Mann and Prof. Luft declare that no conflict of interest exists.

obstructive sleep apnea can also have an aggravating/exacerbating effect. In some patients with androgen deficiency and sleep apnea, testosterone can be returned to normal levels by reestablishing the restorative REM sleep phase using CPAP. Primarily, however, lifestyle modifications are decisive in treatment and prevention (“First, don’t get fat”) and in individual cases growth hormone or testosterone therapy for a period limited to 3 to 6 months can be beneficial, almost as a kick-start (3, 4). Obesity-associated resting tachycardia indicates not only lack of fitness but also a burden on the cardiac atria/chambers and, under stress, long-term activation of the sympathetic nervous system. In such cases percutaneous renal denervation or baroreflex stimulation of the carotid sinus may be beneficial for selected patients, improving insulin sensitivity in a similar way to the treatment of patients with a catecholamine-secreting pheochromocytoma before and after the tumor is removed. In “typical” patients with endocrine hypertension, congenital adrenal hyperplasia (CAH) caused by 11-beta-hydroxylase or 17-alpha-hydroxylase deficiency should also be considered, although most patients with CAH present with 21-hydroxylase deficiency and may develop hypertension, or may only develop it, following over-replacement with dexamethasone or hydrocortisone (exogenous hypercortisolism/Cushing syndrome) and/or with predominant insulin resistance in connection with polycystic ovary syndrome.

DOI: 10.3238/arztebl.2012.0312a

REFERENCES

1. Koch CA, Wofford M, Ayala AR, Pacak K: Chapter 26 – Overview of Endocrine Hypertension. In: Chrousos GP, De Groot L: *Adrenal Physiology and Diseases*. www.endotext.org/adrenal/adrenal26/adrenal frame26.htm
2. Lanfranco F, Motta G, Minetto MA, et al.: Neuroendocrine alterations in obese patients with sleep apnea syndrome. *Int J Endocrinol*. 2010; 2010: 474–518.
3. Ullah MI, Washington T, Kazi M, Tamanna S, Koch CA: Testosterone deficiency as a risk factor for cardiovascular disease. *Horm Metab Res* 2011; 43(3): 153–64.
4. Widdowson WM, et al.: The effect of GH replacement on exercise capacity in patients with growth hormone deficiency: a meta-analysis. *J Clin Endocrinol Metab* 2008; 93: 4413–7.
5. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. *Dtsch Arztebl Int* 2011; 108(43): 725–31.

Prof. Dr. med. habil. Christian A. Koch, FACP, FACE

Mississippi, USA, ckoch@umc.edu

Conflict of interest statement

Prof. Koch has a personal connection with Merck. He has received fees for consultancy from Novo Nordisk. He has received fees for preparing scientific continuing education events from Ipsen. He has received fees for conducting clinical contract studies from Corcept Therapeutics.

Stress: Aspects of Endocrine Hypertension

An increase in endocrine hypertension is to be expected with a global increase in insulin-resistant obesity (few obese persons are insulin-sensitive!) (1). Some mechanisms and biomarkers for this are a deficiency of anabolic hormones (growth hormone and testosterone) and increased levels of leptin, the catabolic stress hormone cortisol, and resting heart rate (2). Concomitant

Biased Reasoning

The authors write, “Nevertheless, its clinical application is limited by the necessity for vascular surgery involving the carotid artery with the patient under

general anesthesia, the need for frequent replacement of the subcutaneously implanted device, and the risk of infection entailed by device implantation” and go on to rate this treatment as a purely experimental approach for a high-risk population. In contrast, renal denervation is valued as a procedure “with a low complication rate.” I find the unstated implications of this description somewhat biased. The short duration of follow-up in the Symplicity trial, involving only 17 of the original 138 patients after two years, is not discussed. In nephrology practice at least, the need to use a contrast substance for RND is a considerable obstacle in cases of stage 3 and above chronic kidney disease (CKD). On the other hand, the complication rates of the latest baroreflex studies (Bisognano J et al., Am Coll Cardiol 2011) are considerably lower than those of the initial study and will probably fall even further in the ongoing BAT study as a result of unilateral implantation (XR-1). It seems necessary to indicate the varying indication and effect profiles that are emerging in this regard. Baroreceptor stimulation probably achieves a greater drop in blood pressure but requires (minor) surgery and replacement, like cardiac pacemakers, which are becoming rarer as technology progresses. RND requires the use of contrast substance, which entails risk. It is faster and cheaper to perform, but its long-term effect requires better evaluation.

DOI: 10.3238/arztebl.2012.0312b

REFERENCES

1. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. Dtsch Arztebl Int 2011; 108(43): 725–31.

PD Dr. med. Joachim Beige, Abteilung Nephrologie und KfH Nierenzentrum am Klinikum St. Georg, Leipzig, Joachim.Beige@kfh-dialyse.de

Conflict of interest statement

Joachim Beige is participating as an investigator in the trial XR-1 to evaluate a new baroreflex stimulation system. This clinical trial involves payment of fees from CVRx for patient recruitment and trial conduct.

A New Procedure, But for Whom?

Before percutaneous renal denervation, an irreversible intervention, is held up to universal acclaim, further questions ought to be clarified. These include the patients in whom a lasting effect can be expected (are renin readings [1] expedient?) and the risks to kidney function (e.g. due to cholesterol embolization). *Figure 2* suggests, with a decreasing number of cases, that for two years after surgery there is an increasing effect in reducing blood pressure when compared to the subgroups used in the original publication (2). eGFR (estimated glomerular filtration rate) was recorded in only a few patients and fell significantly over two years. The procedure is not evaluated, or apparently not suitable, for patients with advanced renal failure, those requiring dialysis, or kidney transplant recipients, in whom resistance to treatment is common. For these patients, information on existing treatment options would be valuable. The table „Drug treatment....“ is

confusing: no superiority of aliskiren over other renin-angiotensin system inhibitors in reducing blood pressure is evidence-based, alpha-1-receptor-blockers and alpha-blockers are identical, doxazosin is not included, and endothelin antagonists are not approved for the treatment of arterial hypertension. Also omitted is the outstanding importance of salt-water balance (3) in resistance to treatment, by taking into consideration diet, duration of drug effects, and patients' renal function. Low doses of aldosterone antagonists are administered for primary hypertension in addition to another diuretic. Of all the above-mentioned substances, the greatest and most reliable antihypertensive effect is that of minoxidil, although minoxidil has multiple side effects. With these caveats in mind, it cannot be disputed that renal denervation may extend the range of antihypertensive tools available. However, pharmacological options should be fully explored before intervention. The trials at hand also show that antihypertensive medication needs to be taken even after renal denervation.

DOI: 10.3238/arztebl.2012.0313a

REFERENCES

1. Brown MJ: Personalised medicine for hypertension. Measuring plasma renin could refine the treatment of resistant hypertension. BMJ 2011; 343: d4697.
2. Symplicity HTN-1 Investigators: Catheter-Based Renal Sympathetic Denervation for Resistant Hypertension Durability of Blood Pressure Reduction Out to 24 Months. Hypertension 2011; 57: 911–7.
3. Appel LJ: Another major role for dietary sodium reduction: improving blood pressure control in patients with resistant hypertension. Hypertension 2009; 54: 444–6.
4. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. Dtsch Arztebl Int 2011; 108(43): 725–31.

Prof. Dr. med. Manfred Anlauf

Bremerhaven/Cuxhaven, Manfred.Anlauf@t-online.de

Prof. Dr. med. Franz Weber, St. Walburga-Krankenhaus, Meschede

Conflict of interest statement

Prof. Anlauf holds shares in Astella Pharma and Novartis. Prof. Weber has received fees for preparing scientific continuing education events from Daiichi Sankyo and Fresenius.

In Reply:

Our article provides an overview of the treatment for resistant hypertension, a problem rarely addressed in research. We therefore consider the information provided by our colleagues a welcome addition. Reversible causes should be identified before any pharmacological or interventional therapy is provided (see *Figure 1*). *Table 1* of the publication contains an overview of the available pharmacological treatment strategies. Substances have been selected according to current guidelines (1). Evaluation of blood pressure-reducing properties within the groups was deliberately omitted. Aldosterone antagonist therapy is particularly important, because its efficacy in reducing blood pressure in patients with resistant hypertension has

been demonstrated in studies (2). Minoxidil causes extremely high levels of side effects, which leads to poor compliance and so considerably restricts widespread clinical use. More detailed discussions of drugs are available and were not the subject of this article. For reasons of space it was only possible to examine common secondary causes of hypertension.

Particular emphasis was placed on interventional renal sympathetic denervation. This procedure is a new, promising treatment option for patients with resistant hypertension for whom pharmacological treatment options have been exhausted. It is important to state again that to date there are no clinically proven pharmacological or non-pharmacological treatment procedures for these patients with high cardiovascular risk. Our article clearly indicates that renal denervation is currently not a standard procedure, but rather an alternative option for the treatment of hypertension in cases of confirmed resistance to treatment. We therefore strongly recommend that our critical correspondents reread our publication.

As mentioned in the article, 206 patients, 106 as part of the randomized Symplicity HTN2 trial, have so far undergone systematic research in clinical trials, and follow-up for a period of 24 months (3, 4). The effects of renal denervation over longer follow-up times (36 months) in the Symplicity HTN1 trial show a sustained drop in blood pressure and do not provide any evidence of renal arterial stenosis or aneurysms (manuscript submitted for publication). In the USA the first patients have been enrolled in the Symplicity HTN3 trial in which randomization is performed after renovasography in a catheterization laboratory and placebo treatment is started. Due to intraprocedural pain, the extent to which blinding and placebo treatment is really possible remains to be seen, as do the ethical and legal considerations arising from this. It is more than unlikely to ascribe a drop in blood pressure over 24 to 36 months to a placebo effect.

Large, randomized trials have shown clearly that reductions in blood pressure over 24 hours during antihypertensive treatment are less marked than office blood pressure values. It is therefore not surprising that this is also the case following renal denervation. The analysis of the effects of renal denervation on 24-hour blood pressure will provide important information on the effect of renal denervation on day and night blood pressure values (manuscript submitted). The international registry, allowing scientific analysis of up to 5000 patients following renal denervation, will provide a valuable contribution to the evaluation of long-term effects and safety and represents a significant addition to randomized trials. In other, ongoing clinical trials, patients with advanced and terminal renal failure, chronic heart failure, milder forms of hypertension, and other cardiovascular diseases with high sympathetic activity are also undergoing systematic research. In the authors' opinion, endpoint trials are needed for these indications, because unlike hypertension no confirmed surrogate parameters for clinical endpoints exist. For

now the procedure should remain restricted to patients following the criteria used in the Symplicity trials (see Box 3).

Baroreflex stimulation is a pathophysiologically interesting concept for the treatment of resistant hypertension. However, it requires long-term pacemaker implantation, which currently requires vascular surgery under general anesthesia. Further development involving unilateral carotid stimulation and therefore longer battery life is currently undergoing clinical testing. It remains to be seen whether this surgery can regularly be performed under local anesthesia and so also reduce the complication rate associated with the procedure.

DOI: 10.3238/arztebl.2012.0313b

REFERENCES

1. Mancia G, De Backer G, Dominiczak A, et al.: 2007 Guidelines for the management of arterial hypertension: The task force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2007; 25: 1105–87.
2. Vlachy J, Sedlak R, Plachy M, et al.: Addition of spironolactone in patients with resistant arterial hypertension (ASPIRANT): a randomized, double-blind, placebo-controlled trial. *Hypertension* 2011; 57: 1069–7.
3. The Symplicity HTN-1 Investigators: Catheter-based renal sympathetic denervation for resistant hypertension: durability of blood pressure reduction out to 24 months. *Hypertension* 2011; 57: 911–7.
4. Esler MD, Krum H, Sobotka PA, et al.: Renal sympathetic denervation in patients with treatment-resistant hypertension (The Symplicity HTN-2 Trial): a randomised controlled trial. *Lancet* 2010; 376: 1903–9.
5. Mahfoud F, Himmel F, Ukena C, Schunkert H, Böhm M, Weil J: Treatment strategies for resistant arterial hypertension. *Dtsch Arztebl Int* 2011; 108(43): 725–31.

Dr. med. Felix Mahfoud, Dr. med. Christian Ukena

Prof. Dr. med. Michael Böhm, Klinik für Innere Medizin III, Kardiologie, Angiologie und Internistische Intensivmedizin, Universitätsklinikum des Saarlandes, Homburg/Saar, felix.mahfoud@uks.eu

Dr. med. Frank Himmel, Prof. Dr. med. Heribert Schunkert

Prof. Dr. med. Joachim Weil, Medizinische Klinik II, Universitätsklinikum Schleswig-Holstein, Campus Lübeck

Conflict of interest statement

The authors of the article received research funding from Ardian/Medtronic Inc., Palo Alto, USA to finance the conduct of clinical trials.

Dr. Mahfoud, Dr. Ukena, and Prof. Böhm are supported by the Ministry for Economics and Science of the Saarland and by the German Research Foundation (DFG; Clinical Research Group KFO 196).

Dr. Mahfoud is supported by the German Hypertension League (Deutsche Hochdruckliga) and has received fees for lectures and consultancy from Berlin-Chemie, Boehringer Ingelheim, Medtronic, Novartis, and Takeda Pharm.

Dr. Himmel is supported by Medtronic within the scope of the AF grant program and has received fees for lectures and consultancy from Medtronic and St. Jude Medical.

Dr. Ukena has received fees for lectures and consultancy from Boehringer Ingelheim and Medtronic.

Prof. Schunkert has received fees for lectures and consultancy from Astra Zeneca, Bayer, Boehringer Ingelheim, Daiichi-Sankyo, Novartis, Pfizer, Sanofi-Aventis, Berlin-Chemie, MSD, Takeda, and Servier.

Prof. Böhm has received fees for lectures and consultancy from Astra Zeneca, Boehringer Ingelheim, Novartis, Sanofi-Aventis, Servier, Medtronic, and Pfizer.

Prof. Weil has received fees for lectures and consultancy from Actelion, Astra Zeneca, Bayer, Daiichi-Sankyo, Novartis, Medtronic, and Pfizer.