Technical Section

TECHNICAL NOTES AND TIPS

EDITORIAL

Differences in technique between different disciplines: a lesson in sharing evidence

Bruce Campbell

Editor, Technical Section

Consultant Vascular Surgeon¹ and Professor²

¹Royal Devon and Exeter NHS Foundation Trust

and ²Peninsula Medical School**

Exeter EX2 5DW

John S McGrath

Consultant Urologist

Exeter Surgical Health Services Research Unit

Royal Devon and Exeter NHS Foundation Trust

Exeter EX2 5DW

The technical note by Mownah *et al* on ultrasound mapping of the long saphenous vein for coronary artery bypass¹ prompts reflection on how different disciplines might profit from observing and reading about the practices of others. It is just one illustration of surgeons in two related specialties approaching the same task in different ways, despite evidence and practice that might guide them otherwise.

The note describes how to map and mark the long saphenous vein using ultrasound prior to harvesting, with the aim of making incisions accurately over it, so reducing the amount of dissection and subsequent wound complications. The tone of the article and the fact that our reviewer judged it to have merit suggest that this is a practice that is unfamiliar to many cardiac surgeons (or at least not in common use by them). For peripheral vascular surgeons, saphenous vein mapping and marking by ultrasound has been normal practice for many years and is well described in their specialist literature.^{2,3} It seems strange that cross-fertilisation has not occurred between vascular and cardiac surgery in this area.

A further, closely related, illustration of different practice relates to the way the long saphenous vein is harvested. Since the 1980s it has been common practice among vascular surgeons to expose the vein through a number of short incisions, based on evidence that wound problems occur less frequently and discharge from hospital is less often delayed.^{4,5} By contrast, most cardiac surgeons continued to harvest the vein using a continuous long incision. The complications from these long incisions stimulated their interest in endoscopic harvesting, which may have inferior long-term outcomes for the grafts.⁶

None of this is intended as a vascular surgical snipe at cardiac surgeons: the technical note on vein mapping by ultrasound was simply a prompt to raise this general issue. Other illustrations of different approaches are not difficult to find: variation in perioperative management for bowel resection provides a good example. Within the last decade, there have been numerous meta-analyses (including Cochrane reviews) demonstrating that routine placement of nasogastric tubes and routine use of mechanical bowel prepara-

tion are unnecessary in elective colorectal surgery.^{7–10} Studies have demonstrated that pre-operative mechanical bowel preparation and delayed feeding after operation are not only unnecessary but can be counterproductive.^{8–10}

On the basis of this evidence, colorectal surgeons largely abandoned these practices, but similar changes have been rather slow to disseminate through other disciplines involved in abdominal surgery. Many urologists continue to advocate pre-operative 'elemental diets', mechanical bowel preparation and routine nasogastric tube placement in procedures such as radical cystectomy. It is only recently that studies have appeared in the urological literature showing that omission of these measures prior to radical cystectomy is safe and associated with an accelerated recovery.^{11,12}

Working within specialty-based 'silos' has impeded the rapid adoption of advances in surgical care. This culture has also bred a perceived but in some cases unnecessary need to repeat clinical trials by specialty-specific groups to confirm findings that should logically apply across similar patient groups. The examples above should perhaps make surgeons of all specialties think about the techniques used and published by related disciplines, from whom they often remain strangely separate. A certain tribalism persists in surgery and medicine, which fuels the belief that the practices of one's own specialty are just a little better than all the others. There may be useful lessons to learn from the technical literature of other disciplines.

References

1. Mownah OA, Middleton PR, Nair RU. Ultrasound mapping of long saphenous vein surface anatomy in coronary artery bypass surgery. *Ann R Coll Surg Engl* 2011; **93**: 415.
2. Salles-Cunha SX, Andros G. Preoperative duplex scanning prior to infrainguinal revascularization. *Surg Clin North Am* 1990; **70**: 41–59.
3. Magee TR, Leopold PW, Campbell WB. Vein marking through ultrasound coupling gel. *Eur J Vasc Surg* 1990; **4**: 491–492.
4. Schwartz ME, Harrington EB, Schanzer H. Wound complications after in situ bypass. *J Vasc Surg* 1988; **7**: 802–807.
5. Wengrovitz M, Atnip RG, Gifford RR *et al*. Wound complications of autogenous subcutaneous infrainguinal arterial bypass surgery: predisposing factors and management. *J Vasc Surg* 1990; **11**: 161–163.
6. Zenati MA, Shroyer AL, Collins JF *et al*. Impact of endoscopic versus open saphenous vein harvest technique on late coronary artery bypass grafting patient outcomes in the ROOBY (Randomized On/Off Bypass) Trial. *J Thorac Cardiovasc Surg* 2011; **141**: 338–344.
7. Slim K, Vicaud E, Launay-Savary MV *et al*. Updated systematic review and meta-analysis of randomized clinical trials on the role of mechanical bowel preparation before colorectal surgery. *Ann Surg* 2009; **249**: 203–209.
8. Guenaga KK, Matos D, Wille-Jørgensen P. Mechanical bowel preparation for elective colorectal surgery. *Cochrane Database Syst Rev* 2009; **21**: CD001544.
9. Gravante G, Caruso R, Andreani SM, Giordano P. Mechanical bowel preparation for colorectal surgery: a meta-analysis on abdominal and systemic complications on almost 5,000 patients. *Int J Colorectal Dis* 2008; **23**: 1,145–1,150.
10. Pineda CE, Shelton AA, Hernandez-Boussard T *et al*. Mechanical bowel preparation in intestinal surgery: a meta-analysis and review of the literature. *J Gastrointest Surg* 2008; **12**: 2,037–2,044.
11. Ruthi RS, Nielsen M, Smith A *et al*. Fast track program in patients undergoing radical cystectomy: results in 362 consecutive patients. *J Am Coll Surg* 2010; **210**: 93–99.
12. Arumainayagam N, McGrath J, Jefferson KP, Gillatt DA. Introduction of an enhanced recovery protocol for radical cystectomy. *BJU Int* 2008; **101**: 698–701.