Published in final edited form as:

Angew Chem Int Ed Engl. 2011 August 16; 50(34): 7906–7909. doi:10.1002/anie.201102404.

Modular Monodentate Oxaphospholane (OxaPhos) Ligands: Utility in Highly Efficient and Enantioselective 1,4-Diboration of 1,3-Dienes

Christopher H. Schuster, Dr. Bo Li, and Prof. Dr. James P. Morken

Department of Chemistry, Boston College, Chestnut Hill, MA 02467 (USA), Fax: (617)-552-2705

James P. Morken: morken@bc.edu

Keywords

Asymmetric catalysis; Ligand design; Boron; Oxidation

Chiral monodentate ligands are important tools for the control of enantioselectivity in catalytic reactions. These compounds are particularly important when transition metal catalysts lack sufficient coordination sites to bind a bidentate ligand and still retain activity. Along these lines, modular, tunable chiral carbenes^[1] and phosphoramidites^[2] have found particular prominence in the field of asymmetric catalysis. Aside from these compound classes, however, there is a relative paucity of effective, tunable monodentate chiral ligands that have found widespread utility. In recent studies on platinum-catalyzed enantioselective diene diboration, we achieved limited success when optimizing chiral phosphoramidite and phosphonite ligands for certain substrates.^[3,4] Because catalytic diboration benefits from the presence of electron-rich monodentate phosphines^[5], we considered that solutions to problematic substrates might arise from the availability of a readily preparable, tunable, chiral, Lewis basic phosphine ligand.^[6] In this manuscript, we describe the synthesis and properties of enantiomerically enriched modular 1,3-oxaphospholanes (termed OxaPhos ligands, Scheme 1). These ligands are readily available from the combination of an enantiopure epoxide, a primary phosphine, and a ketone or the derived ketal.^[7] In addition to describing their synthesis and properties, we also describe their use in the catalytic 1,4-diboration of challenging diene substrates.

The OxaPhos ligands possess two stereocenters, one at carbon and one at phosphorous, and are therefore available in two epimeric forms. The preparation of both epimers of the Et-OxaPhos ligand is described in Scheme 2 and is representative of the synthesis of this ligand class. Enantiomerically enriched terminal epoxide **1** was treated with phenylphosphide to furnish ring-opened secondary phosphine **2**. Without purification, phosphine **2** was subjected to transketalization with the dimethyl ketal of the reacting ketone. This synthesis sequence delivered a 1:1 mixture of tertiary phosphines that were directly oxidized with hydrogen peroxide. The stable phosphine oxides were easily separated by silica gel chromatography and subsequent stereoretentive reduction with phenylsilane^[8] cleanly delivered the isomerically pure tertiary phosphines in good yield.

The crystal structures of the phosphine oxide derivatives of both *cis*- and *trans*-Et-OxaPhos ligands are depicted in Figure 1.^[9] A notable feature of these structures is the invariant nature of the five-membered ring with the *tert*-butyl substituent adopting an equatorial

position in each. Assuming similar conformations predominate in the reduced phosphine derivatives, one can anticipate that the phosphorus lone pair in the *cis*-OxaPhos ligand is better aligned with the adjacent C-O σ^* orbital (OPCO dihedral angle = 151°) compared to the *trans*-OxaPhos ligand (OPCO dihedral = 99°). However, this alignment appears to have little consequence on the ligand electronic properties; after reduction, the derived *trans*- $\text{L}_2\text{Rh}(\text{CO})\text{Cl}$ complexes exhibit nearly identical CO stretching frequencies (1961 cm^{-1} for the *cis* ligand and 1962 cm^{-1} for the *trans*).^[10] Of note, the CO stretching frequencies also suggest that the phosphorous atom in the OxaPhos ligands is relatively basic, comparable to PhPCy_2 (CO stretching frequency for *trans*-(PhPCy_2) $_2\text{Rh}(\text{CO})\text{Cl}$ = 1964 cm^{-1}).^[10b]

To study the utility of oxaphospholanes in catalytic diboration, diene **3** was chosen for analysis (Table 1). Previous studies from our laboratory documented the efficacy of chiral TADDOL-derived phosphonite ligands in diene diboration reactions and with the most optimal ligand (xylyl-TADDOL derived phenyl phosphonite) this substrate under went diboration in only 70% ee.^[3] As shown in Table 1 and discussed below, diboration under the influence of oxaphospholane ligands can be much more selective, and reactivity is outstanding. While diborations with propylene oxide derived ligands and ligands derived from aldehydes were not highly selective (data not shown), when the *tert*-butyl substituted ligands were employed, elevated levels of enantiomeric enrichment were obtained. As depicted in Table 1, the *cis* and *trans* isomers of the OxaPhos ligands favor opposite product enantiomers. Significantly, the enantioselectivity with the *cis* ligand isomer is markedly enhanced as the size of the ketal substituents is increased, (substituents larger than *iso*-butyl could not be incorporated with the current synthesis route) whereas the opposite trend occurs with the *trans* epimer. Importantly, with the *cis*-*t*Bu-OxaPhos ligand, the selectivity with *trans*-1,3-pentadiene (Table 1) far surpasses that obtained with (xylyl)TADDOL-derived phosphonite ligand (97:3 er versus 85:15 er).

To survey the utility of *cis*-*t*Bu-OxaPhos in the diboration of other dienes, the collection of substrates in Table 2 was examined. As shown by the data, *cis*-*t*Bu-OxaPhos is useful for a range of aromatic and aliphatic terminal dienes, affording the derived 1,4-diol in excellent yield and enantioselectivity upon oxidative work-up. The selective diboration reaction could be extended to 3,4-disubstituted dienes as depicted by entries 5–7. The diboration of these substrates occurs in low selectivity with TADDOL-derived ligands, but with *cis*-*t*Bu-OxaPhos, enantiomeric purities of 96–98 % ee were observed. A particularly noteworthy feature is that these substrates deliver difficult-to-access trisubstituted alkene products in a highly diastereo- and enantioselective fashion. The result in entry 8 suggests that the diboration strategy with *cis*-*t*Bu-OxaPhos can deliver chiral, tetrasubstituted alkene products with synthetically useful levels of selectivity. Lastly, for entry 4 it was demonstrated that the intermediate 1,4-bis(boronate) could be isolated in excellent yield and fully characterized.

The *cis*-buten-1,4-diols obtained by diene diboration can be readily oxidized^[11] to butenolides and therefore represent valuable synthetic intermediates.^[12] However, to access these structures on large scale using Pt catalysis, the amount of catalyst employed in the reaction can be a concern. To address this issue, the effect of catalyst composition and loading were examined. Similar to the reactions described above, these reactions were carried out with rigorously dried and deoxygenated solvents and were conducted under argon. However, in contrast to the studies above, the substrate for these experiments was also rigorously deoxygenated. As indicated in entry 1 (Table 3), <5% reaction occurred when a 2:1 L:M ratio was employed. Suspecting that PtL_2 complexes might be inactive and that trace oxygen in the substrate might serve to oxidize a portion of the ligand and provide an active ML_1 complex, the experiment was repeated (entry 2), but after 4 hours (no reaction, TLC analysis) the mixture was briefly exposed (< 5 sec) to air, sealed, and allowed to react for 2 more hours. This experiment provided diol **6** in excellent yield and

enantiomeric purity. Also consistent with the hypothesis that the active species may be an ML_1 complex, is that with 1.95:1 L:M and degassed substrate (entry 3), the reaction proceeds normally. Notably, L:M ratios of 1.1:1 and 0.5:1 (entries 4 and 5) provide reasonable yields and selectivity in diboration of **5**, with the latter experiment indicating that a significant level of ligand accelerated catalysis occurs. Expecting that at lower catalyst loading trace oxygen may be more problematic, an L:M ratio of 2:1 was employed as the amount of catalyst used in experiments was decreased to 0.1 mol% and 0.02 mol% (entries 6–8). These reactions used non-degassed substrate and, as depicted in entry 8, even at a 0.02 mol% loading of Pt(0), the diene diboration proceeds in excellent yields and enantioselection. This level of reactivity is in marked contrast to diborations with TADDOL-derived phosphonite ligands which proceed in <5% conversion at 0.02 mol% loading and in 23 % conversion at the 0.1 % loading level.

The remarkable diboration activity of Pt catalysts derived from *cis*-*i*Bu-OxaPhos raises the concern that the presence of trace amounts of stereoisomeric catalysts (i.e. from incomplete chromatographic resolution during synthesis) might erode selectivity. To study this feature, catalysts of varying diastereomeric purity were examined in the diboration of 1,3-pentadiene. As depicted in Figure 2, these experiments revealed a substantial non-linear effect that favors the more selective *cis* diastereomer of the *i*-Bu-OxaPhos ligand.^[13] Remarkably, even a 1:1 mixture of *cis* and *trans* *i*-Bu-OxaPhos ligands delivers synthetically useful levels of selectivity. Furthermore, the more selective *cis* ligand, even when contaminated with 25% of the less selective *trans* isomer, provides product in nearly indistinguishable levels of selectivity as that derived from ligand that is >97% pure *cis* isomer. Thus, not only is trace contamination by the *trans* stereoisomer of catalyst of little consequence, it is also not necessarily important to achieve substantial resolution during the synthesis outlined in Scheme 1.

In conclusion, OxaPhos ligands are tunable, chiral, monodentate phosphines that can offer very high enantioselectivity in catalytic diboration. Of particular note is that at the 0.02 mol % level of catalyst loading with which these catalysts can be effective, the cost of the catalyst becomes meaningless relative to the substrates and reagents in the diboration reaction.^[14] Considering the impact that phosphoramidite ligands have had on asymmetric catalysis and the observation that OxaPhos ligands can offer advantages, it is anticipated that the latter ligand class may find significant use in catalysis.

Acknowledgments

We are grateful to the NIH (NIGMS GM-59417) for support of this work and to AllyChem for a gift of B₂(pin)₂

References

1. For recent reviews of chiral carbenes: Douthwaite RE. *Coord Chem Rev.* 2007; 251:702. Gade LH, Bellemin-Laponnaz S. *Top Organomet Chem.* 2007; 21:117. Cesar V, Bellemin-Laponnaz S, Gade LH. *Chem Soc Rev.* 2004; 33:619. [PubMed: 15592627] For recent examples of monodentate chiral carbenes, see: Park JK, Lackey HH, Rexford MD, Kovnir K, Shatruck M, McQuade DT. *Org Lett.* 2010; 12:5008. [PubMed: 20919706] Lee KS, Hoveyda AH. *J Org Chem.* 2009; 74:4455. [PubMed: 19445467] Vehlouw K, Wang D, Buchmeiser MR, Blechert S. *Angew Chem.* 2008; 120:2655. *Angew Chem Int Ed.* 2008; 47:2615. Fournier PA, Collins SK. *Organometallics.* 2007; 26:2945.
2. For reviews of chiral phosphoramidites and related compounds, see: Teichert JF, Feringa BL. *Angew Chem.* 2010; 122:2538. *Angew Chem Int Ed.* 2010; 49:2486. Feringa BL. *Acc Chem Res.* 2000; 33:346. [PubMed: 10891052] Selected examples: van den Berg M, Minnard AJ, Schudde EP, van Esch J, de Vries AHM, de Vries JG, Feringa BL. *J Am Chem Soc.* 2000; 122:11539. Jensen JF, Svendsen BY, la Cour TV, Pedersen HL, Johannsen M. *J Am Chem Soc.* 2002; 124:4558.

- [PubMed: 11971692] Ma MFP, Li K, Zhou Z, Tang C, Chan ASC. *Tetrahedron: Asymm.* 1999; 10:3259. Yu RT, Rovis T. *J Am Chem Soc.* 2006; 128:12370. [PubMed: 16984159] Bee C, Han SB, Hassan A, Iida H, Krische MJ. *J Am Chem Soc.* 2008; 130:2746. [PubMed: 18266373] Shirakura M, Sugimoto M. *Angew Chem.* 2010; 122:3915. *Angew Chem Int Ed.* 2010; 49:3827. Teller H, Flügge S, Goddard R, Fuerstner A. *Angew Chem.* 2010; 122:1993. *Angew Chem Int Ed.* 2010; 49:1949.
3. Burks HE, Kliman LT, Morken JP. *J Am Chem Soc.* 2009; 131:9134. [PubMed: 19505078]
 4. Non-enantioselective diene diboration: Ishiyama T, Yamamoto M, Miyaura N. *Chem Commun.* 1996:2073. Clegg W, Thorsten J, Marder TB, Norman NC, Orpen AG, Peakman TM, Quayle MJ, Rice CR, Scott AJ. *J Chem Soc, Dalton Trans.* 1998:1431. Ely RJ, Morken JP. *Org Lett.* 2010; 12:4348. [PubMed: 20828186]
 5. For the accelerating effect of basic phosphines on diboration, see: Alkynes: Thomas RL, Souza FE, Marder TB. *J Chem Soc, Dalton Trans.* 2001:1650. Allenes: Ishiyama T, Kitano T, Miyaura N. *Tetrahedron Lett.* 1998; 39:2357. Burks HE, Liu S, Morken JP. *J Am Chem Soc.* 2007; 129:8766. [PubMed: 17589992] Pelz NF, Woodward AR, Burks HE, Sieber JD, Morken JP. *J Am Chem Soc.* 2004; 126:16328. [PubMed: 15600327]
 6. Reviews involving chiral monodentate phosphines, see: Erre G, Enthaler S, Junge K, Gladiali S, Beller M. *Coord Chem Rev.* 2008; 252:471. Grabulosa A, Granell J, Muller G. *Coord Chem Rev.* 2007; 251:25. Crepy KVL, Imamoto T. *Top Curr Chem.* 2003; 229:1. Komarov IV, Borner A. *Angew Chem.* 2001; 113:1237. *Angew Chem Int Ed.* 2001; 40:1197. Lagasse F, Kagan HB. *Chem Pharm Bull.* 2000; 48:315. [PubMed: 10726849] Hayashi T. *Accs Chem Res.* 2000; 33:354. Pietrusiewicz KM, Zablocka M. *Chem Rev.* 1994; 94:1375. For select recent examples, see: Alberico E, Gladiali S, Taras R, Junge K, Beller M. *Tet Asymm.* 2010; 21:1406. Yamamoto T, Yamada T, Nagata Y, Sugimoto M. *J Am Chem Soc.* 2010; 132:7899. [PubMed: 20481636] Birkholz MN, Dubrouina NV, Shuklov IA, Holz J, Paciello R, Walosh C, Breit B, Borner A. *Tet Asymm.* 2007; 18:2055. Zhang A, Rajanbabu TV. *Org Lett.* 2004; 6:1515. [PubMed: 15101781] Colby EA, Jamison TF. *J Org Chem.* 2003; 68:156. [PubMed: 12515474]
 7. For the synthesis of related achiral or racemic oxaphospholanes, see: Oehme H, Issleib K, Leissring E. *Tetrahedron.* 1972; 28:2587. Marsi KL, Co-Sarno ME. *J Org Chem.* 1977; 42:778. Lopez-Cortina S, Basiulis DI, Marsi KL, Munoz-Hernandez MA, Ordonez M, Fernandez-Zertuche M. *J Org Chem.* 2005; 70:7473. [PubMed: 16149773] For a related chiral, bidentate bis(phosphine), see Tang W, Qu B, Capacci AG, Rodriguez S, Wei X, Haddad N, Narayanan B, Ma S, Grinberg N, Yee NK, Krishnamurthy D, Senanayake CH. *Org Lett.* 2010; 12:176. [PubMed: 19950894]
 8. Marsi KL. *J Org Chem.* 1974; 39:265.
 9. Crystal structures have been deposited with the Cambridge Structure Data Center: CCDC 820558–820561
 10. For the use of CO stretch to measure basicity, see: Vastag S, Heil B, Markó L. *J Mol Cat.* 1979; 5:189. Otto S, Roodt A. *Inorg Chim Acta.* 2004; 357:1. Molloy KG, Petersen JL. *J Am Chem Soc.* 1995; 117:7696. Clarke ML, Cole-Hamilton DJ, Slawin AMZ, Woollins JD. *Chem Commun.* 2000:2065.
 11. (a) Butina D, Sondheimer F. *Synthesis.* 1980:543. (b) Chakraborty TK, Chandrasekaran S. *Tetrahedron Lett.* 1984; 25:2891. (c) Minami I, Tsuji J. *Tetrahedron.* 1987; 43:3903. (d) Mehta G, Roy S. *Chem Commun.* 2005:3210. (e) Yang Y-K, Choi J-H, Tae J. *J Org Chem.* 2005; 70:6995. [PubMed: 16095335]
 12. Seitz M, Reiser O. *Curr Opinion in Chem Biol.* 2005; 9:285.
 13. Considering that PtL₂ complexes appear to be inactive with PtL₁ complexes responsible for catalysis, the origin of the non-linear effect may arise from a higher population of monoligated Pt-(*R,R*)-*i*-BuOxaPhos relative to the diastereomeric PtL₁ complex, or it may be that the Pt-(*R,R*)-*i*-BuOxaPhos is more reactive than the Pt-(*S,R*)-*i*-BuOxaPhos complex. These features will be the subject of a future study.
 14. Pt(dba)₃ is available from K₂PtCl₄ in one step (85% yield). At a price of ca. \$20k/mol for K₂PtCl₄, a one mole scale diene diboration at 0.02 mol % Pt requires ca. \$4 worth of catalyst. The current price of B₂(pin)₂ from AllyChemUSA, Inc. is ca. \$200/mol.

Figure 1.
ORTEP representation of *cis* and *trans* Et-OxaPhos phosphine oxides.

Figure 2.
Correlation between catalyst diastereomer ratio and product.

Scheme 1.
Modular assembly of OxaPhos ligands.

Scheme 2.

Synthesis scheme for assembly of Et-OxaPhos ligands.

Table 1Pt-Catalyzed Enantioselective Diboration/Oxidation of *trans*-Piperylene in the Presence of OxaPhos Ligands.

[a]

ligand	yield [%] ^[b]	er ^[c]
<i>cis</i> -Me-OxaPhos	83	85.5:14.5
<i>trans</i> -Me-OxaPhos	62	20.2:79.8
<i>cis</i> -Et-OxaPhos	85	94.5:5.5
<i>trans</i> -Et-OxaPhos	86	22.8:77.2
<i>cis</i>-<i>i</i>Bu-OxaPhos	86	97.0:3.0
<i>trans</i> - <i>i</i> Bu-OxaPhos	88	38.9:61.1

[a] Reactions employ 1.05 equivalents of B₂(pin)₂ and were carried out at 60 °C for 12 h.

[b] Isolated yield of purified material. Value is an average of two experiments.

[c] Determined by chiral GC. Error ±0.2%.

Table 2

Pt-Catalyzed Enantioselective Diboration/Oxidation in the Presence of (*R,R*)-*i*Bu-OxaPhos.^[a]

6 mol % (*R,R*)-*i*Bu-OxaPhos
3 mol% Pt(dba)₃
B₂(pin)₂, toluene
60 °C, 12 h
then NaOH, H₂O₂

(*R,R*)-*i*Bu-OxaPhos

entry	substrate	product	% yield ^[b]	er ^[c]
1			86	97.0:3.0
2			65	96.1:3.9
3			>95	98.2:1.8
4			>95(>95)	97.5:2.5
5			82	99.0:1.0
6			>95	98.8:1.2
7			61 ^[e]	98.1:1.9
8			82	82.5:17.5

^[a]Reactions employ 1.05 equivalents of B₂(pin)₂ and were carried out at 60 °C for 12 h.^[b]Isolated yield of purified material. Value is an average of two experiments.^[c]Determined by chiral GC, HPLC, or SFC analysis. Error ±0.2%. With (xylyl)TADDOL-PPh, the enantioselectivities of entries 1–5 are 70, 84, 91, 84, and 45 % ee, respectively.^[d]Yield in parentheses is that for the purified 1,4-bis(boronate) intermediate.^[e]2:1 ratio of 1,4:1,2 diboration products obtained. Yield is for the purified 1,4 product.

Table 3

Effect of catalyst loading and composition.^[a]

entry	% ligand	% Pt(dba) ₃	yield [%] ^[b]	ton ^[c]	ee ^[d]
1 ^[e]	2.0	1.0	<5%	-	-
2 ^[f]	2.0	1.0	90	90	96.4:3.6
3 ^[e]	1.95	1.0	97	97	97.4:2.6
4 ^[e]	1.1	1.0	98	98	97.3:2.7
5 ^[e]	0.5	1.0	71	71	95.4:4.6
6	0.2	0.1	98	980	97.3:2.7
7	0.1	0.05	95	1900	97.2:2.8
8	0.04	0.02	95	4750	97.1:2.9

^[a] Reactions under Ar atmosphere with dry deoxygenated solvent, employing 1.05 equivalents of B₂(pin)₂, and carried out at 60 °C for 12 h.^[b] Isolated yield of purified material.^[c] Turnover number = % yield/mol % Pt.^[d] Determined by chiral GC. Error ±0.2%.^[e] Reaction employed degassed substrate.^[f] Reaction employed degassed substrate and briefly exposed to air (<5 sec) at 4 hours.