Published in final edited form as:

J Agric Food Chem. 2012 April 11; 60(14): 3541–3550. doi:10.1021/jf205296s.

Dehydropyrrolizidine Alkaloids, Including Monoesters with an Unusual Esterifying Acid, from Cultivated *Crotalaria juncea* (Sunn Hemp cv. 'Tropic Sun')

Steven M. Colegate^{*,†}, Dale R. Gardner[†], Robert J. Joy[‡], Joseph M. Betz[§], and Kip E. Panter[†]

[†]Poisonous Plant Research Laboratory, Agriculture Research Service, US Department of Agriculture, 1150 East 1400 North, Logan, Utah 84341, USA

[‡]Natural Resources Conservation Service, Plant Materials Center, US Department of Agriculture, P.O. Box 236, Hoolehua, Hawaii 96729, USA

[§]Office of Dietary Supplements, National Institutes of Health, 6100 Executive Blvd., Room 3B01, Bethesda, Maryland 20892, USA

Abstract

Cultivation of *Crotalaria juncea* L. (Sunn Hemp cv. 'Tropic Sun') is recommended as a green manure crop in a rotation cycle to improve soil condition, help control erosion, suppress weeds and reduce soil nematodes. Because *C. juncea* belongs to a genus that is known for the production of toxic dehydropyrrolizidine alkaloids, extracts of the roots, stems, leaves and seeds of 'Tropic Sun' were analyzed for their presence using HPLC-ESI/MS. Qualitative analysis identified previously unknown alkaloids as major components along with the expected macrocyclic dehydropyrrolizidine alkaloid diesters, junceine and trichodesmine. The dehydropyrrolizidine alkaloids occurred mainly as the *N*-oxides in the roots, stems and, to a lesser extent, leaves, but mainly as the free bases in the seeds. Comprehensive spectrometric and spectroscopic analysis enabled elucidation of the unknown alkaloids as diastereoisomers of isohemijunceine, a monoester of retronecine with an unusual necic acid. The dehydropyrrolizidine alkaloid content of the roots, stems, leaves of immature plants was estimated to be 0.05, 0.12 and 0.01% w/w respectively while seeds were estimated to contain 0.15% w/w.

Keywords

Crotalaria juncea; 'Tropic Sun'; pyrrolizidine alkaloids; *N*-oxides; HPLC-ESI/MS; junceine; trichodesmine; isohemijunceine; hemijunceine

INTRODUCTION

Crotalaria juncea L. (sunn hemp) is an extensively cultivated, leguminous plant with a multitude of potential uses/benefits, including non-wood fiber and soil improvement, considered to have originated in India.¹ Agronomic development of the plant in the USA resulted in the 1983 release of the 'Tropic Sun' cultivar that is recommended as a green

*Corresponding Author Tel: +1-435-752-2941. Fax: +1-435-753-5681., steven.colegate@ars.usda.gov.

The authors declare no competing financial interests

Supporting Information

A pdf containing jpg copies of all NMR spectra described herein.

manure crop cultivated in a rotation cycle to improve soil condition, help control erosion, suppress weeds and reduce soil nematodes.^{2, 3}

The dehydropyrrolizidine alkaloid macrocyclic diesters junceine (**1**) (Figure 1), trichodesmine (**2**), senecionine, riddelliine and seneciphylline have been reported to occur in *C. juncea* seeds.⁴ However, another batch of seeds, extracted by the same authors under the same conditions, only yielded **1**. In a study of the total pyrrolizidine alkaloid content and relative toxicities of seeds of various *Crotalaria* species to chicks, Williams and Molyneux determined the total pyrrolizidine alkaloid concentration to be 0.02% based on dry weight of the seeds of *C. juncea* sourced from India.⁵ The low levels and the variation in dehydropyrrolizidine alkaloid content of these earlier reports were later confirmed during the analysis of seeds from several accessions of *C. juncea* collected from various international locations.⁶ This latter report included an analysis of seeds from the 'Tropic Sun' cultivar that only revealed the presence of junceine using HPLC coupled to an evaporative light scattering detector (ca. 1.6 μ moles/g seed or 0.06% w/w).

The 'Tropic Sun' cultivar has undergone 28 years of development since its release in 1983. For this reason, along with increasing attention being given to the potential for long-term or intermittent low-level exposure to dietary dehydropyrrolizidine alkaloids to cause chronic diseases in humans, including cirrhosis, genotoxicity, cancers and pulmonary hypertension,⁷ the 'Tropic Sun' cultivar was re-examined for the presence, type and quantity of potentially toxic dehydropyrrolizidine alkaloids. In contrast to the previous reports on the alkaloid content in seeds, this study examined the roots, stems and leaves in addition to the seeds using an HPLC-electrospray ionization mass spectrometry (ESI/MS) approach.

MATERIALS AND METHODS

Chemicals and Reagents

Acetonitrile was HPLC certified solvent (Honeywell Burdick and Jackson, Muskegon, MI, USA). Water for the HPLC mobile phase was Milli-Q-purified (18.2 M Ω /cm) (Millipore, USA). Methanol was reagent ACS/USP/NF grade (Pharmaco Products, Brookfield, CT, USA). The mobile phase additives; ammonium acetate, formic acid, acetic acid and trifluoroacetic acid were each the highest grade available. Junceine (**1**), trichodesmine (**2**), supinine (**3**), amabiline (**4**) and lasiocarpine (**5**), 7-angelylheliotridine (**6**) and 9-angelylheliotridine (**7**) (Figure 1) were sourced from the stocks of extracted and purified (ca. 80–95% pure based on HPLC-ESI/MS analysis) pyrrolizidine alkaloids kept by the USDA/ARS Poisonous Plant Research Laboratory. Lycopsamine (**8**) was purchased from Planta Analytica (Danbury, CT, USA). The archived seed extract of *C. juncea*, sourced from India and previously analyzed using HPLC,⁵ was supplied by Dr R. J. Molyneux (University of Hawaii at Hilo). Unless otherwise stated, all solid phase extraction cartridges and HPLC columns and guard cartridges were sourced from Phenomenex (Torrance, CA, USA).

Extraction of Plant Samples

The plant samples were supplied by the Natural Resources Conservation Service, Plant Materials Center, US Department of Agriculture from cultivated, seed production fields in Hawaii in April 2011. Composite samples of roots, stems and leaves were each derived from about 10 immature (90 days from planting) plants. The seeds, that are harvested approximately 150 days after planting, were derived from a general seed bank. The air-dried samples were milled to a fine powder.

For the quantitative comparison, an aliquot (ca. 400 – 500 mg) of each composite sample was extracted 5 times with methanol (10 mL) at room temperature and with gentle, inversion

mixing for 16 h. Each successive extract was monitored for dehydropyrrolizidine alkaloid content using HPLC-ESI/MS and then combined to yield the four total extracts i.e., roots, stems, leaves and seeds, which were each concentrated to almost dryness *in vacuo*. The resultant residues were extracted with 0.05 M sulfuric acid (ca. 10 mL) to provide the alkaloid (*N*-oxides as well as free base alkaloids) solution for further cation exchange enrichment.

For the structural elucidation studies, milled seed (160 g) was Soxhlet-extracted with methanol (1.2 L) for 3 periods of 16 h each. Fresh methanol was used for each extraction period to lessen or avoid any degradation of dehydropyrrolizidine alkaloids following extended periods in the boiling methanol. Again, the yield of dehydropyrrolizidine alkaloids after each extraction was monitored using HPLC-ESI/MS to ensure complete extraction. The combined methanol extracts were evaporated to almost dryness and re-extracted with 0.05 M sulfuric acid (3 × 50 mL) to yield the aqueous acid solution of the alkaloids for further purification.

Strong Cation Exchange, Solid Phase Extraction (SCX SPE)

The filtered 0.05 M sulfuric acid-soluble fraction from each of the concentrated methanolic extracts was applied to a separate stack of two conditioned (according to the manufacturer's guidelines) 500 mg/3 mL, Strata SCX 55 μ m 70 Å SCX SPE columns. After loading, the stacked columns were separated and each was washed with water, methanol and, finally, with ammoniated methanol (a 10% solution of saturated ammoniated methanol in methanol) to elute the pyrrolizidine alkaloids and their *N*-oxides. The ammoniated methanol fractions were immediately evaporated to dryness under a stream of nitrogen and the residues reconstituted in methanol (1 mL). Efficiency of alkaloid capture and elution was monitored using HPLC-ESI/MS. The preparative extract of seeds was processed in a similar way except on a more appropriate larger scale 25 mm × 100 mm cation exchange column using Septra SCX, 50 μ m, 65 Å bulk media.

HPLC-ESI/MS

Chromatographic separation of the dehydropyrrolizidine alkaloids and their *N*-oxides (Figure 2) was achieved using a Finnigan Surveyor HPLC system, comprising an Autosampler Plus and a MS Pump Plus (Thermo-Finnigan, San Jose, CA, USA), equipped with a 150 mm × 2 mm i.d., 4 μ m, Synergi Hydro Reversed Phase column, with a 4 mm × 2 mm i.d. AQ C18 guard column. Sample injections (2 μ L) were eluted with a gradient flow (400 μ L/min) in which the initial mobile phase composition of 5% acetonitrile in aqueous 0.05% trifluoroacetic acid/0.5% acetic acid was held for 2 min and then linearly increased to 50% acetonitrile over 13 min. This was held for a further 10 min before returning to the initial mobile phase composition and re-equilibration of the column. The column effluent was monitored using a LCQ Advantage ion trap mass spectrometer (Thermo-Finnigan, San Jose, CA, USA) in the electrospray ionization (ESI), positive ion mode. The mass spectrometer response was tuned to a solution of heliotrine (acquired from stocks held by the USDA/ARS Poisonous Plant Research Laboratory) in methanol. The capillary temperature was 275 °C with a capillary voltage of 46 V. The source voltage was 4.5 kV at a source current of 80 μ A. The sheath and auxiliary gas flows were set at a ratio of 60:10. Data-dependent MS/MS spectra were acquired in the second scan of a two scan sequence in which ions identified in the first total ion scan were isolated and fragmented using 32% applied dissociation energy.

Qualitative Analysis

An even numbered m/z for the MH^+ ion followed by observation of MS/MS fragment ions characteristic of dehydropyrrolizidine alkaloids eg., m/z 94, 120, 122, 136, 137, 138, 156, 180 highlighted HPLC-ESI/MS peaks as probable dehydropyrrolizidine alkaloids.^{8,9}

Observation of a significant dimer ion $[2M + H]^+$ indicated potential dehydropyrrolizidine-*N*-oxide character under these esiMS conditions.⁸ *In situ* treatment, of an analytical sample indicative of *N*-oxides, with indigocarmine-based redox resin reduced the *N*-oxides to their parent dehydropyrrolizidine alkaloids thereby confirming the *N*-oxide character.⁸

Quantitative Analysis

Standard solutions containing lycopsamine (**8**) (69, 34.5, 17.25, 8.63, 4.31 and 2.15 $\mu\text{g/mL}$) and junceine (**1**) (200, 100, 50, 25, 12.5, 6.25 $\mu\text{g/mL}$) were made by adding a solution of **8** (200 μL of 138 $\mu\text{g/mL}$ methanol) to a solution of **1** (200 μL of 400 $\mu\text{g/mL}$ methanol) and serially diluting (1:1) an aliquot (200 μL) in methanol. Calibration standard samples were then prepared by adding lasiocarpine (**5**) solution (10 μL ca. 50 $\mu\text{g/mL}$ methanol), to provide a normalizing injection standard (IS), to an aliquot (100 μL) of each of the standard solutions.

The analytical samples were similarly prepared by adding IS (10 μL) to an aliquot (100 μL) of the redox resin-reduced extract solution. Peak areas, generated from reconstructed ion chromatograms displaying the ions of interest, were adjusted within each run by division by the area of the IS for that run.

Preparative Flash Chromatography

The 10% ammoniated m ethanol eluate from the SCX SPE of seed extract was evaporated to dryness to afford a red gum (5 g) that was re-extracted with chloroform ($3 \times 50 \text{ mL}$). The pooled, orange chloroform extract was filtered, reduced in volume (to about 15 mL) and applied to a Biotage KP silica Samplet (10 g) cartridge of silica gel (Biotage, North America). A narrow orange band was retained on the Samplet cartridge. The loaded Samplet cartridge was fitted to a Biotage Snap KP-silica gel column (50 g) that had been equilibrated with ethyl acetate using the Isolera 1 automated flash chromatography system (Biotage, North America). Elution of retentates from the column assembly was achieved at a flow rate of 50 mL/min, collecting 20 mL fractions, using a linear gradient of 10% ammoniated methanol (0 – 40% over 5 min) into ethyl acetate, hold for 6 min to elute non-dehydropyrrolizidine alkaloid material, then increasing from 40 – 100% ammoniated methanol over 7 min, to elute the dehydropyrrolizidine alkaloids, for a total run time of 18 min. The eluate was simultaneously monitored at 254 and 280 nm that provided an indication, albeit poor due to low UV absorbance, of the alkaloid elution times. The collected fractions were then analyzed for dehydropyrrolizidine alkaloid content using HPLC-ESI/MS.

Preparative and Semi-preparative HPLC

Alkaloid-enriched samples derived using the Biotage flash chromatography system were evaporated to dryness and reconstituted in 0.1% trifluoroacetic acid in water. Aliquots (up to 400 μL) of this solution were applied to a 100 mm $\times$ 21.2 mm i.d., 4 μ , 80 Å, AXIA-packed Synergi Hydro Reversed Phase column with a 15 mm $\times$ 21.2 mm i.d. C18 guard column. Using a Waters Prep LC 2000 System (Waters Corporation, Milford, MA, USA), the alkaloids were eluted with an isocratic (0.1% trifluoroacetic acid/acetonitrile: 93/7) flow (20 mL/min) and collected into bulk fractions determined by monitoring the absorbance at 220

nm. The dehydropyrrolizidine alkaloid content of the fractions was then determined using HPLC-ESI/MS.

Using a Finnigan Surveyor HPLC system, comprising a LC Pump Plus and a UV-VIS Plus (Thermo-Finnigan, San Jose, CA, USA), two isocratic, semi-preparative HPLC modes were used to achieve final separation of target analytes in the dehydropyrrolizidine alkaloid concentrates (up to 200 μ L injections) derived from the preparative HPLC. The column effluent was monitored at 220 nm. Basic separation conditions were achieved using a 250 mm $\times$ 10 mm i.d., 5 μ m, 110 Å Gemini NX C18 column with a 10 mm $\times$ 10 mm i.d. guard column of the same material with acetonitrile (7%) into 20 mM ammonium acetate at pH 9.5 (NH_4OH) (93%) at 5 mL/min. Acidic separation conditions were achieved using a 250 mm $\times$ 10 mm i.d., 4 μ m, 80 Å Synergi Hydro Reversed Phase column with a 10 mm $\times$ 10 mm i.d. Synergi AQ C18 guard column with acetonitrile (7%) into 0.1% trifluoroacetic acid in water (93%) at 5 mL/min.

Structure Elucidation

Low resolution MS and MS/MS data were acquired in the HPLC-ESI/MS mode described in a previous section. The high resolution mass spectrometry (HRMS) measurement was achieved using an Agilent 6220 TOF mass spectrometer in the dual ESI, positive ion mode. One dimensional and 2D (COSY, HSQC and HMBC) ^1H (300 MHz) and ^{13}C (75 MHz) NMR data were acquired using a JEOL Eclipse NMR spectrometer using solutions in CDCl_3 and the residual proton in the chloroform as the lock signal (except for junceine for which d_6 -DMSO was used as the solvent and the crude seed extract which was dissolved in d_4 -methanol).

Junceine (1) (peak 1, Figure 2) and Trichodesmine (2) (peak 6, Figure 2)—The HPLC-ESI/MS data (Table 1) were the same as for the authenticated standards and the ^1H and ^{13}C NMR data were consistent with previous reports.^{12–14}

Isohemijunceine A (9) (peak 3, Figure 2)—Obtained as a colorless oil estimated at > 95% pure using HPLC-ESI/MS and ^1H NMR; ESI(+)/MS and MS/MS (Table 1); HRMS (found, 283.17842; calcd. for $\text{C}_{15}\text{H}_{25}\text{NO}_4$ 283.1784; Δ 0.07 ppm); ^1H , ^{13}C , COSY and HMBC NMR (Table 2).

Isohemijunceine B (10) (peak 5, Figure 2)—Obtained as a colorless oil estimated at 85 – 90% pure using HPLC-ESI/MS and ^1H NMR; ESI(+)/MS and MS/MS (Table 1); HRMS (found, 283.17892; calcd. for $\text{C}_{15}\text{H}_{25}\text{NO}_4$ 283.1784; Δ 1.8 ppm); ^1H , ^{13}C and COSY NMR (Table 3).

Isohemijunceine C (11) (peak 7, Figure 2)—Obtained as a colorless oil estimated at 85 – 90% pure using HPLC-ESI/MS and ^1H NMR; ESI(+)/MS and MS/MS (Table 1); HRMS (found, 283.17912; calcd. for $\text{C}_{15}\text{H}_{25}\text{NO}_4$ 283.1784; Δ 2.5 ppm); ^1H , ^{13}C , COSY and HMBC NMR (Table 4).

Combined NMR and HPLC-ESI/MS Analysis of Seed

Seeds (50 g) were milled to a fine powder. Three sub-samples were dried in an oven at 40 °C for 5 days to determine the moisture content of about 5%. A larger sample (40.99 g) was Soxhlet-extracted with methanol (400 mL) for 20 h. The extraction solvent was removed and the extraction continued twice more with fresh solvent (20 h and 48 h) to lessen any degradation that might result from prolonged boiling in methanol. The combined extracts were evaporated to dryness *in vacuo* to afford a yellow solid (5.7 g, 14%) that was subsequently extracted with 0.05M H_2SO_4 . The aqueous acid solubles were applied to a

conditioned 5g/20mL Strata SCX 55 μm 70 Å SCX SPE column that discharged, under gravity, onto a second conditioned 5g/20mL Strata SCX 55 μm 70 Å SCX cartridge. After complete addition of the acid extracts, the two cartridges were separately washed with methanol (causing elution of an orange colour) and then with 10% satd. ammoniated methanol to elute the alkaloids.

The combined ammoniated methanol eluates were evaporated to dryness to yield a dark red oil (0.127 g) that was reconstituted in d_4 -methanol (700 μL) and the ^1H NMR spectrum recorded. Then an aliquot of *p*-dinitrobenzene (25 μL of 23 mg/mL d_4 -methanol) was added and the spectrum re-acquired. A further aliquot (50 μL) of the *p*-dinitrobenzene was added and the spectrum again re-acquired to provide an additional check on quantitation. The C2 proton signals in both NMR spectra were integrated relative to the *p*-dinitrobenzene singlet, both as a total envelope (ca 6.2 – 5.8 ppm) or as two separate envelopes, about 6.2 ppm for the macrocyclic diesters or about 5.8 ppm for the monoesters, and the amount of dehydropyrrolizidine alkaloids estimated according to Molyneux *et al.*¹⁰

The same sample, suitably diluted with methanol, was then assessed using the HPLC-ESI/MS method using trichodesmine (**2**) (isolated from *C. juncea* in this study; 38.1, 19.05, 9.53, 4.77, 2.38, 1.19 and 0.59 $\mu\text{g/mL}$) and lycopsamine (**8**) (34.5, 17.25, 8.63, 4.32, 2.16, 1.08 and 0.54 $\mu\text{g/mL}$) as the calibration standards for the macrocyclic diesters and monoesters respectively.

RESULTS AND DISCUSSION

Qualitative Analysis of Roots, Stems, Leaves and Seeds

HPLC-ESI/MS analysis (Figure 2) and the MS/MS data (Table 1) showed a differential distribution of free base dehydropyrrolizidine alkaloids and their *N*-oxides within the plant tissues. The seeds contained the free bases as the major pyrrolizidine alkaloid components (> 95%) whereas the roots and stems contained the *N*-oxides with only traces of the free base dehydropyrrolizidine alkaloids. About equal, but relatively low levels of the free base alkaloids and their *N*-oxides were extracted from the leaves. These observations are consistent with previous findings that the dehydropyrrolizidine alkaloids are biosynthesized in roots as their water-soluble *N*-oxides and then transported to the reproductive tissues where they accumulate as the more lipophilic free bases.¹¹

Detection of Previously Identified Alkaloids—Consistent with previous reports^{4, 6} the presence of the macrocyclic diester alkaloids **1** and **2** was indicated by the peaks in reconstructed ion chromatograms showing a protonated molecule (MH^+) at m/z 370 (peak 1, Figure 2) and m/z 354 (peak 6, Figure 2) respectively or by their *N*-oxides showing MH^+ and $[\text{2M} + \text{H}]^+$ ions at m/z 386/771 (peak 4) and m/z 370/739 (peak 12) respectively. Following isolation of the putative junceine and trichodesmine from seed extract, using preparative and semi-preparative reversed phase HPLC, their identities were confirmed by comparison of their ^1H and ^{13}C NMR data with literature reports.^{12–14} Based upon HPLC-ESI/MS/MS, only a minor amount of the previously reported⁴ senecionine (MH^+ m/z 336, peak 16, Figure 2) was observed in these analyses. Contrary to this previous report⁴ there were no HPLC-ESI/MS indications of the related, macrocyclic diester dehydropyrrolizidine alkaloids riddelliine or seneciophylline.

Detection and Identification of Previously Undescribed Alkaloids—At the commencement of this study, the analytical HPLC mobile phase conditions were the same as previously reported⁸ using 0.1% formic acid in a gradient with acetonitrile. The MS/MS fragmentation ions at m/z 156, 138, 120 and 94 observed under these conditions for two unidentified major peaks in the seed extracts, each displaying a MH^+ at m/z 284, were

indicative of dehydropyrrolizidine alkaloid character. However, the MS/MS data were inconsistent with the isobaric (MH^+ m/z 284) diastereoisomers of the known dehydropyrrolizidine alkaloids that lack oxygenation at C7 such as supinine (**3**) and amabiline (**4**) that show an abundant fragmentation ion at m/z 122 under the same conditions. The *N*-oxides of these putative dehydropyrrolizidine alkaloids, with $MH^+/[2M + H]^+$ ions at m/z 300/599 and confirmed by *in situ* reduction back to the parent free base alkaloids, were also observed, especially in the stems and roots.

However, preliminary NMR spectroscopy observations for the second of these peaks, in particular the unusual multiplicity of the C9 proton resonances and the doubling up of most signals in the proton-decoupled ^{13}C NMR spectrum, indicated that it was a mixture of very closely structurally-related compounds. Subsequent HPLC-ESI/MS using trifluoroacetic acid as the acidic modifier rather than formic acid showed the partial resolution of the two free base components of this second peak in the seed extract (peaks 5 and 7, Figure 2) and their respective *N*-oxides in the stem extract (peaks 11 and 13).

Preparative reversed phase HPLC under trifluoroacetic acid-based mobile phase conditions provided (>95% pure) peak 3 (Figure 2), the first, and best separated, of the three observed m/z 284 peaks, and a fraction containing peaks 5, 6 and 7 (Figure 2). Subsequent semi-preparative reversed phase HPLC using the basic mobile phase conditions with the Gemini NX column separated trichodesmine (peak 6, Figure 2) from peaks 5 and 7 that co-elute from this column under these basic conditions. Finally, semi-preparative HPLC using the Synergi Hydro reversed phase column and trifluoroacetic acid-based mobile phase conditions allowed isolation (ca. 85 – 90% pure) of peaks 5 and 7.

Structure Elucidation of Peak 3, Peak 5 and Peak 7 Alkaloids: These alkaloids had identical MS (MH^+ m/z 284) and very similar MS/MS fragmentation profiles (Table 1). High resolution mass measurements supported a molecular formula $C_{15}H_{25}NO_4$ for each. The MS/MS data (Table 1) showed moderately abundant ions at m/z 120 and 156, and a base ion peak at m/z 138 that together support a necine C9/C7 monoester alcohol structure.¹⁵ The fragmentation of m/z 284 to 138 therefore indicates a $C_7H_{14}O_3$ necic acid group that is unusual in that it is at least one oxygen atom less than C7 necic acids previously reported for pyrrolizidine alkaloid monoesters.^{9, 16} The loss of 44 Da from the protonated molecule (m/z 284 $\rightarrow$ 240) under LC-MS conditions is characteristic of a terminal hydroxy(methyl)methine (CH_3CHOH) group on the esterifying acid^{8, 17} analogous to the loss of 45 Da due to the same terminal group under electron ionization MS.¹⁸ Therefore, to be consistent with the putative $C_7H_{14}O_3$ necic acid group, the MS/MS data indicate a monoester with 3-hydroxy-2-*isopropyl*butanoic acid rather than the 2,3-dihydroxy-2-*isopropyl*butanoic acid more usually associated with the lycopsamine (**8**) type of dehydropyrrolizidine alkaloids.¹⁹

Analysis of the 1D and 2D NMR spectroscopic data for the peak 3 alkaloid (Table 2), the peak 5 alkaloid (Table 3) and the peak 7 alkaloid (Table 4) confirmed their common, two dimensional structure as the C9 ester of 1,2-dehydro-7-hydroxypyrrolizidine with 3-hydroxy-2-*isopropyl*butanoic acid. Since this ester forms half of the junceine (**1**) macrocycle, and since it is on the C9 rather than the C7 as in **1**, the peak 3, peak 5 and peak 7 alkaloids are given the trivial names isohemijunceine A (**9**), isohemijunceine B (**10**) and isohemijunceine C (**11**) respectively.

The determination of C9 esterification and the configuration around C7 and C8 was in accord with a review of proton NMR data by Logie *et al.*,²⁰ and a ^{13}C NMR study by Jones *et al.*²¹ Thus, the relatively lower field resonance of the two C9 protons i.e., between δ 4.83 – 4.92 and 4.51 – 4.67 indicated C9 esterification in contrast to expected chemical shifts of

approximately δ 3.6 – 4.3 for C7 esterification. This was supported by the HMBCs (Tables 2 and 4) between the carbonyl C11 and the protons on C9. The hydroxylation of C7, indicated by the MS/MS data (fragment ion at m/z 138 requires either C7 or C9 to be hydroxylated), was confirmed by the downfield chemical shifts of H7 at δ 4.29 – 4.31 and C7 at δ 71.04 – 71.27. The retronecine configuration of the necine base i.e., H8 and H7 both α relative to the plane of the ring, was indicated by the chemical shifts of C1 (δ 132.98 – 133.38), C2 (δ 129.14 – 130.43), C6 (δ 35.97 – 36.08) and C7 (δ 71.04 – 71.27) that are upfield, downfield, downfield and upfield respectively from those expected of a heliotridine-based alkaloid.²¹

The C12 protonation postulated based on the MS/MS data, in contrast to the more usual C12 hydroxylation,¹⁹ was confirmed by NMR experiments i.e., HSQC; HMBCs to C11, C13, C14, C15, C16 and C17; and the COSY interactions with H13 and H15.

While the chromatographic properties and NMR data for isohemijunceines B (**10**) and C (**11**) are very similar, indicating minor diastereoisomeric changes, the better chromatographic separation and the significant differences in the NMR spectra for isohemijunceine A (**9**) indicate a more substantial diastereoisomeric change. Previous reports describing saturated²² and 1,2-unsaturated¹⁸ pyrrolizidine alkaloid C9 monoesters of the lycopsamine (**8**) type i.e., esterified with viridifloric or trachelanthic acids that are 2,3-dihydroxy-2-*isopropyl*butanoic acids diastereoisomeric about C3, have differentiated the stereochemistry of the necic acid on the basis of the difference in chemical shifts ($\Delta\delta$) of the C9 protons, the $\Delta\delta$ of the methyl carbons (C16 and C17) of the *isopropyl* group, and the chemical shifts of C13 and H13 of the terminal methyl(hydroxy)methine group of the acid. However, since the isohemijunceines lack the C12 hydroxylation, it is doubtful these empirical observations can be applied as a general rule to the analogous data for the isohemijunceines (Table 5). Nonetheless, the most obvious difference in the ¹H NMR spectrum of **9** (peak 3, Figure 2) compared to **10** and **11** (peaks 5 and 7) is the large increase in the $\Delta\delta$ of the C9 protons and the downfield shift of the H12 (Table 5). Attempting to assign the relative configurations about C12 and C13 based upon the observations by Wiedenfeld and Röder¹⁸ leads to contradictions in this case and is therefore not possible. However, it is postulated that the significant changes in the ¹H chemical shifts for C9 and C12 of **9** relative to those for **10** and **11** do reflect a diastereoisomerism about C12. Furthermore, because of the very minor NMR spectra changes, the diastereoisomerism leading to isohemijunceines B (**10**) and C (**11**) is therefore expected to be about C13, analogous to supinine (**3**) and amabiline (**4**). There was no evidence for C13-based diastereoisomerism of isohemijunceine A (**9**).

Acetylated Derivatives of the Isohemijunceines: The *N*-oxides of the acetylated derivative of the m/z 284 alkaloids i.e., $MH^+/[2M + H]^+$ m/z 342/683 (peaks 19 and 21, Figure 2), and the much lower abundance free bases i.e., MH^+ m/z 326 (peaks 17 and 20, Figure 2), were observed in the stem and root extracts only. Examination of the MS/MS data for the *N*-oxides and the corresponding free bases (Table 1) supported a 7-acetoxy derivative by virtue of the abundant fragment ion at m/z 180.⁸ Without NMR data, the stereochemistry and hence the relationship of the acetylated derivatives to the three isohemijunceines remains undefined. However, analogous to isohemijunceines A, B and C (**9** – **11**), the MS/MS data for the two *N*-oxide peaks and the two free base peaks are very similar to each other in both cases. Therefore, it can be speculated that peaks 17 and 19 (Figure 2) may be 7-acetylisohemijunceine A and its *N*-oxide respectively while peaks 20 and 21 represent co-eluting 7-acetylisohemijunceines B and C and their *N*-oxides respectively.

Minor and Trace Levels of Additional Dehydropyrrolizidine Alkaloids: By assessing the MS and MS/MS data it is apparent that there are many dehydropyrrolizidine alkaloids, and their *N*-oxides, present at minor to trace levels. Because of their insignificance, at least

to overall levels of dehydropyrrolizidine alkaloids in *C. juncea*, only some of these have been included in Figure 2 and Table 1 but, mostly, there are insufficient data to allow confident speculation on structures.

Another alkaloid with MH^+ at m/z 284 was observed in the seed extract (peak 2, Figure 2). The MS/MS data (Table 1) were very similar to those for the isohemijunceines except for the relative abundances (Figure 3). The increased stability of the protonated molecule, the $[MH^+ - 44]$ ion at m/z 240 (base ion) and the m/z 156 ion were all in contrast to the facile elimination of the necic acid from the isohemijunceines that gives rise to the base ion at m/z 138 in their mass spectra. A possible explanation for this is that this minor alkaloid is the C7 esterified analogue of the isohemijunceines i.e., hemijunceine (**12**). Because of resonance stabilization of the resultant carbocation, the C9 allylic esters would be expected to be more fragile resulting in loss of the entire necic acid to yield the base ion at m/z 138.²³ The more stable C7 ester will have more opportunity for intra-necic acid fragmentation such as the loss of 44 Da to yield the base ion at m/z 240. This effect of C7 esterification versus C9 esterification has been previously recognized in the electron ionization MS of 7 and 9-angelylplatynecine²⁴ and in the reported more facile cleavage of a C9 allylic ester compared to a C7 ester.²⁵ Indeed, under the HPLC-ESI/MS conditions used in this study, 9-angelylheliotridine (**7**) (m/z (% abundance):238 (5, MH^+), 220(1), 156(4), 138(100), 120(14), 96(5), 94(3)) also showed the facile cleavage of the entire necic acid to yield a base ion at m/z 138 whereas the C7 ester (**6**) (m/z (% abundance):238 (4, MH^+), 220(7), 156(7), 138(31), 120(100), 108(6), 94(19)) showed an increase in abundance of the $[MH^+ - H_2O]$ fragment (m/z 220) and the intranecic acid fragmentation ion (m/z 156), and a base ion at m/z 120.

The stem extract included minor amounts of alkaloid *N*-oxides for which the MH^+ and $[2M + H]^+$ ions were observed at m/z 298/595 (peaks 8 and 14, Figure 2). One of the naturally occurring free base analogues (MH^+ m/z 282) was barely observable in the seed extract (peak 10, Figure 2) but was readily detected in the redox resin-reduced stem extract. The MS/MS data acquired for this free base (peak 10, Figure 2) showed a loss of 44 Da (m/z 282 $\rightarrow$ 238) followed by 100 Da (m/z 238 $\rightarrow$ 138) and therefore supported the presence of a methyl(hydroxy)methine derivative of senecioic acid giving rise to the 12,15-dehydro analogue (or co-eluting analogues) of one or more of the isohemijunceines. The MS/MS data for the *N*-oxides (peaks 8 and 14, Table 1), however, were quite different from each other, more so than might be expected for simple C7 versus C9 esterification. Because of the facile loss of a hydroxyl group (m/z 298 (50% abundance) $\rightarrow$ 281(100%)) it is possible that the minor peak 8 may be the *N*-oxide of the macrocyclic diester dicrotaline.⁹ MS/MS data for the reduction product from peak 8 were not acquired due to co-elution with more abundant alkaloids.

Quantitative Analysis of Roots, Stems, Leaves and Seeds

Since the calibration standards are free base dehydropyrrolizidine alkaloids and it has been shown that the esiMS response of an *N*-oxide and the corresponding free base can be quite different,²⁶ the *N*-oxides in the samples were reduced prior to analysis.⁸ HPLC-ESI/MS analysis, using a 0.1% formic acid-based mobile phase with acetonitrile with the Synergi Hydro column, confirmed complete capture of the alkaloids by the top column of the SCX-SPE stack and their subsequent efficient elution by the ammoniated methanol. The peak areas in reconstructed ion chromatograms of the major resultant free base dehydropyrrolizidine alkaloids i.e., junceine (**1**), trichodesmine (**2**), isohemijunceine A (**9**), isohemijunceines B (**10**) and C (**11**), and the acetylisohemijunceines I and II, were quantitated against 6-point calibration curves generated using **1** (for the macrocyclic diesters, $y = 0.23x^2 + 10.71x$, $R^2=0.998$) or lycopsamine (**8**) (for the monoesters, $y = 2.95x^2 + 12.3x$,

$R^2=0.999$) and expressed as a percentage of dry weight of plant (Table 6). The total amount of dehydropyrrolizidine alkaloids in seed was shown to be about 0.15%. The total dehydropyrrolizidine alkaloid content of roots, stems and leaves was determined to be about 0.05, 0.12 and 0.01% respectively. Despite these estimates, which would be expected to fluctuate with the growth stage of the plant, the main observation is that the previously unrecognized monoester dehydropyrrolizidine alkaloids i.e., isohemijunceines A, B and C (**9** – **11**) constitute the greater proportion of the total.

As a more direct comparison with the brief report by Ji *et al.*,⁶ an additional seed sample of the ‘Tropic Sun’ cultivar was extracted and analyzed using the quantitative NMR spectroscopy approach used by Ji *et al.*,⁶ but originally described by Molyneux *et al.*¹⁰ Thus, by quantitating the combined area of the C2 vinylic proton signals at 5.6 – 6.2 ppm against a known amount of *p*-dinitrobenzene, the total dehydropyrrolizidine alkaloid content was estimated to be 0.09% w/w (DW) of seed. A HPLC-ESI/MS analysis of this same sample extract, using 7-point calibration curves generated using trichodesmine (**2**), isolated from ‘Tropic Sun’ seed in this study, for the macrocyclic diesters ($y = 0.47x^2 + 2.43x - 0.16$, $R^2=0.999$) or **8** for the monoesters ($y = 1.63x^2 + 5.86x - 0.42$, $R^2=0.999$), returned an estimate of 0.1% dehydropyrrolizidine alkaloids (w/w, DW). This latter HPLC-ESI/MS estimate comprised 0.007% **1**, 0.009% **2** and 0.08% of the combined isohemijunceines (**9** – **11**). It should be noted that **1** has a poor solubility in chloroform¹³ and may not be totally accounted for in the chloroform-based analysis used in earlier studies.^{5, 6}

Implications of the Presence of Dehydropyrrolizidine Alkaloids in ‘Tropic Sun’

The major use of ‘Tropic Sun’ is as a green manure crop that, among other beneficial attributes, helps control plant-parasitic nematodes.^{2, 3} Related to this, Thoden and Boppre²⁷ have highlighted the potential for at least some of the nematode management properties of plant extracts or mulches to be attributed to the presence of dehydropyrrolizidine alkaloids. They further hypothesize that accumulation (or primary biosynthesis) in the roots as the water-soluble *N*-oxides may be a defense against root-pathogenic nematodes. Thus, the presence of the dehydropyrrolizidine alkaloids in the ‘Tropic Sun’ cultivar may be responsible, at least in part, for the anti-nematode properties of the plant.

The archived seed extract of *C. juncea*, sourced from India, previously analyzed by Williams and Molyneux⁵ was reconstituted in methanol and re-analyzed under the current HPLC-ESI/MS conditions. Not only were the expected macrocyclic diesters **1** and **2** observed but also the new isohemijunceines (**9** – **11**) were readily detected demonstrating that these metabolites are not unique to the ‘Tropic Sun’ cultivar. In their report Williams and Molyneux estimated that toxicity to chicks required levels of at least 1% w/w of pyrrolizidine alkaloids in the seed. They further speculated that levels of 0.1 to 0.3% of pyrrolizidine alkaloids in seed, if constituting greater than 1% of the diet for several weeks, might present a health hazard to animals and fowl.⁵

A major assumption in this study is that the positive ion esiMS responses of **1**, **2** and of the isohemijunceines (**9** – **11**) are similar to those of the calibration standards trichodesmine, junceine and lycopsamine respectively where used. With this *caveat*, the levels of dehydropyrrolizidine alkaloids determined in this study of the ‘Tropic Sun’ cultivar could be a cause for concern if simply compared to levels in *Senecio jacobaea* and *S. vulgaris*, for example, that are known problems to rangeland cattle in the USA with levels of between 0.09 and 0.18% total dehydropyrrolizidine macrocyclic diester alkaloids.¹⁰ However, with over half of the dehydropyrrolizidine alkaloids in ‘Tropic Sun’ being the isohemijunceines that, as monoesters, may be less toxic,²⁸ such simple comparisons may not be valid. Additionally, *Crotalaria* species that have been associated with livestock intoxication have been reported to produce higher levels of alkaloids and include *C. spectabilis* (0.01 – 1.52%

in leaves and 2.41 – 5.35% in seeds)²⁹ and *C. novae-hollandiae* subsp. *novae-hollandiae* (chemotype 2), *C. ramosissima*, *C. retusa* var. *retusa* and *C. crispata*, each producing a mean alkaloid content of 0.6, 2.23, 1.22 and 0.77% w/w respectively (expressed as retrorsine equivalents).³⁰ However, from the aspect of the potential for chronically-developing diseases in humans⁷ resulting from exposure to the *C. juncea* dehydropyrrolizidine alkaloids, it would be of value to determine and assess any such routes of potential exposure such as exposed animal-derived products (meat, milk), bee products (honey and pollen) and seed contamination of grain products.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

This research is supported by the Agricultural Research Service of the U.S. Department of Agriculture and an Interagency Agreement with the Office of Dietary Supplements of the National Institutes of Health.

Joe Stewart of the Utah Water Research Laboratory, Utah State University provided the high resolution mass measurement for the isohemijunceine diastereoisomers. Dr Simon Sham of the Chemistry Department, Utah State University assisted with the setting up of parameters for the 2D NMR experiments. Dr John Beck (Western Regional Research Center, Albany, CA, USA) and Dr Stephen Lee (Poisonous Plant Research Laboratory, Logan, UT, USA) are thanked for their contributions as ARS/USDA internal reviewers. The comments and advice of the anonymous reviewers are gratefully acknowledged.

References

1. Cook, CG.; White, GA. *Crotalaria juncea*: a potential multi-purpose fiber crop. In: Janick, J., editor. Progress in New Crops. ASHS Press; Arlington, VA: 1996. <http://www.hort.purdue.edu/newcrop/proceedings1996/v3-389.html>
2. United States Department of Agriculture. [Last accessed November 30th 2011] Sunn Hemp: A cover crop for southern and tropical farming systems. Soil Quality-Agronomy Technical Note No 10. 1999. http://soils.usda.gov/sqi/management/files/sq_atn_10.pdf
3. USDA Natural Resources Conservation Service. [Last accessed November 30th 2011] ‘Tropic Sun’ Sunn Hemp. 2009. <http://www.plant-materials.nrcs.usda.gov/pubs/hipmcrb8433.pdf>
4. Adams R, Gianturco M. The alkaloids of *Crotalaria juncea*. J Am Chem Soc. 1956; 78:1919–1921.
5. Williams MC, Molyneux RJ. Occurrence, concentration and toxicity of pyrrolizidine alkaloids in *Crotalaria* seeds. Weed Sci. 1987; 35:476–481.
6. Ji X, Khan I, Mosjidis JA, Wang H, Livant P. Variability for the presence of pyrrolizidine alkaloids in *Crotalaria juncea* L. Pharmazie. 2005; 60:620–622. [PubMed: 16124407]
7. Edgar JA, Colegate SM, Boppré M, Molyneux RJ. Pyrrolizidine alkaloids in food: a spectrum of potential health consequences. Food Addit Contam: Part A. 2011; 28:308–324.
8. Colegate SM, Edgar JA, Knill AM, Lee ST. Solid phase extraction and LCMS profiling of pyrrolizidine alkaloids their *N*-oxides: a case study of *Echium plantagineum*. Phytochem Anal. 2005; 16:108–119. [PubMed: 15881119]
9. Bull, LB.; Culvenor, CCJ.; Dick, AT. The Pyrrolizidine Alkaloids: their Chemistry, Pathogenicity and other Biological Properties. North-Holland; Amsterdam: 1968. p. 60-66.
10. Molyneux RJ, Johnson AE, Roitman JN, Benson ME. Chemistry of toxic range plants. Determination of pyrrolizidine alkaloid content and composition in *Senecio* species by nuclear magnetic resonance spectroscopy. J Agric Food Chem. 1979; 27:494–499. [PubMed: 447923]
11. Hartmann, T.; Witte, L. Chemistry, biology, and chemoeology of the pyrrolizidine alkaloids. In: Pelletier, SW., editor. Alkaloids: Chemical and Biological Perspectives. Vol. 9. Pergamon Press; Oxford: 1995. p. 155-233.

12. Peñaloza GC, Peláez JCA. Evaluation of biological activity of *Crotalaria juncea* extracts through a *Drosophila melanogaster* model. *Vitae*. 2008; 15:279–284.
13. Atal CK, Sharma RK, Culvenor CCJ, Smith LW. Alkaloids of *Crotalaria rubiginosa* Willd. Trichodesmine and junceine. *Aust J Chem*. 1966; 19:2189–2191.
14. Roeder E, Sarg T, El-Dahmy S, Abdel Ghani A. Pyrrolizidine alkaloids from *Crotalaria aegyptiaca*. *Phytochemistry*. 1993; 34:1421–1423.
15. Culvenor CCJ, Edgar JA, Frahn JL, Smith LW, Ulubelen A, Doganca S. The structure of anadoline. *Aust J Chem*. 1975; 28:173–178.
16. Röder E. Medicinal plants in Europe containing pyrrolizidine alkaloids. *Pharmazie*. 1995; 50:83–98. [PubMed: 7700976]
17. Williams MT, Warnock BJ, Betz JM, Beck JJ, Gardner DR, Lee ST, Molyneux RJ, Colegate SM. Detection of high levels of pyrrolizidine-*N*-oxides in the endangered plant *Cryptantha crassipes* (Terlingua Creek Cat's-eye) using HPLC-ESI-MS. *Phytochem Anal*. 2011; 22:532–540. [PubMed: 21433162]
18. Wiedenfeld H, Röder E. Pyrrolizidine alkaloids from *Ageratum conyzoides*. *Planta Med*. 1991; 57:578–579. [PubMed: 17226207]
19. Weber S, Eisenreich W, Bacher A, Hartmann T. Pyrrolizidine alkaloids of the lycopsamine type: biosynthesis of trachelanthic acid. *Phytochemistry*. 1999; 1005–1014.
20. Logie CG, Grue MR, Liddell JR. Proton NMR spectroscopy of pyrrolizidine alkaloids. *Phytochemistry*. 1994; 37:43–109.
21. Jones AJ, Culvenor CCJ, Smith LW. Pyrrolizidine alkaloids- A carbon-13 NMR study. *Aust J Chem*. 1982; 35:1173–1184.
22. Mohanraj S, Herz W. High resolution ¹H-NMR and ¹³C-NMR spectra of saturated pyrrolizidine monoester alkaloids. *J Nat Prod*. 1982; 45:300–336.
23. Pedersen E, Larsen E. Mass spectrometry of some pyrrolizidine alkaloids. *J Mass Spectrom*. 1970; 4:249–256.
24. Stelljes ME, Kelley RB, Molyneux RJ, Seiber JN. GC-MS determination of pyrrolizidine alkaloids in four *Senecio* species. *J Nat Prod*. 1991; 54:759–773.
25. Roitman JN. The pyrrolizidine alkaloids of *Senecio triangularis*. *Aust J Chem*. 1983; 36:1203–1213.
26. Boppré M, Colegate SM, Edgar JA. Pyrrolizidine alkaloids of *Echium vulgare* honey found in pure pollen. *J Agric Food Chem*. 2005; 53:594–600. [PubMed: 15686407]
27. Thoden TC, Boppré M. Plants producing pyrrolizidine alkaloids: sustainable tools for nematode management? *Nematology*. 2010; 12:1–24.
28. Culvenor CCJ, Edgar JA, Jago MV, Outteridge A, Peterson JE, Smith LW. Hepato- and pneumotoxicity of pyrrolizidine alkaloids and derivatives in relation to molecular structure. *Chemico-Biol Interact*. 1976; 12:299–324.
29. Johnson AE, Molyneux RJ, Merrill GB. Chemistry of toxic range plants. Variation in pyrrolizidine alkaloid content of *Senecio*, *Amsinckia*, and *Crotalaria* species. *J Agric Food Chem*. 1985; 33:50–55.
30. Fletcher MT, McKenzie RA, Blaney BJ, Reichmann KG. Pyrrolizidine alkaloids in *Crotalaria* taxa from northern Australia: risk to grazing livestock. *J Agric Food Chem*. 2009; 57:311–319. [PubMed: 19061310]

Figure 1. Structures of some dehydropyrrolizidine alkaloids observed in *Crotalaria juncea*, or used in this study.

Figure 2.

The HPLC-ESI/MS base ion (m/z 200 – 800) chromatograms of the aqueous acid-soluble fraction of the methanolic extract of the stems (**A**) and seeds (**B**) of *Crotalaria juncea* (cv. 'Tropic Sun'). Peak numbers identify dehydropyrrolizidine alkaloids and are referred to in the text and/or in Table 1.

Figure 3.

The HPLC-ESI/MS, reconstructed ion chromatogram (**A**) displaying m/z 284 for the extract of seeds of *Crotalaria juncea* (cv. 'Tropic Sun'). The peak numbers in **A** correspond to Figure 2. The subtle structural difference is highlighted by the MS/MS fragmentation profile of : **B**. hemijunceine (**12**) (peak 2); and **C**. the isohemijunceines A (**9**), B (**10**) and C (**11**) (peaks 3, 5 and 7 respectively).

Table 1

Positive Ion Electrospray Ionization, Ion Trap MS/MS Data for the Dehydropyrrolizidine Alkaloids Detected in Extracts of *Crotalaria juncea* (cv. 'Tropic Sun').

Peak Number ^a	Identity	MH ⁺ /[2M + H] ⁺ (m/z)	MS/MS ^b (% abundance)
1	junceine (1)	370	370(20), 352(15), 340(7), 324(100), 282(8), 267(9), 266(58), 265(7), 244(14), 226(53), 223(20), 222(36), 190(5), 180(23), 162(8), 122(18), 121(7), 120(28)
2	hemijunceine (12)	284	284(42), 240(100), 214(2), 156(45), 138(36), 120(10), 108(5), 94(2)
3	isohemijunceine A (9)	284	284(<1), 240(2), 222(1), 204(1), 156(13), 138(100), 120(18), 94(38)
4	junceine- <i>N</i> -oxide	386(30)/771(100) ^c	386(14), 368(96), 358(6), 340(100), 298(20), 282(12), 281(8), 264(22), 260(10), 256(6), 242(6), 239(10), 238(21), 225(7), 224(4), 137(10), 120(10)
5	isohemijunceine B (10)	284	284(1), 266(1), 240(2), 222(1), 156(9), 138(100), 120(17), 94(28)
6	trichodesmine (2)	354	354(21), 336(3), 308(100), 266(14), 265(9), 228(8), 222(30), 210(21), 192(5), 164(14), 148(3), 138(1), 136(3), 122(4), 121(5), 120(7)
7	isohemijunceine C (11)	284	284(<1), 240(1), 222(1), 156(9), 138(100), 120(14), 94(33)
8	dehydroisohemijunceine- <i>N</i> -oxide I	298(70)/595(100)	298(51), 282(9), 281(100), 264(9), 238(8), 220(29), 172(4), 155(17), 154(27), 137(18), 136(26), 124(7), 120(3), 118(6), 111(5), 109(9), 106(44)
9	isohemijunceine A- <i>N</i> -oxide	300(50)/599(100)	300(57), 283(4), 282(13), 256(100), 172(45), 155(22), 154(22), 139(13), 138(18), 137(28), 136(40), 118(3), 112(6), 108(5), 94(5)
10	dehydroisohemijunceine I	282	282(14), 254(1), 238(2), 164(2), 156(1), 138(100), 120(33), 110(2), 94(73)
11	isohemijunceine B- <i>N</i> -oxide	300(42)/599(100)	300(51), 283(4), 282(16), 256(100), 172(40), 155(22), 154(19), 150(2), 138(9), 137(29), 136(37), 112(4), 111(6), 109(2), 108(3), 94(4)
12	trichodesmine- <i>N</i> -oxide	370/739(20)	370(1), 352(5), 324(100), 298(7), 282(4), 281(6), 264(11), 244(6), 239(14), 238(23), 226(6), 209(3), 180(2), 176(3), 164(2), 138(4), 137(9), 120(6)
13	isohemijunceine C- <i>N</i> -oxide	300(45)/599(100)	300(53), 282(11), 256(100), 172(40), 155(20), 154(23), 139(3), 138(11), 137(34), 136(33), 121(2), 120(2), 109(3), 108(4), 94(4)
14	dehydroisohemijunceine - <i>N</i> -oxide II	298(89)/595(100)	298(63), 281(2), 280(4), 238(1), 237(3), 172(100), 155(16), 154(37), 138(11), 137(33), 136(44), 126(3), 112(3), 111(3), 109(1), 108(2), 94(4)
15	unidentified macrocyclic diester <i>N</i> -oxide	354(40)/707(100)	354(100), 337(4), 336(3), 326(8), 308(7), 292(6), 267(3), 222(2), 156(2), 155(11), 154(10), 138(39), 137(40), 119(3), 118(19), 109(2), 108(6)
16	senecionine	336	336(86), 318(4), 308(94), 290(14), 199(7), 153(9), 138(71), 120(100)
17	acetylisohemijunceine I	326	198(14), 180(83), 162(3), 120(100), 118(2)
18	unknown <i>N</i> -oxide	368(85)/735(100)	368(100), 340(55), 306(15), 264(31), 247(14), 237(12), 236(66), 222(13), 219(14), 218(42), 206(13), 154(26), 136(18), 122(11), 121(22), 120(30), 119(57), 118(50)
19	acetylisohemijunceine - <i>N</i> -oxide I	342(40)/683(100)	342(51), 324(28), 298(100), 282(6), 214(72), 197(14), 180(14), 179(8), 178(28), 154(9), 137(4), 136(8), 121(5), 120(9)

Peak Number ^a	Identity	MH ⁺ /[2M + H] ⁺ (m/z)	MS/MS ^b (% abundance)
20	acetylisohemijunceine II	326	198(17), 180(61), 162(4), 120(100), 118(1)
21	acetylisohemijunceine -N-oxide II	342(44)/683(100)	342(38), 324(23), 298(100), 282(4), 214(55), 197(15), 180(7), 179(6), 178(23), 154(5), 137(4), 136(8), 120(7)
22	unidentified macrocyclic diester	338	338(50), 310(100), 292(3), 222(1), 138(11), 120(11),
23	unidentified dihydro macrocyclic diester	340	340(100), 312(40), 278(2), 140(3), 122(8)

^a peak numbers refer to Figure 2

^b using a dissociation energy of 32% for the MH⁺ ion

^c numbers in parentheses are % relative abundances

Table 2

^1H , ^{13}C , ^1H - ^1H (COSY) and ^1H - ^{13}C (HMBC) NMR Spectroscopic Data for Isohemijunceine A (9) Extracted from *Crotalaria juncea* (cv. 'Tropic Sun').

Carbon ^a	^1H δ ^b	^{13}C δ ^c	COSY ^d (proton number)	HMBC ^{d, e} (proton number)
1	na ^f	133.38	na	9d (m), 9u (w), 3u (w)
2	5.85 (s)	129.83	3u ^g (w), 3d (w), 8(w), 9u (w), 9d (w)	9d (m), 9u (w), 3u (w)
3	3.92 (d, $J_{3d,3u}=15.4$)	63.1	2 (w), 3u, 8 (w), 9u (w)	2 (w), 5u (m)
	3.40 (dd, $J_{3u,3d}=15.5$, $J_{3u,2}=7.4$)		2 (w), 3d, 8 (m), 9u (w)	
5	3.26 (m)	54.01	5u, 6 (w-m)	3u (w), 6 (w), 7 (m)
	2.73 (ddd, $J_{5u,5d}=11.6$, $J_{5u,6}=9.1$ and 6)		5d, 6 (m-s)	
6	ca 1.86 - 2.01 (m, 2H)	35.97	5u (m-s), 5d (w-m), 7 (w)	5d (m), 5u (m)
7	4.29 (t, $J=2.7$)	71.27	6 (w), 8 (m)	5d, 6,
8	4.16 (bs)	78.94	2 (w), 3u (m), 3d (w), 7 (m)	2 (w), 9d, 9u (w), 7 (w-m), 5d, 6d, 6u
9	4.92(d, $J=13.0$)	61.26	2 (w), 9u	5u (w),
	4.51(d, $J=13.0$)		2 (w), 3d (w), 3u (w), 9d	
11	na	173.43	na	9d (m), 12, 13, 15 (w-m)
12	2.36 (dd, $J=8.2$, 6.2)	59.16	13 (m-s), 15(w)	13, 15, 14, 16, 17
13	4.09 (p, $J=6.3$)	66.91	12 (m-s), 14	12, 14, 15 (w), 16(w), 17 (w)
14	1.21 (d, $J=6.3$)	19.7	13	12, 13
15	ca 2.0 - 2.13 (m)	27.49	12 (w), 16/17	12, 13, 14 (w), 16, 17
16	0.96 (d, $J=5$)	20.63	15	12, 15, 17
17	0.94 (d, $J=4.8$)	20.81	15	12, 15, 16

^a carbon numbering shown in Figure 1

^b bs = broad singlet; m = multiplet; d = doublet; t = triplet; q = quartet; J in Hz

^c assignments of ^{13}C resonances were unambiguously established *via* a gradient enhanced HSQC experiment

^d strong interactions unless otherwise annotated (m) medium; (w) weak

^e data shown as proton to carbon correlations

^f na = not applicable

^g u = upfield; d = downfield partner of a geminal proton pair

Table 3

^1H , ^{13}C , and ^1H - ^1H (COSY) NMR Spectroscopic Data for Isohemijunceine B (10) Extracted from *Crotalaria juncea* (cv. 'Tropic Sun').

Carbon ^a	^1H δ ^b	^{13}C δ ^c	COSY ^d (proton number)
1	na ^e	132.98	na
2	5.87 (s)	129.14	3u ^f (m), 3d (m), 7 or 8 (m), 9u (m), 9d (m)
3	4.05 (d, $J_{3d,3u}=15$)	62.6	2 (m), 3u, 7 or 8 (w), 9d (w-m), 9u (w-m)
	3.47(bd, $J_{3d,3u}=15$)		2 (m), 3d, 7 or 8 (m), 9d (w), 9u (w)
5	3.41 (bm)	54.04	5u, 6 (m)
	2.82 (bm)		5d, 6
6	ca2.03 (m, 2H)	35.98	5u, 5d (m), 7 (w)
7	4.35(bs, 2H))	71.04	6 (m), 2 (m), 3u (m), 3d (w)
8		78.89	
9	4.83 (d, $J=13.2$)	61.16	2 (m), 3d (w), 3u (w), 9u
	4.67 (dd, $J=13.1, 1.5$)		2 (m), 3d (w), 3u (w), 9d
11	na	174.44	na
12	2.13 (m)	59.75	13 (m-s)
13	4.05(m)	66.25	12 (m), 14
14	1.22 (d, $J=6.3$)	22.28	13
15	ca 2.09 (m)	27.79	16, 17
16	1.01 (d, $J=6.4$)	21.03	15
17	0.93 (d, $J=6.5$)	20.06	15

^a carbon numbering shown in Figure 1

^b bs = broad singlet; m = multiplet; d = doublet; t = triplet; q = quartet; J in Hz

^c assignments of ^{13}C resonances were unambiguously established *via* a gradient enhanced HSQC experiment

^d strong (s) interactions unless otherwise annotated (m) medium; (w) weak

^e na = not applicable

^f u = upfield; d = downfield partner of a geminal proton pair.

Table 4

^1H , ^{13}C , ^1H - ^1H (COSY) and ^1H - ^{13}C (HMBC) NMR Spectroscopic Data for Isohemijunceine C (11) Extracted from *Crotalaria juncea* (cv. ‘Tropic Sun’).

Carbon ^a	^1H δ ^b	^{13}C δ	COSY ^c (proton number)	HMBC ^{d, e} (proton number)
1	na ^f	132.99	na	2 (w-m), 3d, 3u, 9d, 9u
2	5.87 (bs)	130.43	3d ^g , 3u (m), 8 (w-m), 9d (m), 9u	3d, 3u, 9d, 9u
3	4.03 (bd)	63.06	2 (m), 3u, 8 (w)	2, 5u,
	3.46 (bd)		2 (m), 3d, 8	
5	3.40 (m)	54.01	5u, 6d, 6u	3d (w), 3u, 7
	2.82 (m)		5d, 6d, 6u	
6	ca 1.9 – 2.03 (2H, m)	36.08	5d, 5u, 7	5u
7	4.35 (bs)	71.16	6d, 6u,	3d (m-s), 5d, 6,
8	4.31 (bs)	78.89	2 (w-m), 3d (m), 3u, 5u	2, 3d (m), 5d, 6, 7 (m), 9d, 9u (w),
9	4.87 (bd, $J = 13.2$)	61.36	2, 9u, 3d (m), 3u (m)	2,
	4.63 (bd, $J = 13.2$)		2, 9d, 3d (m-s), 3u (m)	
11	na	174.84	na	9d, 9u (w), 12, 13, 15
12	ca 2.10 – 2.19 (m)	59.52	13	13, 14, 15, 16, 17
13	4.05 (m)	66.41	12, 14	12, 14, 15 (w)
14	1.20 (d, $J_{14,13} = 6.3$)	22.29	13	12 (w), 13
15	ca 2.02 – 2.1 (m)	27.85	16, 17	12, 13, 16, 17
16	1.01 (d, $J_{16,15} = 6.6$)	21.0	15	12, 15, 17
17	0.93 (d, $J_{17,15} = 6.6$)	20.02	15	12, 15, 16

^a carbon numbering shown in Figure 1

^b m = multiplet; s = singlet; d = doublet; t = triplet; q = quartet; b = broad; J in Hz

^c assignments of ^{13}C resonances were unambiguously established *via* a gradient enhanced HSQC experiment

^d strong interactions unless otherwise annotated (m) medium; (w) weak

^e data shown as proton to carbon correlations

^f na = not applicable

^g u = upfield; d = downfield partner of a geminal proton pair

Table 5

Some Significant Differences in the NMR Spectra for the Isohemijunceines A (9), B (10) and C (11) of Potential Stereochemical Diagnostic Value.¹⁸

Isohemijunceine	$\Delta\delta$ C9 protons	δ H12	δ C13 $^1\text{H}/^{13}\text{C}$	^{13}C $\Delta\delta$ C16/C17
A (9)	0.41	2.36	4.09/66.91	0.18
B (10)	0.16	2.13	4.05/66.25	0.97
C (11)	0.24	2.10 – 2.19	4.05/66.41	0.98

Table 6

Total Concentrations of Major Dehydropyrrolizidine Alkaloids in Roots, Stems, Leaves and Seeds of *Crotalaria juncea* (cv. 'Tropic Sun') Following Reduction of *N*-Oxides to Free Bases.

Dehydropyrrolizidine Alkaloid	Alkaloid Concentration (% w/w)			
	Plant Part			
	roots	stems	leaves	seeds
junceine (1) ^a	0.021	0.026	0.003	0.03
trichodesmine (2) ^a	0.007	0.013	0.001	0.02
isohemijunceine A (9) ^b	0.007	0.009	0.001	0.03
isohemijunceine B (10) and C (11) ^b	0.014	0.023	0.002	0.07
acetylisohemijunceine I and II ^b	0.004	0.044	0.001	0.00
Total (% w/w)	0.053	0.115	0.008	0.15

^a quantitated from the junceine calibration curve

^b quantitated from the lycopsamine calibration curve