Published in final edited form as:

Org Lett. 2012 June 1; 14(11): 2862–2865. doi:10.1021/ol301107m.

Total Syntheses of Graphisin A and Sydowinin B¹

Andrew Little and John A. Porco Jr.*

Department of Chemistry and Center for Chemical Methodology and Library Development CMLD-BU), Boston University, Boston Massachusetts 02215

Abstract

Efficient syntheses of the highly substituted benzophenone graphisin A and the xanthone sydowninin B are described. Key steps involve aryl anion addition to substituted benzaldehyde derivatives, subsequent methyl ester installation, and dehydrative cyclization. Oxidation of graphisin A led to a spirodienone derived from a highly substituted benzoquinone intermediate.

Acremoxanthone A² (**1**), xanthoquinodin A³ (**2**), and acremonidin A (**3**)⁴ (Figure 1) are members of a class of heterodimeric natural products originally isolated from *Acremonium* sp. BCC31806, *Humicola* sp. FO888 and *Acremonium* sp., LL-Cyan 416, respectively. These natural products share structurally related anthraquinone and xanthone-derived monomer units which are linked by a unique bicyclo[3.2.2] ring system.⁵ We have initiated synthetic studies toward these targets with syntheses of the requisite xanthone and benzophenone fragments present in the natural product frameworks. In our initial studies, we have targeted the highly functionalized, tetra-*ortho*-substituted benzophenone graphisin A⁶ (**4**) and the corresponding cyclized xanthone sydowninin B⁷ (**5**). Herein, we report the total synthesis of the highly substituted benzophenone⁸ graphisin A using a suitably functionalized benzophenone as well as preparation of the derived xanthone sydowninin B *via* dehydrative cyclization.

Our initial plan (Figure 2) was to prepare benzophenone nitrile **6** *via* aryl anion addition to a substituted benzaldehyde according to literature precedent.⁹ We envisioned that acidic methanolysis¹⁰ of aryl nitrile **6** followed by global deprotection should provide graphisin A (**4**). Sydowninin B (**5**) could then be accessed through dehydrative cyclization.¹¹

In our forward synthesis, benzaldehyde **7** was regioselectively brominated and protected as a *bis*-MOM ether in 82% yield over two steps (Scheme 1). Regioselective deprotonation of **8** with *n*-BuLi, followed by addition of aldehyde **9** at 0 °C, provided benzylic alcohol **10** in 56% yield with the remainder of the mass balance consisting of byproducts derived from competitive lithium-halogen exchange of the *ortho* aryl bromide. Oxidation of **10** with 2-iodoxybenzoic acid (IBX) provided bromobenzophenone **11** (75%). We had initially planned to utilize **11** directly in palladium-catalyzed carbonylative methyl ester formation (Pd(dppf)Cl₂, CO, NEt₃, DMF/MeOH);¹² however, this substrate was found to be unreactive likely due to the highly electron rich and *bis-ortho*-substituted aryl ring. Efforts to

*porco@bu.edu.

Supporting Information Available. Detailed experimental procedures and spectral data for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

install the methyl ester on either bromobenzaldehyde **9** or its corresponding acetal-protected derivatives were also unsuccessful. Similarly, anion chemistry (*e.g.* lithium halogen exchange, Grignard formation) of aryl bromide substrates was also found to be unproductive.^{8c} However, compound **11** was found to undergo smooth cyanation using Pd(PPh₃)₄ and Zn(CN)₂ under microwave heating to afford aryl nitrile **6** in 60% yield.¹³

Initial attempted hydrolysis of aryl nitrile **6** to the derived acid or ester **12** (aq NaOH or H₂SO₄) resulted in either decomposition or partial MOM deprotection (Scheme 2). Attempted partial hydrolysis of the primary amide with K₂CO₃ and H₂O₂¹⁴ resulted in no reaction. Similarly, treatment of **6** with methyl triflate in CH₂Cl₂ followed by a methanolysis to access the imidate in a Ritter-type reaction was also unsuccessful.¹⁵ Attempts to hydrolyze the nitrile in the presence of methanol (BF₃,¹⁶ FeCl₃,¹⁷ TMSCl,¹⁸ and H₂SO₄¹⁹) resulted in mixtures of partially and fully MOM-deprotected compounds with the aryl nitrile remaining intact.

In light of the apparent incompatibility of the MOM protecting groups with nitrile hydrolysis conditions, we elected to fully deprotect benzophenone **6** to tetraphenol **13** prior to hydrolysis (Scheme 2). Accordingly, treatment of nitrile **6** with *p*-TsOH in MeOH at 40 °C produced a deep red solution whereby the protecting groups were globally removed to afford benzophenone **13** in 50% yield. Purification of polyphenol **13** on silica gel was found to be difficult due its high polarity and propensity to form an intractable, red-colored salt with residual *p*-TsOH. An example of a highly substituted benzophenone nitrile hydrolysis has been reported.²⁰ However, standard acidic hydrolysis⁹ (H₂SO₄ or HCl, MeOH, 60 °C) of **13** resulted in recovered starting material and minor formation of xanthone **14**, whereas standard basic conditions²¹ (NaOH) resulted in decomposition.

At this juncture, we elected to optimize the xanthone formation / nitrile methanolysis sequence. We observed dehydrative cyclization to the xanthone when attempting to hydrolyze nitrile **13** with DOWEX-50 resin in water under microwave heating.²² We later observed clean conversion of **13** to xanthone **14** using microwave heating with aqueous ZnCl₂.²³ A ground state model of nitrile **13** (Figure 3) shows that the benzophenone exists substantially in a bisected conformation with hydrogen bonding maintained between the C-6 and C-10 phenols and the C-8 ketone. However, internal hydrogen-bonding of the phenols may be disrupted in water such that rotation can occur prior to cyclization to form xanthone **14**. Xanthone **14** was finally subjected to acidic methanolysis conditions (3M H₂SO₄, MeOH, 60 °C) which afforded sydownin B (**5**) in 50% yield (two steps).

To enable installation of the requisite methyl ester and access graphisin A (**4**) without dehydrative cyclization, we implemented the alternative route shown in Scheme 3. Bromoaldehyde²⁴ **9** was reduced and benzyl-protected to afford aryl bromide **15** in 80% yield (two steps). Regioselective deprotonation of **8** and subsequent quenching with DMF provided benzaldehyde **16** (78%). Lithium halogen exchange of aryl bromide **15** at –78 °C in ether with *t*-BuLi followed by a rapid quench with aldehyde **16** provided benzyl alcohol **17** (61%). Subsequent IBX oxidation afforded ketone **18** in 78% yield (Scheme 4). Initial attempted conditions for selective benzyl deprotection of **18** (H₂, Pd(OH)₂, MeOH) resulted in complete conversion to the over-reduced product **19**, analogous to a product²⁵ reported by Snider and coworkers for a similar substrate.²⁶ However, hydrogenolysis of **18** using Pd/C in THF/MeOH (3:1) led to the desired hydroxybenzophenone **20** (Scheme 5). It is apparent that sole use of a polar protic solvent activates **20** toward hemiketal formation which undergoes further hydrogenolysis to the corresponding dihydroisobenzofuran **19**.

A three-step oxidation-methylation sequence was next conducted on the protected benzophenone **20** involving sequential Dess-Martin and Pinnick oxidations followed by

treatment with excess TMSCH₂N₂ in 1:1 MeOH/Et₂O to provide methyl ester **12** in 65% yield over 3 steps (Scheme 5).²⁵ Compound **12** was fully deprotected with *p*-TsOH in MeOH to provide graphisin A (**4**) in 77% yield. Analytical data for **4** including ¹H and ¹³C NMR spectra were in agreement with those reported for the natural product.⁶ Graphisin A (**4**) could also be converted to sydownin B (**5**) in 58% yield by dehydrative cyclization with aqueous ZnCl₂ under microwave conditions (120 °C).²³

It has been proposed that the biosynthesis of various members of the blennolide/secalonic acid²⁷ family of natural products proceeds through oxidation/reduction of functionalized benzophenone intermediates resembling graphisin A.²⁸ Accordingly, we conducted preliminary investigations as to whether graphisin A (**4**) could be converted to dienone **22** *via* oxidation to benzoquinone **21** followed by *oxa*-Michael addition (Scheme 6).²⁹ Treatment of graphisin A (**4**) with phenyliodine diacetate (PIDA) resulted in clean oxidative cyclization to spirodienone **23**.³⁰ Oxidations of hydroxybenzophenones to similar spirodienones are known^{25,31} and are thought to occur *via* a phenoxide radical that is aligned to favor the 5-membered ring due to a bisected conformation of the highly substituted benzophenone. Spirodienone **23** was treated under acidic conditions²⁵ to provide a mixture of rearranged acid **24** (57%) and depsidone **25** (12%). Acid **24** could be further cyclized to depsidone **25** by heating with TFAA in toluene (68%). In a similar fashion, oxidation of **4** with FeCl₃ provided a mixture of **24** (39%) and **25** (44%) as the sole products.

Following reported dehydrogenation conditions that likely proceed through quinone intermediates,³² treatment of graphisin A (**4**) with Pd/C-O₂ also provided spirodienone **23** in 89% yield. This result supports the intermediacy of benzoquinone **21** enroute to **23**. While both 6-*endo-trig* and 5-*exo-trig* cyclization modes of **21** are possible,³³ the observed isomer **23** is likely favored due to the bisected conformation³⁴ of proposed intermediate **21** which places one *ortho*-phenol in favorable alignment with the π*-orbital of the C-7 quinone carbon (Figure 4).

In conclusion, an efficient synthesis of graphisin A has been accomplished employing aryl anion addition to a benzaldehyde, followed by oxidation to a highly substituted benzophenone, subsequent methyl ester installation, and global deprotection. Graphisin A was further converted by dehydrative cyclization under microwave conditions to the xanthone sydownin B. Oxidation of graphisin A led to formation of a spirodienone likely from a highly substituted and bisected benzoquinone intermediate. Further studies employing graphisin A and sydownin B toward the synthesis of acremoxanthones and related natural products are currently in progress and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support from the National Institute of Health (RO1 CA137270 and CA 099920) is gratefully acknowledged. We thank Prof. P. Pittayakhajonwut, Bioresources Research Laboratory, Thailand for providing a sample of natural graphisin A and Tian Qin (Boston University) for helpful discussions.

References

- (1). Presented in part at the 240th American Chemical Society National Meeting; Boston. Aug 22-26, 2010; abstract ORGN 410
- (2). Isaka M, Palasarn S, Auncharoen P, Komwijit S, Jones EB. Tetrahedron Lett. 2008; 50:284–287.
- (3). Tabata N, Tomoda H, Matsuzaki K, Omura S. J. Am. Chem. Soc. 1993; 115:8558–8564.

- (4). He H, Bigelis R, Solum E, Carter G. J. Antibiot. 2003; 56:923–930. [PubMed: 14763558]
- (5). (a) Milat ML, Praange T, Ducrot PH, Tabet JC, Einhorn J, Blein JP, Lallelmand JY. J. Am. Chem. Soc. 1992; 114:1478–1479. (b) Jalal MA, Hossain MB, Robeson DJ, Helm D. J. Am. Chem. Soc. 1992; 114:5967–5971.
- (6). Pittayakhajonwut P, Sri-indrasutdhi V, Dramaie A, Lapanun S, Suvannakad R, Tantichareon M. Aust. J. Chem. 2009; 62:389–391.
- (7). Hamasaki T, Sato Y, Hatsuda Y. Agr. Biol. Chem. 1975; 39:2341–2345.
- (8). (a) Nicolaou KC, Bunnage ME, Koide K. J. Am. Chem. Soc. 1994; 116:8402–8403. (b) Kaiser F, Schwink L, Velder J, Schmalz HG. J. Org. Chem. 2002; 67:9248–9256. [PubMed: 12492326] (c) Slavov N, Cvengros J, Neudorfl JM, Schmalz HG. Angew. Chem. Int. Ed. 2010; 49:7588–7591. For syntheses of highly substituted benzophenones, see:
- (9). Hollinshead SP, Nichols JB, Wilson JW. J. Org. Chem. 1994; 59:6703–6709.
- (10). Clarke HT, Taylor ER. Org. Synth. Collect. 1943; 2:588.
- (11). (a) Jeso V, Nicolaou KC. Tetrahedron Lett. 2009; 50:1161–1163. [PubMed: 20161287] (b) Sousa ME, Pinto MM. Curr. Med. Chem. 2005; 12:2447–2479. [PubMed: 16250872] (c) Diderot NT, Silvere N, Etienne T. Adv. Phytomed. 2006; 2:273–298. (d) Pinto MMM, Castanheiro RAP. Natural Prod. 2009:520–675. For recent reviews of xanthone syntheses, see:
- (12). Brennfuhrer A, Neumann H, Beller M. Angew. Chem. Int. Ed. 2009; 48:4114–4133.
- (13). Tschaen DM, Desmond R, King AO, Fortin MC, Pipik B, King S, Verhoeven TR. Synth. Commun. 1994; 24:887–890.
- (14). Katritzky AR, Pilarski B, Urogdi L. Synthesis. 1989; 12:949–950.
- (15). Booth BL, Jibodu KO, Proenca MF. J. Chem. Soc. Perkin Trans. 1. 1983:1067–1068.
- (16). Jayachitra G, Yasmeen N, Srinivasa K, Ralte SL, Singh AK. Synth. Commun. 2003; 33:3461–3466.
- (17). Srinivasan R, Rao KS, Jayachitra G, Ralte SL. Synthetic Commun. 2006; 36:2883–2886.
- (18). Luo FT, Jeevanadam A. Tetrahedron Lett. 1998; 39:9455–9456.
- (19). Anzalone L, Hirsch JA. J. Org. Chem. 1985; 50:2128–2133.
- (20). Storm JP, Andersson CM. J. Org. Chem. 2000; 65:5264–5274. [PubMed: 10993355]
- (21). Staab HA, Hauck R, Popp B. Eur. J. Org. Chem. 1998; 1:631–642.
- (22). Lindstrom U. Green Chem. 2006; 8:22–24.
- (23). Manjula K, Pasha MA. Synth. Commun. 2007; 37:1545–1550. For Zn (II)-mediated dehydration of amides in water, see:
- (24). Goddard ML, Tabacchi R. Tetrahedron Lett. 2006; 47:909–911.
- (25). Yu M, Snider BB. Org. Lett. 2011; 13:4224–4227. [PubMed: 21793536]
- (26). We observed aryl and MOM ether peak broadening in the ^1H and ^{13}C NMR spectra for compound **19** similar to a 2,6-disubstituted aryl dihydroisobenzofuran reported in reference 25 (see Supporting Information for details).
- (27). (a) Zhang W, Krohn K, Flörke U, Pescitelli G, Di Bari L, Antus S, Kurtan T, Rheinheimer J, Draeger S, Schulz B. Chem. Eur. J. 2008; 14:4913–4923. [PubMed: 18425741] (b) Gerard EMC, Bräse S. Chem. Eur. J. 2008; 14:8086–8089. [PubMed: 18720484] (c) Qin T, Johnson RP, Porco JA. J. Am. Chem. Soc. 2011; 133:1714–1717. [PubMed: 21265529]
- (28). (a) Kurobane I, Vining LC. Tetrahedron Lett. 1978; 16:1379–1382. (b) Kurobane I, Vining LC, McInnes AC. J. Antibio. 1979:1256–1266. (c) Bräse S, Encinas A, Keck J, Nising CF. Chem. Rev. 2009; 109:3903–3990. [PubMed: 19534495]
- (29). (a) Franck B, Stockigt U, Franckowiak G. Chem. Ber. 1973; 106:1198–1220. (b) Nicolaou KC, Li A. Angew. Chem. Int. Ed. 2008; 120:6681–6684. (c) Bröhmer MC, Bourcet E, Nieger M, Bräse S. Chem. Eur. J. 2011; 17:13706–13711. [PubMed: 22076837] (d) Tisdale EJ, Kochman DA, Theodorakis EA. Tetrahedron Lett. 2003; 44:3281–3284. (e) Suzuki Y, Fukuta Y, Ota S, Kamiya M, Sato M. J. Org. Chem. 2011; 76:3960–3967. [PubMed: 21469707] For intramolecular *oxa*-Michael reactions to form tetrahydroxanthenes, see: For xanthone formation, see:
- (30). See Supporting Information for details on characterization.

- (31). (a) Hendrickson JB, Ramsay MVJ, Kelly TR. J. Am. Chem. Soc. 1972; 94:6834–6843. (b) Lewis JR, Paul JG. J. Chem. Soc., Perkin Trans 1. 1981:770–775. (c) Adeboya MO, Edwards RL, Lassoe T, Maitland DJ, Shields S, Whalley AJS. J. Chem. Soc. Perkin Trans. 1996:1419–1426.
- (32). (a) Andrus MB, Hicken EJ, Meredith EL, Simmons BL, Cannon JF. Org. Lett. 2003; 5:3859–3862. [PubMed: 14535728] (b) Gao W, He Z, Qian Y, Zhao J, Huang Y. Chem. Sci. 2012; 3:883–886.
- (33). Baldwin JE. J. Chem. Soc. Chem. Commun. 1976:734.
- (34). Nichols AL, Zhang P, Martin SF. Org. Lett. 2011; 13:4696–4699. [PubMed: 21812455] For examples of cyclization of hindered 2-hydroxybenzoyl-*p*-quinones to xanthenes, see:

Scheme 1.
Synthesis of aryl nitrile **6**

Scheme 2.
Synthesis of sydownin B *via* an aryl nitrile

Scheme 3.
Alternative route to graphisin A

Scheme 4.
Dihydroisobenzofuran formation

Scheme 5.
Synthesis of sydownin B *via* graphisin A

Scheme 6.
Oxidation of graphisin A

Figure 1.
Acremoxanthone A and related natural products

Figure 2.
Initial synthetic plan

Figure 3.
DFT minimized model of nitrile **13**

Figure 4.
DFT minimized model of proposed benzoquinone intermediate **21**