Clinical update

The role of pulse oximetry in chiropractic practice: a rationale for its use

Michael W. Hall DC, FIACN^{a,*},
Anne M. Jensen DC, ICSSD, MSc (Social Science), MSc (Psych)^b

^a Professor, Parker University, Dallas, TX

^b PhD Candidate, Departments of Primary Care and Continuing Professional Education, University of Oxford, UK

Received 8 January 2011; received in revised form 20 August 2011; accepted 20 October 2011

Key indexing terms:

Oximetry;
Diagnostic techniques
and procedures;
Chiropractic

Abstract

Objective: Pulse oximetry is used regularly to assess oxygen saturation levels. The objective of this commentary is to discuss a rationale for using pulse oximetry in chiropractic practice.

Discussion: Pulse oximetry may offer doctors of chiropractic a way to monitor patients' oxygen saturation levels. Quantification of saturation values with heart rate may give clinical aid to the management of chiropractic patients. Markedly reduced saturation levels may necessitate medical referral, whereas mildly reduced levels could lead to changes in chiropractic management.

Conclusions: Pulse oximetry has the potential to be an integral part of chiropractic practice.

© 2012 National University of Health Sciences.

Introduction

Pulse oximetry is a standard measure of peripheral arterial oxygen saturation in many health care venues. It has become a routine assessment in critical care units,¹ labor and delivery units,² paediatrics,³ dentistry,⁴ and even veterinary medicine.^{5,6} Some professionals believe it is so important that it has been called the *fifth vital sign*.⁷⁻¹¹ The prevalence of undersaturation in the general population has yet to be determined, but is estimated to be approximately 30% in those presenting to hospital emergency departments.^{7,8}

A pulse oximeter is a simple, inexpensive, and noninvasive method for quickly measuring a patient's heart rate and oxygen saturation values. It is used to detect hypoxemia¹ or deficient oxygenation in the blood.¹² More specifically, it measures the percentage of hemoglobin bound to oxygen in arterial blood, denoted as SpO_2 .¹ Hypoxemia often leads to hypoxia (ie, reduced tissue oxygenation¹³), which results in oxidative stress, a condition associated with a wide variety of degenerative processes.¹⁴

Doctors of chiropractic (DCs) are trained to assess vital signs and the clinical signs of oxidative stress, such as cyanosis, pallor, fatigue, and mental confusion. However, previous studies have found that clinicians using these signs alone failed to detect hypoxemia/hypoxia in emergency department patients.^{7,8} Furthermore, it has been shown that the absence of these positive findings does not eliminate the possibility of

* Corresponding author. Parker University, 2500 Walnut Hill Lane, Dallas, TX 75229. Tel.: +1 972 438 6932; fax: +1 214 902 2438.

E-mail address: MHall@parker.edu (M. W. Hall).

serious cardiopulmonary or cerebrovascular disease.^{7,8,15} Pulse oximetry, which has been shown to accurately and reliably detect undersaturation,¹⁶ may improve DCs' ability to assess patients' risk of potentially life-compromising illnesses. The purpose of this article is to provide a rationale supporting the regular use of pulse oximetry in chiropractic practice.

Conceptual model of pulse oximetry in a chiropractic setting

Pulse oximetry is used regularly in acute care settings, such as emergency departments and critical care units. Although there are not much data to date on its use in outpatient settings, it is becoming more widely used in such settings as general practices,¹⁷⁻¹⁹ pediatrics,²⁰⁻²² dentistry,^{23,24} sport science,²⁵ acupuncture,²⁶ and even at home.^{21,27} The authors believe that there are strong arguments for its use within the chiropractic clinical practice.

The principal reasons that pulse oximetry should be used in chiropractic practice include the following:

1. Pulse oximetry allows for quick and reliable retrieval of oxygen saturation and heart rate values previously not readily obtained or used.
2. Pulse oximetry allows for immediate assessment of potentially hypoxic patients.
3. Pulse oximetry allows for important changes in patient management potentially influencing outcomes and safety.

The use of an oximeter in a chiropractic practice affords the practitioner a quick capture of both heart rate and oxygen saturation. Heart rate varies with many aspects of the clinical examination process, such as temperature, anxiety, pain, and altered sympathetic/parasympathetic regulation.²⁸ If not for an oximeter, oxygen saturation altogether would not be captured during the routine chiropractic examination.

Use of palpation of peripheral pulses to check for patency of peripheral arteries has been shown to be of questionable accuracy.²⁹⁻³² A more reliable method to assess cardiovascular health is warranted. It has been recommended that patients at risk for hypoxic conditions be monitored using pulse oximetry.^{12,14,33-37} Hypoxia is commonly found in the elderly³⁸ and those at risk for coronary vascular disease,³⁹ coronary artery disease,³⁹ sleep disordered breathing,³⁹ disorders of balance, hypertension,³⁹ chronic migraines, and anx-

iety and mood disorders,³⁹ all of which can be present within a population of chiropractic patients. See [Table 1](#) for a complete list of clinical signs and symptoms indicating use of pulse oximetry.

How it works

Pulse oximetry depends upon the optical characteristics of pulsating arterial blood.¹ Using the Beer-Lambert law on the absorption of light, the pulse oximeter has a pair of small light-emitting diodes that are placed on a translucent part of the patient's body, such as an earlobe or fingertip,⁵⁷ and left in place for 5 minutes. One light-emitting diode emits red light and the other emits infrared light, which are absorbed at significantly different rates by oxygenated hemoglobin compared with deoxygenated hemoglobin.¹ With each heart beat, the change in blood volume causes pulsating changes in the amounts of red and infrared light absorbed.¹ From this information, SpO₂ can be calculated using an algorithm programmed into the device. Most pulse oximeters display both the SpO₂ (in percentage saturation) and heart rate (in beats per minute) ([Fig 1](#)). It must be noted that detection of a pulse is imperative for the operation of a pulse oximeter¹; therefore, attention must be given to the correct placement and use of the oximeter for its results to be meaningful.

Pulse oximetry norms have been established for SpO₂, with values greater than 95% considered normal.¹ For most devices, accuracy to within 2% has become an acceptable standard; and most are considered sufficiently accurate and reliable for many clinical purposes.^{14,16,58-60}

Interpreting results

Ideally, an oxygen saturation value of 100% or near 100% saturation is preferred for optimum efficiency. However, daytime SpO₂ values greater than 95% are considered normal by many.^{7,8,57,61} Conversely, daytime values less than 90% are cause for alarm and immediate attention.^{12,13,33,34} Because SpO₂ values may fall 4% to 6% or more at night, daytime values of 90% to 95% may mean nighttime values less than 90%, which would also be cause for concern.^{13,34} Chiropractors that detect daytime saturation values less than 97% should monitor this patient's SpO₂ each

Table 1 Signs and symptoms indicating use of pulse oximetry

History
<i>Inadequate external respiration</i>
Decreased O ₂ in environment
High altitudes ⁴⁰
Enclosures without outside ventilation ⁴¹
Smoke inhalation ⁴¹
Toxic gas inhalation ⁴¹
<i>Inadequate mechanical ventilation</i>
Pain ^{26,42}
Emphysema ⁴²
Pleurisy ⁴³
Sleep disordered breathing (eg, apnea) ⁴⁴
<i>Traumatic injuries</i>
Rib fractures ^{41,42}
Pneumothorax ⁴¹
Flail chest ⁴⁵
Crushing neck or chest injuries ^{41,42}
<i>Other conditions</i>
Airway obstruction ⁴⁶
Anxiety ^{42,47}
Depression ⁴⁸
Chronic migraines ⁴⁹
Chronic illness ⁵⁰⁻⁵²
<i>Inadequate oxygen diffusion</i>
Pulmonary edema ⁴¹
Pneumonia ⁵³
COPD ⁵⁴
Pulmonary emboli ⁴¹
<i>Inadequate oxygen transport</i>
Anemia
Inadequate hemoglobin ⁵⁵
Inadequate # RBCs ⁵⁵
Poisoning
Carbon monoxide poisoning ⁴¹
<i>Inadequate circulation</i>
Shock ⁴¹
Arteriosclerosis
Hypothermia ⁴¹
<i>Physical examination</i>
<i>Signs and symptoms of respiratory compromise</i>
Dyspnea ⁵⁵
Accessory muscle use ⁴²
Inability to speak in full sentences
Adventitious breath sounds ⁵⁵
Irregular breathing pattern ⁵⁵
Abdominal breathing only ⁴²
Increased or decreased respiratory rate ⁵⁵
Shallow breathing ⁵⁵
Flared nostrils or pursed lips
Unusual anatomy (eg, barrel chest) ⁵⁵

Table 1 (continued)

History
<i>Signs and symptoms of hypoxia⁵⁵</i>
Restlessness
Altered or deteriorating mental status
Increased heart rate
Increased or decreased respiratory rate
Cyanosis
<i>Other signs or symptoms</i>
Hypertension ⁵⁶

Critical signs and symptoms indicating use of pulse oximetry. COPD, chronic obstructive pulmonary disease; RBC, red blood cell.

visit, whereas values less than 90% should prompt immediate referral for medical assessment.

The pulse oximeter also displays the patient's heart rate and, when correlated with oxygen saturation level, can round out the clinical picture. See Table 2 for a suggested action plan for interpreting results. Other factors can also influence oxygen saturation levels; therefore, the chiropractic physician must bear this in mind when designing a patient management plan.

Furthermore, it must be noted that there is no evidence to date to suggest that chiropractic treatments can influence the outcome of pulse oximetry assessments. However, no evidence does not mean that there

Fig 1. Pulse oximeter—placement on right index finger.

Table 2 Suggested interpretation of SpO₂ values

Daytime SpO ₂ (%) ³⁴	Heart Rate	Interpretation	Possible Risks	Suggested Interventions
>97	Normal	Good health markers ¹		Education/awareness Chiropractic adjustments (spine, ribs, chest) ⁶² Monitor periodically ¹
	Elevated	Autonomic arousal ^{47,63}	Tension Stress Anxiety Fear	Education/awareness Chiropractic adjustments (spine, ribs, chest) ⁶² Monitor periodically ¹ Lifestyle/exercise/stress reduction ⁶³
	Decreased	Training induced ^{64,65}	Athletic heart syndrome	Education/awareness Chiropractic adjustments (spine, ribs, chest) ⁶² Monitor periodically ¹
94-97	Normal, increased, or decrease	Early stages of lowered oxygenation ^{1,11}	Fatigue Malaise Migraine Sleep disturbance	Education/awareness Chiropractic adjustments (spine, ribs, chest) ⁶² Monitor each visit ¹ Breathing exercises ⁶⁶ Lifestyle/exercise/stress reduction ⁶³ Possibly refer for medical assessment ¹¹
<93	Normal or increase	Cardiac dysfunction ^{39,67}	TIA Stroke Neurological compromise Dementia CAD CVD	Education/awareness Interprofessional communication/referral ¹¹ Prescription oxygen ¹⁷ Ventilation assistance ¹⁷ Chiropractic adjustments (spine, ribs, chest) ⁶² Breathing exercises ⁶⁶ Lifestyle/exercise/stress reduction ⁶³ Monitor each visit ¹
	Decreased	Autonomic abnormality ²⁸	All of the above	Immediate referral ¹¹

SpO₂, oxygen saturation by pulse oximetry; TIA, transient ischemic attack; CAD, coronary artery disease; CVD, cardiovascular disease.

is no benefit; it simply means there is no evidence to date. Research in this domain is warranted.

Limitations and suggested research

Although pulse oximetry may be a useful diagnostic tool, it does not offer a comprehensive description of the complete oxygen transport system. Pulse oximetry is one part of the thorough assessment of a patient's oxygenation status, which should also include assessment of oxygen delivery and perfusion systems, and the patient's ability to ventilate.⁵⁷ Specific physiologic factors may have an influence

on the accuracy of the SpO₂ values obtained and, therefore, must be considered.

The oxygen-carrying capacity of blood may be hindered by certain disease states, such as anemia⁶⁸ or reduced or abnormal hemoglobin.^{1,57} In addition, increased pH or decreased body temperature may increase hemoglobin's oxygen affinity, thereby making oxygen less available to tissues.⁵⁷ Low perfusion states will also result in inaccurate SpO₂ readings due to low signal of the pulsating flow of blood.¹ Therefore, caution is recommended with conditions such as peripheral neuropathies⁵⁷ or peripheral artery disease.⁶⁹ Error may also be introduced with tremor or excessive movement because of signal loss, which may result in an underestimation of SpO₂.^{1,57}

Sensor malpositioning can also result in inaccuracies. False-positive readings are common if the sensors detect excessive ambient light, such as from sunlight, fluorescent lights, infrared heating lights, or examination lights.¹ In adults, fingertip probes have been proven to be more accurate than earlobe probes,¹ whereas in children, ear probes were found to be more accurate than finger or toe probes.³ Any discoloration of the nail bed or a dark nail polish may affect transmission of the light, resulting in an underestimation of SpO₂.^{1,57} On the other hand, darkly pigmented skin does not seem to impact pulse oximetry whereas severe jaundice or hyperbilirubinemia might have an impact if SpO₂ is less than 90%.¹

The use of pulse oximetry may also be misleading in patients whose ability to ventilate is compromised, such as those with chronic obstructive pulmonary disease^{54,70} or acute asthma.^{71,72} Although it may be helpful to monitor an acute exacerbation of these conditions, additional assessment must also be undertaken.^{54,70-72} In addition to these physiological limitations, pulse oximetry also has technological limitations. As with any measuring device, the instrument is only as accurate as its empirical calibration. Despite this limitation, if using the same instrument to compare the same patient at different points in time, the degree of error can be minimized.¹ Error is especially minimized if saturation readings are greater than 80%.¹

Because pulse oximetry has some accuracy limitations, practitioners must understand them fully to ensure effective use. Within these constraints, pulse oximetry can offer a valuable, noninvasive measure of arterial saturation.

Finally, because this article is a commentary, it does not offer evidence of the usefulness of pulse oximetry in chiropractic clinical practice. The purpose of this article is to propose a rationale for the use of this technology to assess patients presenting to DCs. Investigational studies are needed to establish if pulse oximetry has a place in a chiropractic setting. Future research should focus on the prevalence of under-saturation within chiropractic settings and on implementing pulse oximetry as an outcome measure to investigate chiropractic's therapeutic effect.

Conclusion

As primary health care providers, DCs should be vigilant in detecting signs of oxidative stress or

hypoxemia in their patients. It is the authors' belief that the utilization of pulse oximetry should be considered a standard part of the clinical examination in the chiropractic office. Oxygen saturation values are important determinants of health and the status of peripheral tissues. Quantification of saturation values with heart rate may give clinical aid to the management of the chiropractic patient. When used appropriately, pulse oximetry may offer DCs a safe and effective way to monitor patients' oxygen saturation levels.

Funding sources and potential conflicts of interest

No funding sources or conflicts of interest were reported for this study.

References

1. Jensen LA, Onyskiw JE, Prasad NGN. Meta-analysis of arterial oxygen saturation monitoring by pulse oximetry in adults. *Heart Lung* 1998;27(6):387-408.
2. van Dijk EJ, Vermeer SE, de Groot JC, van de Minkelis J, Prins ND, Oudkerk M, et al. Arterial oxygen saturation, COPD, and cerebral small vessel disease. *J Neurol Neurosurg Psychiatry* 2004;75(5):733-6.
3. Bilan N, Behbahan AG, Abdinia B, Mahallei M. Validity of pulse oximetry in detection of hypoxaemia in children: comparison of ear, thumb and toe probe placements. *East Mediterr Health J* 2010;16(2):218-22.
4. Jafarzadeh H, Rosenberg PA. Pulse oximetry: review of a potential aid in endodontic diagnosis. *J Endod* 2009;35(3):329-33.
5. Allen JL. Pulse oximetry: everyday uses in a zoological practice. *Vet Rec* 1992;130(16):354-5.
6. Koenig J, McDonnell W, Valverde A. Accuracy of pulse oximetry and capnography in healthy and compromised horses during spontaneous and controlled ventilation. *Can J Vet Res* 2003;67(3):169-74.
7. Mower WR, Myers G, Nicklin EL, Kearin KT, Baraff LJ, Sachs C. Pulse oximetry as a fifth vital sign in emergency geriatric assessment. *Acad Emerg Med* 1998;5(9):858-65.
8. Mower WR, Sachs C, Nicklin EL, Baraff LJ. Pulse oximetry as a fifth pediatric vital sign. *Pediatrics* 1997;99(5):681-6.
9. Neff TA. Routine oximetry: a fifth vital sign? *Chest* 1988;94(2):227.
10. Tierney LM, Whooley MA, Saint S. Oxygen saturation: a fifth vital sign? *West J Med* 1997;166(4):285-6.
11. Tozzetti C, Adembri C, Modesti PA. Pulse oximeter, the fifth vital sign: a safety belt or a prison of the mind? *Intern Emerg Med* 2009;4(4):331-2.
12. Ali K, Cheek E, Sills S, Crome P, Roffe C. Day-night differences in oxygen saturation and the frequency of

- desaturations in the first 24 hours in patients with acute stroke. *J Stroke Cerebrovasc Dis* 2007;16(6):239-44.
13. Roffe C, Sills S, Halim M, Wilde K, Allen MB, Jones PW, et al. Unexpected nocturnal hypoxia in patients with acute stroke. *Stroke* 2003;34(11):2641-5.
 14. Lee WW, Mayberry K, Crapo R, Jensen RL. The accuracy of pulse oximetry in the emergency department. *Am J Emerg Med* 2000;18(4):427-31.
 15. Maneker AJ, Petrack EM, Krug SE. Contribution of routine pulse oximetry to evaluation and management of patients with respiratory illness in a pediatric emergency department. *Ann Emerg Med* 1995;25(1):36-40.
 16. Louw A, Cracco C, Cerf C, Harf A, Duvaldestin P, Lemaire F, et al. Accuracy of pulse oximetry in the intensive care unit. *Intensive Care Med* 2001;27(10):1606-13.
 17. Roberts CM, Franklin J, O'Neill A, Roberts RPL, Ide J, Hanley ML, et al. Screening patients in general practice with COPD for long term domiciliary oxygen requirement using pulse oximetry. *Respiratory Medicine* 1998;92(11):1265-8.
 18. Potter VAJ. Pulse oximetry in general practice: how would a pulse oximeter influence patient management? *Eur J Gen Pract* 2007;13(4):216-20.
 19. Ingram G, Munro N. The use (or otherwise) of pulse oximetry in general practice. *Br J Gen Pract* 2005;55(516):501-2.
 20. Schidlow DN, Anderson JB, Klitzner TS, Beekman Iii RH, Jenkins KJ, Kugler JD, et al. Variation in interstage outpatient care after the Norwood procedure: a report from the Joint Council on Congenital Heart Disease National Quality Improvement Collaborative. *Congen Heart Dis* 2011;6(2):98-107.
 21. Felemban O, Leroux K, Aubertin G, Miandy F, Damagnez F, Amorim B, et al. Value of gas exchange recording at home in children receiving non-invasive ventilation. *Pediatr Pulmonol* 2011;46(8):802-8.
 22. Reséndiz Gómez F, Gómez Vera J, Orea Solano M. Pulse oxymetry in the diagnosis of asthma in children. *Oximetría de pulso en el diagnóstico de asma en el niño* 2004;51(3):102-6.
 23. Pozzobon MH, de Sousa Vieira R, Alves AMH, Reyes-Carmona J, Teixeira CS, de Souza BDM, et al. Assessment of pulp blood flow in primary and permanent teeth using pulse oximetry. *Dental Traumatol* 2011;27(3):184-8.
 24. Dunn-Russell T, Adair SM, Sams DR, Russell CM, Barenie JT. Oxygen saturation and diffusion hypoxia in children following nitrous oxide sedation. *Pediatr Dent* 1993;15(2):88-92.
 25. Marca Fuertes C, Canales MG, Miguel-Tobal F, Escudero PM. The pulse oximetry and its use in maximal exercise test. *La pulsioximetría y su aplicación en pruebas de esfuerzo máximo*; 2010.
 26. Ahmedov S. Pulse oximetry and acupuncture in patients with pain syndrome. *Med Acupunct* 2010;22(3):219-22.
 27. Kacmarek RM. Noninvasive monitoring of respiratory function outside of the hospital. *Respir Care* 1990;35(7):719-27.
 28. Guyton AC, Hall JE. Human physiology and mechanisms of disease. 10th ed. Philadelphia: W.B. Saunders Co.; 2000.
 29. Cournot M, Boccalon H, Cambou JP, Guilloux J, Taraszkiwicz D, Hanaire-BROUTIN H, et al. Accuracy of the screening physical examination to identify subclinical atherosclerosis and peripheral arterial disease in asymptomatic subjects. *J Vasc Surg* 2007;46(6):1215-21.
 30. Lundin M, Wiksten JP, Peräkylä T, Lindfors O, Savolainen H, Skyttä J, et al. Distal pulse palpation: is it reliable? *World J Surg* 1999;23(3):252-5.
 31. Myers KA, Scott DF, Devine TJ, Johnston AH, Denton MJ, Gilfillan IS. Palpation of the femoral and popliteal pulses: a study of the accuracy as assessed by agreement between multiple observers. *Eur J Vasc Surg* 1987;1(4):245-9.
 32. Brearley S, Shearman CP, Simms MH. Peripheral pulse palpation: an unreliable physical sign. *Ann R Coll Surg Engl* 1992;74(3):169-71.
 33. Cripps T, Rocker G, Stradling J. Nocturnal hypoxia and arrhythmias in patients with impaired left ventricular function. *Br Heart J* 1992;68(4):382-6.
 34. Eguchi K, Kario K, Hoshida S, Ishikawa J, Morinari M, Shimada K. Nocturnal hypoxia is associated with silent cerebrovascular disease in a high-risk Japanese community-dwelling population. *Am J Hypertens* 2005;18(11):1489-95.
 35. Lavie L. Oxidative stress and inflammatory consequences in obstructive sleep apnoea. *Int J Respir Care* 2006;2(1):21-4.
 36. Muraki I, Tanigawa T, Yamagishi K, Sakurai S, Ohira T, Imano H, et al. Nocturnal intermittent hypoxia and metabolic syndrome; the effect of being overweight: the CIRCS study. *J Atheroscler Thromb* 2010;17(4):369-77.
 37. Roffe C, Frohnhofen H, Sills S, Hodsoll J, Allen MB, Jones PW. Frequency of nocturnal hypoxia in clinically stable patients during stroke rehabilitation. *Clin Rehabil* 2010;24(3):267-75.
 38. Roth T, Holtmann BJ. Prevalence of nocturnal oxygen desaturation in patients admitted to a geriatric rehabilitation unit. *J Sleep Res* 1995;4(1):194-6.
 39. Sommermeier D, Schwaibold M, Schöller B, Grote L, Hedner J, Bolz A. Prediction of cardiovascular risk from peripheral pulse wave. IFMBE proceedings. Munich, Germany: Springer; 2009. p. 891-2.
 40. Roach RC, Hackett PH. Frontiers of hypoxia research: acute mountain sickness. *J Exp Biol* 2001;204(18):3161-70.
 41. Bledsoe BE, Porter RS, Cherry RA. Essentials of paramedic care update. 2nd ed. Upper Saddle River, NJ: Prentice Hall, 2003.
 42. Braun SR. Respiratory rate and pattern. In: Walker HK, Hall WD, Hurst JW, editors. Clinical methods: the history, physical, and laboratory examinations. 3rd ed. Boston: Butterworths; 1990. p. 226-30.
 43. Yim JY, Lee KH, Kyung JH, Chang JH, Cheon SH. A study of effect on pulmonary function of pleural effusion in tuberculous pleurisy patients. *Tuberc Respir Dis* 1996;43(4):491-9.
 44. Chen NH, Chen MC, Li HY, Chen CW, Wang PC. A two-tier screening model using quality-of-life measures and pulse oximetry to screen adults with sleep-disordered breathing. *Sleep Breathing* 2010;1-8.
 45. Bastos R, Calhoun JH, Baisden CE. Flail chest and pulmonary contusion. *Semin Thorac Cardiovasc Surg* 2008;20(1):39-45.
 46. Regal MAH, El Ghoneimy YA, Aljehani YM. Respiratory distress in patients with central airway obstruction. *Indian J Thorac Cardiovasc Surg* 2010;1-6.
 47. Hoehn-Saric R. Anxiety and somatization. *Int Congr Ser* 2006;368-72.
 48. Cyranowski JM, Hofkens TL, Swartz HA, Salomon K, Gianaros PJ. Cardiac vagal control in nonmedicated depressed women and nondepressed controls: impact of depression status, lifetime trauma history, and respiratory factors. *Psychosom Med* 2011;73(4):336-43.

49. Post MC, Van Gent MWF, Plokker HWM, Westermann CJJ, Kelder JC, Mager JJ, et al. Pulmonary arteriovenous malformations associated with migraine with aura. *Eur Respir J* 2009;34(4):882-7.
50. Morden NE, Sullivan SD, Bartle B, Lee TA. Skeletal health in men with chronic lung disease: rates of testing, treatment, and fractures. *Osteoporos Int* 2011;22(6):1855-62.
51. Sykes S, Kingshott R, Primhak R. Awake and asleep oxygen saturations in infants with chronic neonatal lung disease. *Acta Paediatr Int J Paediatr* 2011;100(8):1087-91.
52. Doering LV. The effect of positioning on hemodynamics and gas exchange in the critically ill: a review. *Am J Crit Care* 1993; 2(3):208-16.
53. Foran M, Ahn R, Novik J, Tyer-Viola L, Chilufya K, Katamba K, et al. Prevalence of undiagnosed hypoxemia in adults and children in an under-resourced district hospital in Zambia. *Int J Emerg Med* 2010;3(4):351-6.
54. Schermer T, Leenders J, in't veen H, van den Bosch W, Wissink A, Smeele I, et al. Pulse oximetry in family practice: indications and clinical observations in patients with COPD. *Fam Pract* 2009;26(6):524-31.
55. Bickley LS. Bates' guide to physical examination and history taking. 9th ed. Philadelphia: Lippincott Williams & Wilkins; 2005.
56. Tanigawa T. Obstructive sleep apnea: its prevention and screening may contribute to the prevention of hypertension, diabetes and cardiovascular diseases. *EPMA J* 2011;2(1):83-9.
57. Schutz SL. Oxygen saturation monitoring by pulse oximetry. In: Lynn-McHale DJ, Carlson KK, editors. *AACN procedure manual for critical care*. 4th ed. Philadelphia: W. B. Saunders; 2001.
58. Kamlin COF, Dawson JA, O'Donnell CPF, Morley CJ, Donath SM, Sekhon J, et al. Accuracy of pulse oximetry measurement of heart rate of newborn infants in the delivery room. *J Pediatr* 2008;152(6):756-60.
59. Muñoz X, Torres F, Sampol G, Rios J, Martí S, Escrich E. Accuracy and reliability of pulse oximetry at different arterial carbon dioxide pressure levels. *Eur Respir J* 2008;32(4): 1053-9.
60. Wilson BJ, Cowan HJ, Lord JA, Zuege DJ, Zygun DA. The accuracy of pulse oximetry in emergency department patients with severe sepsis and septic shock: a retrospective cohort study. *BMC Emerg Med* 2009;10:9.
61. Ogburn-Russell L, Johnson JE. Oxygen saturation levels in the well elderly: altitude makes a difference. *J Gerontol Nurs* 1990;16(10):26-30.
62. Budgell BS. Reflex effects of subluxation: the autonomic nervous system. *J Manipulative Physiol Ther* 2000;23(2):104-6.
63. Hoehn-Saric R. Physiologic responses in anxiety. *Curr Psychiatry Rev* 2007;3(3):196-204.
64. Somanathan L, Khalil I. Fitness monitoring system based on heart rate and SpO2 level; 2010.
65. Puffer JC. The athletic heart syndrome: ruling out cardiac pathologies. *Phys Sports Med* 2002;30(7):41-7.
66. Urell C, Emtner M, Hedenström H, Tenling A, Breidenskog M, Westerdaal E. Deep breathing exercises with positive expiratory pressure at a higher rate improve oxygenation in the early period after cardiac surgery—a randomised controlled trial. *Eur J Cardiothorac Surg* 2011;40(1):162-7.
67. Hayashi M, Fujimoto K, Urushibata K, Uchikawa SI, Imamura H, Kubo K. Nocturnal oxygen desaturation correlates with the severity of coronary atherosclerosis in coronary artery disease. *Chest* 2003;124(3):936-41.
68. Severinghaus JW, Koh SO. Effect of anemia on pulse oximeter accuracy at low saturation. *J Clin Monit* 1990;6(2):85-8.
69. Joyce WP, Walsh K, Gough DB, Gorey TF, Fitzpatrick JM. Pulse oximetry: a new non-invasive assessment of peripheral arterial occlusive disease. *Br J Surg* 1990;77(10):1115-7.
70. McGovern JP, Sasse SA, Stansbury DW, Causing LA, Light RW. Comparison of oxygen saturation by pulse oximetry and co-oximetry during exercise testing in patients with COPD. *Chest* 1996;109(5):1151-5.
71. Ribeiro de Andrade C, Duarte MC, Camargos P. Correlations between pulse oximetry and peak expiratory flow in acute asthma. *Braz J Med Biol Res* 2007;40(4):485-90.
72. Solé D, Komatsu MK, Carvalho KVT, Naspitz CK. Pulse oximetry in the evaluation of the severity of acute asthma and/or wheezing in children. *J Asthma* 1999;36(4):327-33.