Published in final edited form as:

Biol Psychiatry. 2012 July 1; 72(1): 8–18. doi:10.1016/j.biopsych.2012.01.012.

Defensive mobilization in specific phobia: Fear specificity, negative affectivity and diagnostic prominence

Lisa M. McTeague, Ph.D., Peter J. Lang, Ph.D., Bethany C. Wangelin, M.S., Marie-Claude Laplante, Ph.D., and Margaret M. Bradley, Ph.D.

University of Florida

Abstract

Background—Understanding of exaggerated responsivity in specific phobia—its physiology and neural mediators—has advanced considerably. However, despite strong phenotypic evidence that prominence of specific phobia relative to co-occurring conditions (i.e., principal versus non-principal disorder) is associated with dramatic differences in subjective distress, there is yet no consideration of such comorbidity issues on objective defensive reactivity.

Methods—A community sample of specific phobia (N=74 principal phobia; N=86 non-principal phobia) and control (n=76) participants imagined threatening and neutral events while acoustic startle probes were presented and eye-blink responses (orbicularis oculi) recorded. Changes in heart rate, skin conductance level, and facial expressivity were also measured.

Results—Principal specific phobia patients far exceeded controls in startle reflex and autonomic reactivity during imagery of idiographic fear scenes. Distinguishing between single and multiple phobias within principal phobia and comparing these to non-principal phobia revealed a continuum of decreasing defensive mobilization: single phobia patients were strongly reactive, multiple phobia intermediate, and non-principal patients reliably attenuated—the inverse of measures of pervasive anxiety and dysphoria (i.e., negative affectivity). Further, as more disorders supplanted specific phobia from principal disorder, overall defensive mobilization was systematically more impaired.

Conclusions—The exaggerated responsivity considered characteristic of specific phobia is limited to those patients for whom circumscribed fear is the most impairing condition, and coincident with little additional affective psychopathology. As specific phobia is superseded in severity by broad and chronic negative affectivity, defensive reactivity progressively diminishes. Focal fears may still be clinically-significant, but not reflected in objective measures of defensive mobilization.

Keywords

imagery; anxiety; specific phobia; fear; comorbidity; depression; anhedonia; anxiety sensitivity; chronicity; emotional reactivity; narrative imagery; mental imagery; diagnostic subtypes;

© 2012 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved

Corresponding author: Peter J. Lang, PhD Center for the Study of Emotion & Attention University of Florida PO Box 112766 Gainesville, FL 32611 plang@phhp.ufl.edu Telephone: (352) 392-2439 Fax: (352) 392-6047.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Financial Disclosures All authors report no biomedical financial interests or potential conflicts of interest.

psychophysiology; startle; heart rate; facial expressivity; skin conductance; corrugator; EMG; SCL

Introduction

Specific phobia is considered the prototypical anxiety disorder of defensive hyperreactivity, a view supported by extensive evidence of pronounced mobilization to fear cues—in reflex psychophysiology (1–5), electro-cortical response (6–7), neural circuitry activation (8–11)—from a variety of elicitation procedures (e.g., pictures (2), movies (12), imagery (3), and conditioning (8)). Surprisingly, however, this literature does not include examination of the broader clinical constellation within which specific phobias are often embedded. Much of the research is based on non-patient samples, and the minority of physiological investigations addressing clinically-significant specific phobia have assessed presence or absence of the disorder irrespective of its severity relative to co-occurring conditions. As such, there is a dearth of research utilizing objective measures of emotional responding that considers critical features such as diagnostic primacy/prominence and comorbidity. In phenotypic studies careful consideration of ranked severity (i.e., principal versus non-principal problem) across anxiety disorders has revealed important variation in liability for co-occurring disorders. For example, in a sample of over 1,000 anxiety patients Brown and colleagues observed that as a whole, 70% of patients with a specific phobia diagnosis had a comorbid anxiety or mood disorder (13). The comorbidity rate dropped to 33% among the subset of patients for whom specific phobia was the principal disorder. In other words, whether specific phobia is the principal (i.e., worst) disorder or not, covaries dramatically with the level of functional interference and, perhaps, the burden that affective psychopathology imposes on defensive reflex physiology.

Narrative Imagery

The current investigation of specific phobia examines whether the presence/absence as well as number of fears and gradations in diagnostic primacy (i.e., principal, secondary, tertiary) reflect differences in defensive reflex physiology during narrative imagery. Script-driven emotional imagery is a valuable tool in studies of anxiety disorders, permitting presentation of both standard and idiographic threat challenges, akin to methods of imaginal exposure therapy (14). Physiological arousal during aversive imagery parallels anticipatory reactions to threatening events (15), similarly mobilizing the autonomic nervous system (e.g., heart rate, skin conductance), communicating threat through facial musculature (e.g., corrugator “frown” muscle), and prompting somatic reflexive action (e.g., startle potentiation) (16–17). Animals confronting survival threat show similar reactions, mediated by the brain's defense circuit (centered on the amygdala) (18–19) and neuroimaging studies suggest a comparable circuit (20–22) underlies human fear.

In a series of imagery investigations, Lang and colleagues (23–33) assessed differences in defensive arousal within several anxiety disorders. In general, each principal diagnostic group (e.g., social phobia, posttraumatic stress disorder (PTSD), panic disorder) showed greater defensive reactivity than control participants. However, consideration of within-diagnosis features revealed dramatic differences in defensive mobilization. That is, reactivity was robust in patients with focal affective disruptions (e.g., social phobia limited to structured performance situations), whereas reactivity was increasingly reduced as the principal disorder features were more generalized (e.g., apprehension extending to routine social interaction) and coincident with increased disorder severity and duration, poorer prognosis, and higher anxiety and depressive disorder comorbidity. Blunted reflex responding was also related to symptom elevations across numerous domains including anhedonia, unspecified/trait anxiety, anger, and functional interference. The confluence of

dimensional and categorical dysphoria was termed *negative affectivity* (27–29) to highlight the synergy of multiple pathologies as opposed to isolated disorders in modulating defensive reflex physiology. Taken together, these findings suggest that defensive engagement during imagery might be compromised by prolonged and diffuse anxious hyperarousal and accompanying negative affectivity (30–33).

The Research Problem

In the current study, a similar distress-related reflex pattern was expected within specific phobia—varying as a function of phobia precedence and comorbid symptomatology. First, principal specific phobia patients were compared to control participants with the expectation that similar to preceding studies (2–11), principal phobia would be characterized by exaggerated defensive mobilization (i.e., potentiated startle and autonomic action) during imagery of idiographic fear narratives, whereas patients and controls would react similarly during threatening imagery for which defensive mobilization is normal and adaptive (e.g., facing an attacking animal).

Next, principal phobia patients were distinguished according to whether they endorsed a single or multiple specific phobias. Further, another set of patients who had at least one specific phobia exclusive of their principal problem (i.e., non-principal/additional phobia) were identified. Concerning number of fears within principal phobia, competing hypotheses were tested: As shown in a non-patient investigation of individuals endorsing solitary or numerous fears (25), multiple phobia patients might be putatively more fearful than individuals with an isolated phobia, and hence show the most robust physiological reactivity during aversive imagery. Alternatively, in a clinical sample greater negative affectivity could be expected with multiple fears and correspondingly, reduced reactivity. These debilitating symptom features might be yet more extreme in the non-principal phobia group whose foremost difficulties could include far more generalized anxiety and dysphoria (e.g., generalized anxiety disorder (GAD), panic disorder with agoraphobia (PDA)), thus prompting the most pronounced attenuation of defensive action.

Method

Participants

Participants were assessed at the University of Florida Fear and Anxiety Disorders Clinic: 160 treatment-seeking adults with a diagnosis of specific phobia (N=74 with principal specific phobia; N=86 with non-principal/additional specific phobia¹) and 76 healthy community controls. Fear focus was distributed as follows: animal 19.4%, blood-injury-injection 15.6%, situational 40%, natural environment 18.1%, other 6.9%.

Diagnostic Classification

Diagnostic groups were established using the Anxiety Disorder Interview Schedule for DSM-IV (ADIS-IV; 34), a structured interview for assessing current anxiety, mood, substance use, and somatoform disorders and for screening psychosis and major physical disease. For multiple Axis I disorders, diagnostic primacy was determined by clinician-rated severity (ranging from 0, *No features present*, to 5, *Diagnosis present; severe*) reflecting both distress and interference. Controls denied current or lifetime diagnoses of psychiatric

¹Among the non-principal specific phobia group, principal disorders were limited to anxiety, mood, and adjustment disorders: GAD 24.4%; PDA; 19.8%; PTSD secondary to repeated trauma 14.0%; generalized social phobia 12.8%; obsessive compulsive disorder 9.3%; panic disorder without agoraphobia 8.1%; PTSD secondary to single trauma 4.7%; anxiety disorder not otherwise specified (NOS) 4.0%; adjustment disorder with mixed anxiety and depression 1.2%; recurrent major depressive disorder 1.2%; depressive disorder NOS 1.2%; circumscribed performance phobia 1.2%.

illness. Inter-rater reliability (via videotape) was calculated for 20% of patients, yielding agreement at 100% for principal and 82.35% for non-principal phobia diagnosis among three masters- or doctoral-level clinicians.

Patients whose foremost clinical complaint was specific phobia (i.e., principal phobia) were further classified according to whether the patient indicated a single phobia (N=50) or multiple phobias (N=24)².

Procedure

The University of Florida Institutional Review Board (IRB-01) approved the study. Participants provided informed consent, completed questionnaires and interview in the morning; psychophysiological assessment and clinical debriefing followed in the afternoon.

Experimental Stimuli—Twenty-four narrative imagery texts were used (35). Analyses focused on two idiographic, “personal” threat narratives representing each participant’s primary clinical fear; or for controls their “worst fear” experiences. Standard scenes included: two panic attack (crowded checkout line, driving alone), four survival threat (physical attack by animal/human), two neutral (watching documentary, reading magazine) events. Filler scripts were low arousal or engaging pleasant scenes to impede an overall unpleasant arousal context. Scripts were ~20 words designed to quickly reveal affect and reflect active participation. A woman recorded the scenes using minimal prosody for presentation over earphones (Telephonics TDH-49, Telephonics Corporation, Huntington, New York).

Imagery Assessment—Seated in a quiet, dimly lit room, with electrodes placed, participants were instructed to listen to the auditory scripts with eyes closed, vividly imagining the events described, as if actively involved. Throughout the recording session, soft tones cued participants to relax, breathe slowly, and silently repeat the word “one” to stabilize between-trial physiological activity (36). Imagery scripts were interspersed every 36 seconds in the tone series, with content pseudorandomized so that no more than 2 stimuli of the same hedonic valence (pleasant, neutral, unpleasant), or content category (e.g., panic attack) were presented consecutively. The script series was repeated in a counterbalanced order.

Trials consisted of a 1-second baseline, a 6-second auditory script, and 12 seconds of imagery. Startle probes (50-ms 95 dB(A) white noise, instantaneous rise-time) were presented at 4–5.5 or 10–11.5 seconds post-script onset, or both, and on 25% of inter-trial intervals, at 22–23.5 seconds post-imagery offset.

Following imagery assessment (approximately 45 minutes) participants rated each scene for experienced pleasure and emotional arousal (37).

Experimental Control and Data Collection

A computer running VPM software (38) controlled stimulus presentation and data acquisition. Bioamplifiers recorded electromyograph (EMG) potentials at left orbicularis

²The vast majority (78%) of principal specific phobia patients were diagnosed only with specific phobia(s). More specifically, 82% of the single, principal phobia patients endorsed no other disorders, while the remaining 18% were diagnosed with a range of disorders, typically anxiety, adjustment, and/or mood syndromes, of lesser intensity than the specific phobia. Among the principal phobia patients with multiple disorder-level phobias, 62.5% met criteria for two phobias and no other disorders and 8% endorsed three phobias and no other disorders. The remaining patients endorsed two (16.7%) or three (12.5%) specific phobias in addition to a range of less severe Axis I disorders.

occuli and corrugator supercilii, skin conductance level (SCL) and electrocardiogram (EKG) as reported previously (26).

Data Reduction and Analysis

Univariate ANOVAs and Tukey Honestly Significant Difference (HSD) tests for planned comparisons determined group differences in demographic and questionnaire data.

Using VPM software EMG, SCL [$\log[SCL+1]$], and EKG r-r intervals [converted to beats-per-minute] were reduced into half-second bins. Responses were determined by subtracting amplitude during the 1 second prior to script presentation from averages during the 12-second imagery period.

Startle blinks from orbicularis oculi EMG represented the magnitude difference between onset and peak muscle potential (39), standardized within subject in relation to the mean and standard deviation of inter-trial probe responses (26).

Using SPSS (SPSS, Inc., Chicago, Illinois), omnibus repeated measures ANOVAs were conducted separately for each physiological measure, with diagnostic status as a between-subjects factor and imagery content as a within-subjects factor. Analyses were initially performed with control versus principal specific phobia as a between-subjects factor. Startle and autonomic reactivity during imagery have been shown to strongly covary with rated emotional arousal (30–32), thus contents were entered according to the mean linear increase in arousal reported by the patients (i.e., neutral, panic attack, survival threat, idiographic/personal threat). Significant overall group effects were followed up with between-group tests by contents to specify which imagery scenarios evoked different sensitivities in patients and controls, facilitating comparisons to preceding studies that utilized different prompts (23–26). Within-group comparisons explicated interactions.

Finally, the data for 86 non-principal phobia patients were included and analyses were repeated for all patients, this time also accounting for presence of single or multiple phobias in the principal phobia group (i.e., single phobia, multiple phobia, non-principal phobia). Wilks' lambda addressed sphericity issues (40).

Results

Principal Specific Phobia and Control Groups

Affective Judgments—Across groups rated displeasure reliably increased from neutral, to panic attack, survival threat and personal threat at the extreme, $F(3,143)=258.88$ $p<.001$ (Table 1). Controls rated personal and survival threat scenes equally aversive, *all ns*; whereas patients rated personal threat more aversive than all other contents, *all ps<.001*; Content $\times$ Diagnosis interaction, $F(3,143)=3.82$, $p<.05$; Diagnosis $F(1,145)=0.90$, *ns*. Patients rated personal threat scenes more unpleasant than controls, $p<.05^\dagger$.

Both controls and patients rated personal threat scenes most arousing followed by survival threat, panic attack and neutral scenes, Content $F(3,143)=259.67$, $p<.001$; Content $\times$ Diagnosis $F(3,143)=2.64$, $p=.05$; Diagnosis $F(1,145)=3.81$, $p=.05$. Paralleling hedonic valence ratings, patients endorsed higher arousal than controls during personal threat imagery, $p<.05^\dagger$.

[†]One-tailed test based on prediction that principal phobia patients would exceed controls during personal threat imagery.

Startle Reflex Potentiation—Blink magnitude during imagery relative to ITI acoustic startle probes (Figure 1 top panel) was consistently larger during unpleasant compared to neutral imagery, Content $F(3,131)=10.17$, $p<.001$ (all unpleasant versus neutral comparisons, $ps<.001$). Extreme responses differed by group, Diagnosis $F(1,133)=6.26$, $p<.05$; Content $\times$ Diagnosis $F(3,131)=3.35$, $p<.05$; Content $\times$ Diagnosis (Cubic Contrast) $F(1,133)=4.58$, $p<.05$: whereas controls responded with similar robustness to both survival and personal threat imagery, $F(1,67)=1.0$, ns , principal phobia patients showed the greatest reflex responding during personal threat imagery—reliably exceeding responding elicited by survival threat imagery $F(1,66)=8.05$, $p<.01$, and surpassing control responses to personal threat imagery, $F(1,133)=8.81$, $p<.01$ ³.

Autonomic and Facial Responses—Heart rate (Figure 1 bottom panel) increased above neutral during unpleasant imagery, Content $F(3,142)=21.94$, $p<.001$. Whereas both groups showed the most extreme accelerations to personal threat followed by survival threat and then panic attack imagery, patients also showed a significant decrease during neutral imagery—yielding more pronounced affective discrimination in heart rate, Content $\times$ Diagnosis $F(3,142)=4.20$, $p<.01$, Content $\times$ Diagnosis (Linear Contrast) $F(1,144)=95.35$, $p<.001$, Diagnosis $F(1,144)=0.48$, ns . Conspicuous increases to personal threat, Diagnosis $F(1,144)=10.21$, $p<.01$, as well as decreases to neutral, Diagnosis $F(1,145)=7.67$, $p<.01$, demonstrated by patients both differed reliably from control responses.

Enhanced sympathetic activation was evident in increased SCL during unpleasant relative to neutral imagery (Figure 1 middle panel), Content $F(3,140)=18.53$, $p<.001$. Modulation of SCL differed by groups, Diagnosis $F(1,142)=7.79$, $p<.01$; Content $\times$ Diagnosis $F(3,140)=5.47$, $p<.01$, Content $\times$ Diagnosis (Quadratic Contrast) $F(1,142)=7.79$, $p<.001$: patients showed progressive SCL increases from neutral, to panic, survival, and finally to personal threat, while controls showed similar magnitude increases to both panic and survival threat imagery. Similar to startle responses, patients' heightened SCL—specifically during personal threat imagery—exceeded that of control participants $F(1,142)=15.05$, $p<.001$.

Principal phobia patients and controls showed the same pattern of facial frowning, covarying strongly with their shared pattern of rated displeasure; Content, $F(3,144)=14.05$, $p<.001$, Content $\times$ Diagnosis $F(3,144)=0.32$, ns ; Diagnosis $F(1,146)=0.004$, ns .

Fear Generalization, Phobia Primacy & Defensive Reactivity

Affective Judgments—All three phobia groups endorsed a similar pattern of rated displeasure, Content $F(3,147)=204.37$, $p<.001$; Content $\times$ Diagnosis, $F(6,294)=1.37$, ns ; Diagnosis, $F(2,149)=5.27$, $p<.001$, with the exception that panic attack imagery was characterized as more aversive by non-principal than single phobia patients (Table 2).

The same sensitivity to panic attack imagery emerged in more extreme ratings of emotional arousal by the non-principal relative to the single phobia group, Content $F(3,148)=163.67$, $p<.001$; Content $\times$ Diagnosis, $F(6,296)=2.79$, $p<.05$; Content $\times$ Diagnosis Quadratic Trend, $F(2,150)=5.18$, $p<.01$; Diagnosis, $F(2,143)=1.42$, ns . Importantly, across patient groups personal threat scenes were rated as equally extreme in both unpleasantness and arousal.

³No group differences emerged for blink magnitude to intertrial startle probes or for baseline SCL, or corrugator activity, $F_s=0.12$ – 02.53 . Consistent with preceding studies (30–32) heart rate was higher for patients ($M=73.38$, $SD=9.90$) than controls ($M=65.12$, $SD=10.02$), Diagnosis $F(1,144)=25.08$, $p<.001$, and as such analyses for heart rate change were calculated on residuals secondary to removing the trial-specific baseline (1-second average prior to script onset) effects via linear regression.

Startle Reflex Potentiation—Whereas affective modulation was similar between groups, Content $F(3,141)=9.32$, $p<.001$; Content $\times$ Diagnosis $F(6,282)=1.59$, ns ; marked differences were evident in magnitude of responding during imagery relative to ITI startle probes, Diagnosis, $F(2,143)=4.30$, $p<.05$, with the single phobia group exceeding the non-principal phobia group, $p<.05$ —the multiple phobia group intermediate (Figure 2 top panel). Follow-up pairwise tests (Table 2) clarified that the overall group difference was foremost attributable to exaggerated responses in the single phobia group, specifically during personal threat imagery⁴.

Autonomic and Facial Responses—Single and multiple phobia patients showed heart rate deceleration to neutral and incremental acceleration starting with panic, survival, and finally the most pronounced acceleration during personal threat imagery. Consistent with their subjective aversion and arousal ratings, non-principal phobia patients showed the second largest heart rate increase during panic attack imagery, Content $F(3,153)=24.44$, $p<.001$; Content $\times$ Diagnosis $F(6,306)=2.31$, $p<.05$; Diagnosis, $F(2,155)=0.49$, ns . Concordant with the startle findings, the only between-group difference in response magnitude was observed for personal threat imagery due to the single group showing larger accelerations than the non-principal group—the multiple phobia group, again, intermediate (Table 2; Figure 2 bottom panel).

Reliable sympathetic reactivity was indexed in SCL change during imagery, Content $F(3,150)=20.19$, $p<.001$. Single phobia patients showed a strong linear skin conductance increase from neutral to personal threat, whereas the multiple $F(1,22)=8.19$, $p<.01$, and non-principal phobia, $F(1,83)=26.14$, $p<.001$ groups demonstrated less differentiation among the lower arousing contents, Diagnosis $F(2,152)=2.90$, $p=.06$; Content $\times$ Diagnosis $F(6,300)=3.43$, $p<.01$. Finally, the single phobia group demonstrated more pronounced reactivity than the non-principal group, specifically in response to personal threat imagery—the multiple phobia group, again, evinced intermediate reactivity (Table 2; Figure 2 middle panel).

In contrast to the concordant hyperreactivity in single phobia patients across startle, SCL, and heart rate indices—specifically during personal threat imagery—corrugator changes, Content $F(3,153)=16.70$, $p<.001$, Diagnosis, $F(2,155)=0.07$, ns ; Content $\times$ Diagnosis $F(6,306)=1.12$, ns , suggested consonant, reliable facial expressivity across patient and control groups.

Diagnostic Primacy & Comorbid Negative Affectivity

Questionnaire measures of non-specific trait anxiety (STAI), cognitive/somatic symptoms of depression (BDI) as well as anhedonia (MASQ Anhedonia), interoceptive sensitivity (ASI), trait anger (STAXI), broad fearfulness (FSS), and interference across functional domains (IIRS)⁵ reliably increased from controls at the minimum, to single phobia, multiple phobia and finally non-principal/additional phobia at the extreme (Table 3; Figure 3)⁶. The latter patients also surpassed principal phobia patients in total number of Axis I disorders and frequency of comorbid anxiety as well as mood disorders. To further specify the pattern of

⁴No differences emerged between patient groups in resting physiology and intertrial interval startle probe responses (Supplemental Table 1).

⁵Similar to other studies that endeavored to utilize the IIRS for assessing functional interference in lifestyle domains associated with mental as opposed to physical health conditions (49–50), the IIRS was worded: “How much do your mental health problems (e.g., anxiety, depression) and/or their treatment interfere with your...?”

⁶For ease of comparison with the extant literature, the BDI and STAI scores were presented in conjunction with the MASQ Anhedonia subscale in Table 3. As evident in Figure 3, the other 4 MASQ subscales (Mixed Anxiety/Depression Symptoms, General Anxiety, General Depression, Anxious Arousal) showed the same pattern across groups as the BDI and STAI and, not surprisingly, all six measures were highly intercorrelated, $rs=0.69–0.89$, $ps<.001$.

comorbidity across the phobia spectrum, subgroups were compared in terms of presence or absence of single episode versus recurrent major depression, which reflected marginal differences across subgroups for transient depression in comparison to stark differences in refractory depression, with much greater prevalence of the latter in non-principal phobia. By nature of the analyses undertaken here, all patients were positive for a significant fear disorder. To further characterize the anxiety/mood comorbidity, disorders were defined as predominantly broad distress/anxious-misery (i.e., GAD, PTSD, OCD, PDA, recurrent major depression) or not, drawing on the findings of epidemiological (51–56) as well as psychophysiological (32–33) studies. Similar to the presence of more intractable depression, the frequency of broad distress disorders was far greater in non-principal (85%) than principal phobia (6–8%). In fact, for 84% of non-principal patients, at least one anxious-misery disorder superseded specific phobia in severity. Consistent with protracted functional interference, the non-principal group reported the lowest educational attainment (Table 3). This overall pattern of broad dysphoria and impairment (indexed both dimensionally and categorically) in the non-principal phobia group was consistent with clinician estimates of poorer treatment prognosis.

Interestingly, features more pertinent to the specific phobia diagnosis did not distinguish subgroups (Table 3). Irrespective of phobia type, when queried about their particular phobic object/context during interview, patients reported experiencing equivalently severe fear and avoidance and consequent distress and functional interference. Furthermore, patients recalled experiencing disorder-level dysfunction of similar duration, on average 16 years⁷.

Regarding other factors that might mitigate defensive engagement, self-reported ability to generate vivid mental imagery was equivalent across participants (Table 3). Furthermore, medication and substance use patterns exerted no discernible influence on physiological patterns⁸.

Gradations in Specific Phobia Prominence

In light of the observed pattern of decreasing physiological reactivity during personal threat imagery across single and multiple principal phobia and finally non-principal phobia, post hoc analyses were conducted to assess whether a more refined spectrum of reactivity might be revealed if the latter group were further considered according to the relative primacy of specific phobia within the diagnostic profile. Toward this aim non-principal phobia patients were distinguished as those whose specific phobia was the second-most (N=32), third-most (N=29), or fourth (or more)-most severe disorder (N=25) and repeated-measures were performed on neutral and personal threat conditions. While subjective, autonomic, and facial expressivity measures were similarly representative of the group mean across the non-principal phobia subtypes, startle potentiation during personal threat imagery systematically decreased as the specific phobia diagnosis grew less prominent in the diagnostic profile

⁷As some patients had more than one diagnosed specific phobia, chronicity refers to the earliest onset phobia. Patient groups also did not differ in duration of “worst” phobia (M= 15.46 years, SD=13.04).

⁸Psychotropic usage mirrored broad symptom severity with single (32%) and multiple phobia (29%) reporting appreciably lower rates than non-principal phobia (55%) patients ($X^2(2) = 9.04, p < .05$). Most commonly, these medications were selective serotonin-reuptake inhibitors (single 14%; multiple 13%; non-principal 36%, $X^2(2) = 10.05, p < .05$) and/or benzodiazepines (single 19%; multiple 14%; non-principal 32%, $X^2(2) = 4.05, ns$). The effects of these and less frequently endorsed compounds (e.g., serotonin-norepinephrine reuptake inhibitors 5.5% beta blockers, 3.7%, tricyclics 1.9%) were assessed by comparing resting and imagery reactivity among the medicated and non-medicated patients. Considering either general psychotropic usage or more specific classes of drugs, no reliable effects emerged, consistent with prior psychophysiological studies of anxiety and depression (57–59). Reported usage of both prescription and over-the-counter medications for promoting physical health, as well as recreational substance use were also collected but low frequencies of endorsement precluded statistical analysis. As previously demonstrated in samples characterized by highly comorbid anxiety and depression (60), non-principal patients (26%) were significantly more likely, pairwise comparisons, $ps < .05$, than control (6.6%), and single phobia patients (6.0%) to be current smokers, with multiple phobia patients intermediate (16.7%), $X^2(3) = 16.26, p < .001$. Importantly, no physiological effects were observed comparing resting and imagery reactivity of smokers and non-smokers.

(Figure 4 top panel), Diagnostic subtype, $F(4,142)=3.88$, $p<.01$, Content $\times$ Diagnostic Subtype, $F(4,142)=3.88$, $p<.05$. Whereas responses elicited during neutral imagery did not vary across groups, $F=1.90$, *ns*, reliable differences were observed during personal threat imagery, $F(4,142)=3.92$, $p<.01$: probe reflex responses in the single group exceeded both groups for whom specific phobia was superseded by at least two other disorders ($ps<.05$). Coincident with progressive startle response diminution, measures of broad negative affectivity (ASI, MASQ subscales, BDI) systematically and reliably increased ($Fs=13.52$ – 39.51 , $ps<.001$; Figure 4 bottom panel). Furthermore, the precedence of an anxious-misery disorder over specific phobia showed a parallel increase among non-principal patients, $X^2(2)=17.34$, $p<.001$: phobia as second-most (56%), third-most (93%), or fourth (or more)-most severe disorder (100%). Finally, duration of anxious-misery showed the same pattern, $F(2,70)=3.90$, $p<.05$, with patients whose phobia was most distant from the principal disorder (years $M=19.8$; $SD=13.4$) recalling those symptoms longer than the other two groups (phobia as second-most (years $M=10.7$; $SD=11$), third-most (years $M=11.6$; $SD=13.3$)).

Discussion

Consistent with the extant conceptual and empirical literature, principal specific phobia relative to control participants demonstrated amplified defensive mobilization to their worst fears, showing greater startle potentiation, skin conductance increases, and heart rate acceleration during personal threat imagery, concordant with more extreme ratings of aversion and emotional arousal. Responses to standard panic attack and survival threat imagery were similar in patients and controls consistent with functional neuroimaging data (9,20,61) and suggesting that specific phobia does not broadly sensitize fear circuitry.

Considering the number and relative primacy of specific phobia revealed that single principal phobia patients demonstrated the most robust personal threat reactivity, greatly exceeding the non-principal phobia patients in overall responsivity. Multiple phobia patients were intermediate—not exceeding the single phobia group as in the McNeil et al. (25) analogue study, but consistent with intermediate negative affectivity—showed somewhat less reactivity. For all three patient groups their personal, “worst” threatening scenes were highest in facial expressivity (corrugator) and similarly rated most arousing and unpleasant. The observed spectrum of decreasing physiological reactivity (i.e., startle reflex, heart rate, SCL) from single to multiple and finally non-principal phobia is not surprising considering the inverse increase in dimensional and categorical negative affectivity, particularly comorbidity of refractory depression and anxious-misery disorders—a pattern previously observed within social phobia (30), PTSD (31), and panic disorder with agoraphobia (PDA) (32).

In contrast to the influence of broad distress, features essential to meeting threshold for specific phobia diagnosis (e.g., fear-related avoidance, impairment) did not distinguish subgroups, suggesting that relative hyper- and hypo-reactivity did not reflect differences in phobia intensity. Furthermore, principal (single and multiple) and non-principal phobia patients recalled phobia-related dysfunction of similarly long duration (average 16 years). This differs markedly from our results for other anxiety disorders: generalized social phobia (31), PTSD (32), PDA (33) all showed hypo-reactivity amidst more enduring and broader disorder-related distress. This difference is not wholly unforeseen: specific phobia is characterized by phasic defensive mobilization in the face of fear-relevant cues (62), which can typically be readily avoided leading to rapid dissipation of distress. In contrast, generalized social phobia, PTSD, and PDA, as well as other disorders of anxious misery are marked by persistent, uncontrollable symptom exacerbation often only moderately ameliorated by escape/avoidance. Such sustained distress may ultimately take a toll on the

defensive systems subserving chronic hyperarousal, whereby protective homeostasis follows quickly—in the event of fear resolution. In concert with this, in the current research the strongest, most punctate responding was observed in those with the most isolated or encapsulated, fear-specific psychopathology (i.e., principal single phobia). Meanwhile the least physiologically reactive patients, also most disparate from principal phobia, endorsed symptom levels on the BDI and STAI beyond the 99th percentile relative to non-clinical samples. Furthermore, among those with non-principal phobia, patients with the most impaired responding (rightmost group in Figure 4) recalled the most chronic and pervasive anxious misery.

Animal research varying stressor intensity and duration (63–67) has provided some support for the hypothesis that the stress of chronic anxiety and depression may relate to dampening of defensive reflexes. For example, rats exposed to brief and/or less threatening stress demonstrate hypervigilance and hyperarousal, whereas rats exposed to longer duration stress develop more generalized anxiety and depressive-like symptoms, including passivity and reduced movement and communication behaviors (64–67). Taken together, the cumulative, chronic stress inherent in pervasive anxiety and dysphoria and the associated functional impairments may have a debilitating effect on the integrity of the underlying fear/defense system—a consequence not produced by intermittent focal fear episodes.

Alternatively, however, these defensive and subjective response profiles may be stable, time-invariant dispositions throughout the trajectories of affective dysfunction, potentially reflecting genetic underpinnings (68–69). In a recent latent class analysis of two epidemiological samples Vaidyanathan and colleagues (70) found that specific phobia diagnoses typically occur in one of two diagnostic profiles, one marked by multiple phobic disorders (and few other disorders)—the other by a wide array of Axis I disorders concurrent with specific phobia. The former group is more akin to the highly reactive principal phobia patients in the current study, while the latter group resembles the non-reactive, non-principal phobia patients. Vaidyanathan and colleagues speculated that rather than reflecting gross severity of psychopathology (71), these classes distinguished individuals fundamentally disposed to different disorder combinations—and as our findings strongly suggest, different defensive propensities as well.

Conclusion

Simply considering the presence or absence of specific phobia within the clinical presentation provides an incomplete picture of the associated defensive mobilization. The most robust defensive responding to fear-relevant imagery is evident in those patients with the most focal fearfulness and the least overall distress—a finding paradoxical in relation to the majority of conceptual and empirical work in anxiety disorders. Conversely as specific phobia is superseded in severity by more pervasive and chronic anxious apprehension and dysphoria, defensive reactivity progressively diminishes. Essentially, the exaggerated responsivity considered characteristic of specific phobia is limited to those patients for whom their circumscribed fear is the most impairing condition, and coincident with little additional affective psychopathology. Taken together, the primacy of focal fears and the extent of comorbid negative affectivity must be considered simultaneously to capture the associated phenotypic and endophenotypic constellation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by a National Institute of Mental Health grants P50 MH 72850 to the Center for the Study of Emotion and Attention (CSEA), University of Florida, Gainesville, FL (P.J. Lang Director & Principal Investigator), R21 MH082702 to Peter J. Lang, and NRSA Research Fellowship (F31 MH069048) to Lisa M. McTeague. Special thanks to Bruce N. Cuthbert for assistance with study design and interpretation of findings. Special thanks to the following individuals for their assistance in data collection: Cyd C. Strauss, Denise M. Sloan, Eleni Dimoulas, Jose M. Soler-Baillo, Reid Scott, Greg Perlman, Joshua R. Shumen, Esther Jean-Baptiste, and Hailey W. Bulls.

References

1. Grillon C. Startle reactivity and anxiety disorders: aversive conditioning, context, and neurobiology. *Biol Psychiatry*. 2002; 52(10):958–75. 15. [PubMed: 12437937]
2. Hamm AO, Cuthbert BN, Globisch J, Vaitl D. Fear and the startle reflex: blink modulation and autonomic response patterns in animal and mutilation fearful subjects. *Psychophysiology*. 1997; 34:97–107. [PubMed: 9009813]
3. Lang PJ, Levin DN, Miller GA, Kozak MJ. Fear behavior, fear imagery, and the psychophysiology of emotion: the problem of affective response integration. *J Abnorm Psychol*. 1983; 92:276–306. [PubMed: 6619405]
4. Lang PJ, Levin DN, Miller GA, Kozak MJ. Fear behavior, fear imagery, and the psychophysiology of emotion: the problem of affective response integration. *J Abnorm Psychol*. 1983; 92:276–306. [PubMed: 6619405]
5. Sarlo M, Palomba D, Angrilli A, Stegagno L. Blood phobia and spider phobia: two specific phobias with different autonomic cardiac modulations. *Biol Psychol*. 2002; 60:91–108. [PubMed: 12270586]
6. Leutgeb V, Schäfer A, Schienle A. Late cortical positivity and cardiac responsivity in female dental phobics when exposed to phobia-relevant pictures. *Int J Psychophysiol*. 2011; 79:410–6. [PubMed: 21238507]
7. Michalowski JM, Melzig CA, Weike AI, Stockburger J, Schupp HT, Hamm AO. Brain dynamics in spider-phobic individuals exposed to phobia-relevant and other emotional stimuli. *Emotion*. 2009; 9:306–15. [PubMed: 19485608]
8. Schweckendiek J, Klucken T, Merz CJ, Tabbert K, Walter B, Ambach W, Vaitl D, Stark R. Weaving the (neuronal) web: fear learning in spider phobia. *Neuroimage*. 2011; 54:681–8. [PubMed: 20673801]
9. Lipka J, Miltner WH, Straube T. Vigilance for threat interacts with amygdala responses to subliminal threat cues in specific phobia. *Biol Psychiatry*. 2011; 70:472–8. [PubMed: 21601831]
10. Ahs F, Pissioti A, Michelgård A, Frans O, Furmark T, Appel L, Fredrikson M. Disentangling the web of fear: amygdala reactivity and functional connectivity in spider and snake phobia. *Psychiatry Res*. 2009; 172:103–8. [PubMed: 19321315]
11. Wendt J, Lotze M, Weike AI, Hosten N, Hamm AO. Brain activation and defensive response mobilization during sustained exposure to phobia-related and other affective pictures in spider phobia. *Psychophysiology*. 2008; 45:205–15. [PubMed: 17995911]
12. Ritz T, Wilhelm FH, Meuret AE, Gerlach AL, Roth WT. Airway response to emotion- and disease-specific films in asthma, blood phobia, and health. *Psychophysiology*. 2011; 4:121–35. [PubMed: 20573055]
13. Brown TA, Campbell LA, Lehman CL, Grisham JR, Mancill RB. Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. *J Abnorm Psychol*. 2001; 110:585–99. [PubMed: 11727948]
14. Foa EB, Hembree E, Rothbaum BO. *Prolonged Exposure Therapy for PTSD: Emotional Processing of Traumatic Experiences Therapist Guide (Treatments That Work)*. Oxford University Press; New York: 2007.
15. Lang PJ, Kozak MJ, Miller GA, Levin DN, McLean A Jr. Emotional imagery: Conceptual structure and pattern of somato-visceral response. *Psychophysiology*. 1980; 17:179–92. [PubMed: 7375619]

16. Miller MW, Patrick CJ, Levenston GK. Affective imagery and the startle response: probing mechanisms of modulation during pleasant scenes, personal experiences, and discrete negative emotions. *Psychophysiology*. 2002; 39:519–29. [PubMed: 12212644]
17. Vrana SR, Lang PJ. Fear imagery and the startle-probe reflex. *J Abnorm Psychol*. 1990; 99:189–97. [PubMed: 2348014]
18. Walker DL, Davis M. Role of the extended amygdala in short-duration versus sustained fear: a tribute to Dr. Lennart Heimer. *Brain Struct Funct*. 2008; 213:29–42. [PubMed: 18528706]
19. Lang PJ, Davis M. Emotion, motivation, and the brain: Reflex foundations in animal and human research. *Prog Brain Res*. 2006; 156:3–29. [PubMed: 17015072]
20. Sabatinelli D, Bradley MM, Fitzsimmons JR, Lang PJ. Parallel amygdala and inferotemporal activation reflect emotional intensity and fear relevance. *Neuroimage*. 2005; 24:1265–70. [PubMed: 15670706]
21. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biol Psychiatry*. 2003; 54:504–14. [PubMed: 12946879]
22. Costa VD, Lang PJ, Sabatinelli D, Versace F, Bradley MM. Emotional imagery: Assessing pleasure and arousal in the brain's reward circuitry. *Hum Brain Mapp*. 2010; 31:1446–57. [PubMed: 20127869]
23. Weerts TC, Lang PJ. Psychophysiology of fear imagery: Differences between focal phobia and social performance anxiety. *J Consult Clin Psychol*. 1978; 46:1157–9. [PubMed: 701554]
24. Cook EW 3rd, Melamed BG, Cuthbert BN, McNeil DW, Lang PJ. Emotional imagery and the differential diagnosis of anxiety. *J Consult Clin Psychol*. 1988; 56:734–40. [PubMed: 3192790]
25. McNeil DW, Vrana SR, Melamed BG, Cuthbert BN, Lang PJ. Emotional imagery in simple and social phobia: Fear versus anxiety. *J Abnorm Psychol*. 1993; 102:212–25. [PubMed: 8315134]
26. Cuthbert BN, Lang PJ, Strauss C, Drobos D, Patrick CJ, Bradley MM. The psychophysiology of anxiety disorder: Fear memory imagery. *Psychophysiology*. 2003; 40:407–22. [PubMed: 12946114]
27. Lang, PJ.; McTeague, LM.; Cuthbert, BN. Fearful imagery and the anxiety disorder spectrum. In: Rothbaum, BO., editor. *Pathological Anxiety: Emotional Processing in Etiology and Treatment*. Guilford Press; New York: 2005. p. 56-77.
28. Lang, PJ.; McTeague, LM.; Cuthbert, BN. Fear, anxiety, depression, and the anxiety disorder spectrum: A psychophysiological analysis. In: Treat, T.; Baker, T., editors. *Psychological Clinical Science: Recent Advances in Theory and Practice. Integrative Perspectives in Honor of Richard M. McFall*. Lawrence Erlbaum Associates; Mahwah, NJ: 2007. p. 167-195.
29. Lang PJ, McTeague LM. The anxiety disorder spectrum: Fear imagery, physiological reactivity, and differential diagnosis. *Anxiety Stress Coping*. 2009; 22:5–25. [PubMed: 19096959]
30. McTeague LM, Lang PJ, Laplante MC, Cuthbert BN, Strauss CC, Bradley MM. Fearful imagery in social phobia: Generalization, comorbidity, and physiological reactivity. *Biol Psychiatry*. 2009; 65:374–82. [PubMed: 18996510]
31. McTeague LM, Lang PJ, Laplante MC, Cuthbert BN, Shumen JR, Bradley MM. Aversive imagery in posttraumatic stress disorder: Trauma recurrence, comorbidity, and physiological reactivity. *Biol Psychiatry*. 2010; 67:346–56. [PubMed: 19875104]
32. McTeague LM, Lang PJ, Laplante MC, Bradley MM. Aversive imagery in panic disorder: Agoraphobia severity, comorbidity, and defensive physiology. *Biol Psychiatry*. 2011; 70:415–24. [PubMed: 21550590]
33. McTeague LM, Lang PJ. The anxiety spectrum & the reflex physiology of defense: From circumscribed fear to broad distress. *Depression & Anxiety*. In press.
34. Brown, TA.; Barlow, DH.; DiNardo, PA.; Barlow, DH. *The Anxiety Disorder Interview Schedule for DSM-IV: Adult Version*. Oxford University Press; New York, NY: 1994.
35. Bradley, MM.; Lang, PJ. Technical Report. D-1. University of Florida; Gainesville, FL: 2007. Affective Norms for English Text (ANET): Affective ratings of text and instruction manual.
36. Benson, H. *The relaxation response*. Morrow; New York: 1975.
37. Bradley MM, Lang PJ. Measuring emotion: The Self-Assessment Manikin and the Semantic Differential. *J Behav Ther Exp Psychiatry*. 1994; 25:49–59. [PubMed: 7962581]

38. Cook, EW, 3rd. VPM reference manual. Birmingham, AL: 2000.
39. Globisch J, Hamm A, Schneider R, Vaitl D. A computer program for scoring reflex eyeblink and electrodermal responses written in Pascal. *Psychophysiology*. 1993; 39:S30.
40. Vasey MW, Thayer JF. The continuing problem of false positives in repeated measures ANOVA in psychophysiology: A multivariate solution. *Psychophysiology*. 1987; 24:479–86. [PubMed: 3615759]
41. Spielberger, CD.; Gorsuch, RL.; Lushene, PR.; Vagg, PR.; Jacobs, GA. Manual for the State-Trait Anxiety Inventory (STAI). Consulting Psychologists Press; Palo Alto, CA: 1983.
42. Beck, AT.; Steer, RA.; Brown, GK. Manual for the Beck Depression Inventory. 2nd edition. The Psychological Corporation; San Antonio, TX: 1996.
43. Watson D, Weber K, Assenheimer JS, Clark LA, Strauss ME, McCormick RA. Testing a tripartite model: I. Evaluating the convergent and discriminant validity of anxiety and depression symptom scales. *J Abnorm Psychol*. 1995; 104:3–14. [PubMed: 7897050]
44. Reiss S, Peterson RA, Gursky DM, McNally RJ. Anxiety sensitivity, anxiety frequency and the prediction of fearfulness. *Behav Res Ther*. 1986; 24:1–8. [PubMed: 3947307]
45. Wolpe, J.; Lang, P. Fear Survey Schedule. Educational and Industrial Testing Service; San Diego, CA: 1969.
46. Spielberger, CD. Manual for the State-Trait Anger Expression Inventory. Psychological Assessment Resources; Odessa, FL: 1988.
47. Devins GM. Using the illness intrusiveness ratings scale to understand health-related quality of life in chronic disease. *J Psychosom Res*. 2010; 68:591–602. [PubMed: 20488277]
48. Sheehan PW. A shortened form of Betts' questionnaire upon mental imagery. *J Clin Psychol*. 1967; 23:380–89.
49. Beiling PJ, Rowa K, Summerfeldt LJ, Swinson RP. Factor structure of the Illness Intrusiveness Rating Scale in patients diagnosed with anxiety disorders. *J Psychopathol Behav*. 23:223–230.
50. Wuyek LA, Antony MM, McCabe RE. Psychometric properties of the panic disorder severity scale: Clinician-administered and self-report versions. *Clin Psychol Psychother*. 2010; 18:234–243. [PubMed: 21110405]
51. Krueger RF. The structure of common mental disorders. *Arch Gen Psychiatry*. 1999; 56:921–6. [PubMed: 10530634]
52. Slade T, Watson D. The structure of common DSM-IV and ICD-10 mental disorders in the Australian general population. *Psychol Med*. 2006; 36:1593–600. [PubMed: 16882356]
53. Vollebergh WA, Iedema J, Bijl RV, de Graaf R, Smit F, Ormel J. The structure and stability of common mental disorders: The NEMESIS study. *Arch Gen Psychiatry*. 2001; 58:597–603. [PubMed: 11386990]
54. Hettema JM, Prescott CA, Myers JM, Neale MC, Kendler KS. The structure of genetic and environmental risk factors for anxiety disorders in men and women. *Arch Gen Psychiatry*. 2005; 62:182–9. [PubMed: 15699295]
55. Kendler KS, Prescott CA, Myers J, Neale MC. The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Arch Gen Psychiatry*. 2003; 60:929–37. [PubMed: 12963675]
56. Chantarujikapong SI, Scherrer JF, Xian H, Eisen SA, Lyons MJ, Goldberg J, Tsuang M, True WR. A twin study of generalized anxiety disorder symptoms, panic disorder symptoms and post-traumatic stress disorder in men. *Psychiatry Res*. 2001; 103:133–69. [PubMed: 11549402]
57. Dichter GS, Tomarken AJ, Shelton RC, Sutton SK. Early- and late-onset startle modulation in unipolar depression. *Psychophysiology*. 2004; 41:433–40. [PubMed: 15102129]
58. Forbes EE, Miller A, Cohn JF, Fox NA, Kovacs M. Affect-modulated startle in adults with childhood-onset depression: relations to bipolar course and number of lifetime depressive episodes. *Psychiatry Res*. 2005; 134:11–25. [PubMed: 15808286]
59. Rottenberg J, Gross JJ, Gotlib IH. Emotion context insensitivity in major depressive disorder. *J Abnorm Psychol*. 2005; 114:627–39. [PubMed: 16351385]

60. Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and mental illness: A population-based prevalence study. *JAMA*. 2000; 284:2606–10. [PubMed: 11086367]
61. Larson CL, Schaefer HS, Siegle GJ, Jackson CA, Anderle MJ, Davidson RJ. Fear is fast in phobic individuals: Amygdala activation in response to fear-relevant stimuli. *Biol Psychiatry*. 2006; 60:410–7. [PubMed: 16919528]
62. Davis M, Walker DL, Miles L, Grillon C. Phasic vs sustained fear in rats and humans: Role of the extended amygdala in fear vs anxiety. *Neuropsychopharmacology*. 2010; 35:105–35. [PubMed: 19693004]
63. Radley JJ, Morrison JH. Repeated stress and structural plasticity in the brain. *Ageing Res Rev*. 2005; 4:271–87. [PubMed: 15993654]
64. Uchida S, Hara K, Kobayashi A, Funato H, Hobara T, Otsuki K, Yamagata H, McEwen BS, Watanabe Y. Early life stress enhances behavioral vulnerability to stress through the activation of REST4-mediated gene transcription in the medial prefrontal cortex of rodents. *J Neurosci*. 2010; 30:15007–18. [PubMed: 21068306]
65. Avgustinovich DF, Kovalenko IL, Kudryavtseva NN. A model of anxious depression: Persistence of behavioral pathology. *Neurosci Behav Physiol*. 2005; 35:917–24. [PubMed: 16270173]
66. Rygula R, Abumaria N, Flügge G, Fuchs E, Rütger E, Havemann-Reinecke U. Anhedonia and motivational deficits in rats: Impact of chronic social stress. *Behav Brain Res*. 2005; 162:127–34. [PubMed: 15922073]
67. Dwivedi Y, Mondal AC, Payappagoudar GV, Rizavi HS. Differential regulation of serotonin (5HT)2A receptor mRNA and protein levels after single and repeated stress in rat brain: Role in learned helplessness behavior. *Neuropharmacology*. 2005; 48:204–14. [PubMed: 15695159]
68. Ressler KJ, Mercer KB, Bradley B, Jovanovic T, Mahan A, Kerley K, Norrholm SD, Kilaru V, Smith AK, Myers AJ, Ramirez M, Engel A, Hammack SE, Toufexis D, Braas KM, Binder EB, May V. Post-traumatic stress disorder is associated with PACAP and the PAC1 receptor. *Nature*. 2011; 470:492–7. [PubMed: 21350482]
69. Grillon C, Dierker L, Merikangas KR. Fear-potentiated startle in adolescent offspring of parents with anxiety disorders. *Biol Psychiatry*. 1998; 44:990–7. [PubMed: 9821563]
70. Vaidyanathan U, Patrick CJ, Iacono WG. Patterns of comorbidity among mental disorders: a person-centered approach. *Compr Psychiatry*. 2011; 52:527–35. [PubMed: 21111407]
71. Krueger RF, Finger MS. Using item response theory to understand comorbidity among anxiety and unipolar mood disorders. *Psychol Assess*. 2001; 13:140–51. [PubMed: 11281035]

Figure 1.

Mean startle reflex responses (standardized to the distribution of responses during intertrial intervals; top panel), skin conductance level change (middle panel), and heart rate change (residuals; bottom panel) during neutral, panic attack, survival threat, and personal threat imagery for control and principal specific phobia groups. Error bars refer to standard error of the mean.

Figure 2.

Mean startle reflex responses (standardized to the distribution of responses during intertrial intervals; top panel), skin conductance level change (middle panel), and heart rate change (residuals; bottom panel) during neutral, panic attack, survival threat, and personal threat imagery for control and principal (single and multiple phobia) and non-principal specific phobia groups. Error bars refer to standard error of the mean.

Figure 3.

Mean subscale scores on the Mood and Anxiety Symptom Scale (MASQ; 43) for control and principal (single and multiple phobia) and non-principal specific phobia groups. Given the different ranges of the subscales, total scores were standardized across participants.

Figure 4.

Mean fear potentiation of startle reflexes (startle response magnitude during personal threat minus neutral imagery; top panel) and Beck Depression Inventory total score (42) for principal and non-principal specific phobia patients according to diagnostic prominence. Error bars refer to standard error of the mean.

Table 1

Mean Responses and Standard Deviations to Imagery Scenes by Control and Principal Specific Phobia Groups

Response Modality/Imagery scene	Control	Principal Specific phobia	Group Effect
Emotional Valence (1–9)			
Neutral	6.93 (1.53) *	6.91 (1.29) *	$F(1, 150) = 0.15, ns$
Panic Attack	4.00 (1.08) *	3.96 (1.30) *	$F(1, 149) = 0.12, ns$
Survival Threat	2.68 (0.99)	2.95 (1.24) *	$F(1, 150) = 3.20, ns$
Personal Threat	2.65 (1.41)	2.18 (1.67)	$F(1, 148) = 3.35, p < .05^{\dagger}$
Emotional Arousal (1–9)			
Neutral	2.31 (1.54) *	2.73 (1.74) *	$F(1, 147) = 2.40, ns$
Panic Attack	5.37 (1.90) *	5.86 (1.61) *	$F(1, 147) = 2.82, ns$
Survival Threat	6.69 (1.53) *	6.64 (1.65) *	$F(1, 147) = 0.05, ns$
Personal Threat	7.69 (1.67)	8.17 (1.36)	$F(1, 145) = 3.63, p < .05^{\dagger}$
Startle Reflex (T-score)			
Neutral	49.76 (5.01) *	51.15 (7.02) *	$F(1, 133) = 2.75, ns$
Panic Attack	51.51 (7.20) *	55.25 (14.08) *	$F(1, 133) = 3.80, ns$
Survival Threat	54.57 (8.38)	57.34 (16.09) *	$F(1, 133) = 1.62, ns$
Personal Threat	53.58 (7.33)	61.55 (20.87)	$F(1, 133) = 8.81, p < .01$
Heart Rate Δ (bpm)			
Neutral	-0.25 (2.07) *	-1.17 (1.98) *	$F(1, 145) = 7.67, p < .01$
Panic Attack	0.10 (2.25) *	-0.07 (1.91) *	$F(1, 145) = 0.26, ns$
Survival Threat	0.23 (1.16) *	0.26 (1.29) *	$F(1, 145) = 0.27, ns$
Personal Threat	1.21 (2.26)	2.61 (3.01)	$F(1, 144) = 10.21, p < .01$
SCL Δ (log ($\mu S + 1$))			
Neutral	-0.006 (0.036) *	-0.011 (0.058) *	$F(1, 143) = 0.35, ns$
Panic Attack	0.004 (0.046) *	-0.006 (0.057) *	$F(1, 142) = 1.26, ns$
Survival Threat	0.004 (0.022) *	0.009 (0.051) *	$F(1, 142) = 0.69, ns$
Personal Threat	0.045 (0.086)	0.139 (0.189)	$F(1, 142) = 15.05, p < .001$
Corrugator EMG Δ (μV)			
Neutral	-0.07 (0.79) *	-0.32 (1.10) *	$F(1, 146) = 2.43, ns$
Panic Attack	0.49 (1.40) *	0.54 (2.53) *	$F(1, 146) = 0.02, ns$
Survival Threat	0.96 (1.98)	0.93 (1.62) *	$F(1, 146) = 0.01, ns$
Personal Threat	1.17 (2.60)	1.45 (3.29)	$F(1, 146) = 0.33, ns$

Note. Valence rated on Self-Assessment Manikin (SAM) (40) 1=Completely unhappy, 9=Completely happy; Arousal rated on SAM 1=Completely relaxed, 9=Completely aroused; SCL=skin conductance level; Δ =change; μS =microsiemen; bpm=residual beats per minute after removal of baseline effects; EMG=electromyographic; μV =microvolt;

* Within-group comparison to personal threat significant at $p < .05$

† =one-tailed planned between-group comparison

Table 2

Mean Responses and Standard Deviations to Imagery Scenes by Control and Principal and Non-principal Specific Phobia Groups

Response Modality/Imagery scene	Principal Specific Phobia: Single Fear	Principal Specific phobia: Multiple Fears	Non-principal/Additional Specific Phobia	Group Effect
Emotional Valence (1–9)				
Neutral	7.09 (1.27) ^C	6.52 (1.29)	6.84 (1.46) ^a	$F(2, 154) = 1.43$, <i>ns</i>
Panic Attack	3.96 (1.30)	3.96 (1.34)	3.27 (1.56)	$F(2, 152) = 4.32$, $p < .05$
Survival Threat	2.92 (1.22)	3.02 (1.30)	2.53 (1.10)	$F(2, 155) = 2.66$, <i>ns</i>
Personal Threat	2.03 (1.51)	2.50 (1.97)	1.83 (1.18)	$F(2, 150) = 1.97$, <i>ns</i>
Emotional Arousal (1–9)				
Neutral	2.70 (1.80)	2.79 (1.65)	2.68 (1.63)	$F(2, 154) = 0.04$, <i>ns</i>
Panic Attack	5.82 (1.77) ^C	5.94 (1.24)	6.69 (1.55) ^a	$F(2, 154) = 5.41$, $p < .05$
Survival Threat	6.62 (1.73)	6.66 (1.51)	7.11 (1.51)	$F(2, 155) = 1.79$, <i>ns</i>
Personal Threat	8.17 (1.41)	8.15 (1.27)	7.87 (1.82)	$F(2, 151) = 0.63$, <i>ns</i>
Startle Reflex (T-score)				
Neutral	52.01 (7.68)	49.52 (5.32)	49.86 (6.34)	$F(2, 143) = 2.82$, <i>ns</i>
Panic Attack	56.20 (16.38)	53.45 (8.04)	51.81 (7.34)	$F(2, 143) = 2.26$, <i>ns</i>
Survival Threat	59.02 (18.44)	54.12 (9.78)	54.31 (11.95)	$F(2, 143) = 1.79$, <i>ns</i>
Personal Threat	63.83 (24.33) ^C	57.19 (10.94)	53.97 (10.97) ^a	$F(2, 143) = 5.25$, $p < .01$
Heart Rate Δ (bpm)				
Neutral	−1.07 (2.00)	−1.39 (1.96)	−0.64 (1.84)	$F(2, 156) = 1.55$, <i>ns</i>
Panic Attack	−0.11 (2.02)	0.00 (1.71)	0.37 (1.49)	$F(2, 156) = 1.47$, <i>ns</i>
Survival Threat	0.12 (1.28)	0.55 (1.28)	0.05 (1.12)	$F(2, 156) = 1.61$, <i>ns</i>
Personal Threat	2.84 (3.26) ^C	2.17 (2.42)	1.29 (2.56) ^a	$F(2, 155) = 4.21$, $p < .05$
SCL Δ (log ($\mu S + 1$))				
Neutral	−0.015 (0.059)	−0.001 (0.055)	0.005 (0.049)	$F(2, 153) = 2.41$, <i>ns</i>
Panic Attack	−0.001 (0.058)	−0.015 (0.055)	0.005 (0.046)	$F(2, 152) = 1.46$, <i>ns</i>
Survival Threat	0.013 (0.046)	0.002 (0.061)	0.002 (0.050)	$F(2, 152) = 0.52$, <i>ns</i>
Personal Threat	0.158 (0.186) ^C	0.099 (0.192)	0.052 (0.093) ^a	$F(2, 152) = 7.98$, $p < .01$

Response Modality/Imagery scene	Principal Specific Phobia: Single Fear	Principal Specific phobia: Multiple Fears	Non-principal/Additional Specific Phobia	Group Effect
Corrugator EMG Δ (μ V)				
Neutral	-0.27 (1.08)	-0.43 (1.16)	0.14 (1.26)	$F(2, 155) = 3.01$, <i>ns</i>
Panic Attack	0.43 (0.88)	0.76 (4.28)	0.40 (1.31)	$F(2, 155) = 0.31$, <i>ns</i>
Survival Threat	0.99 (1.80)	0.81 (1.18)	0.86 (1.86)	$F(2, 155) = 0.09$, <i>ns</i>
Personal Threat	1.44 (2.55)	1.47 (4.50)	0.86 (1.73)	$F(2, 155) = 1.03$, <i>ns</i>

Note. Valence rated on Self-Assessment Manikin (SAM) (40) 1=Completely unhappy, 9=Completely happy; Arousal rated on SAM 1=Completely relaxed, 9=Completely aroused; SCL=skin conductance level; Δ =change; μ S=microsiemen; bpm=residual beats per minute after removal of baseline effects; EMG=electromyographic; μ V=microvolt;

^a = Post hoc between-group comparison to single phobia group is significant at $p < .05$

^b = Post hoc between-group comparison to multiple phobia group is significant at $p < .05$

^c = Post hoc between-group comparison to non-principal phobia group significant at $p < .05$

^d =one-tailed planned between-group comparison.

Table 3

Questionnaire and interview responses (means and standard deviations) for Control and Principal and Non-principal Specific Phobia Groups

Measure	Control	Principal Specific Phobia: Single Fears	Principal Specific Phobia: Multiple Fears	Non-principal/Additional Specific Phobia	Group Effect
Questionnaire Measures of Broad Distress					
STAI-Trait (20–80)	30.80 (8.57) ^{bcd}	37.17 (10.42) ^{ad}	42.29 (11.13) ^a	56.55 (11.00) ^{ab}	$F(3, 226) = 90.09, p < .001$
BDI Total (0–63)	3.51 (4.55) ^{bcd}	8.08 (7.46) ^{ad}	12.38 (8.66) ^{ad}	21.80 (11.71) ^{abc}	$F(3, 230) = 63.21, p < .001$
MASQ Anhedonia (22–110)	46.26 (13.32) ^{cd}	52.92 (15.12) ^d	59.86 (17.27) ^{ad}	77.09 (15.23) ^{abc}	$F(3, 227) = 62.80, p < .001$
ASI Total (0–64)	9.06 (6.83) ^{bcd}	21.55 (10.91) ^{acd}	29.57 (12.34) ^{abd}	38.41 (13.64) ^{abc}	$F(3, 216) = 93.33, p < .001$
FSS Total (103–515)	156.47 (37.42) ^{bcd}	195.00 (50.88) ^{ad}	217.69 (54.76) ^{ad}	265.89 (75.92) ^{abc}	$F(3, 229) = 48.41, p < .001$
STAXI-Trait (10–40)	14.42 (3.94) ^d	15.73 (5.46) ^d	17.22 (4.60)	19.65 (5.83) ^{ab}	$F(3, 227) = 15.04, p < .001$
IIRS Total (13–91)	18.24 (11.64) ^{bcd}	26.13 (14.68) ^{ad}	29.68 (14.69) ^{ad}	51.66 (19.44) ^{abc}	$F(3, 213) = 60.20, p < .001$
QMI Total (35–245)	82.96 (29.70)	88.15 (23.80)	97.09 (39.15)	89.36 (32.70)	$F(3, 224) = 1.37, ns$
Interview Measures of Specific Phobia					
Fear of worst phobia (0–8)		6.72 (2.20)	6.43 (2.15)	6.75 (1.87)	$F(2, 156) = 0.23, ns$
Avoidance of worst phobia (0–8)		6.77 (1.91)	6.61 (2.69)	6.73 (2.07)	$F(2, 156) = 0.05, ns$
Interference secondary to worst phobia (0–8)		5.01 (2.40)	5.68 (2.35)	5.21 (1.99)	$F(2, 157) = 0.77, ns$
Distress secondary to worst phobia (0–8)		6.02 (2.03)	6.30 (2.05)	5.53 (2.05)	$F(2, 157) = 1.73, ns$
Specific phobia chronicity (Years)		15.16 (14.80)	13.88 (11.70)	17.48 (13.14)	$F(2, 151) = 0.86, ns$
Prognosis (1–4)		1.44 (0.50) ^d	1.75 (0.61) ^d	2.50 (0.75) ^{bc}	$F(2, 157) = 43.54, p < .001$
Interview Measures of Comorbidity					
Axis I disorders (Count)		1.28 (0.67) ^{cd}	2.58 (0.83) ^{bd}	3.64 (1.30) ^{bc}	$F(2, 157) = 76.35, p < .001$
Comorbid anxiety (excluding specific phobia) disorder (%)		10.0 ^{cd}	20.83 ^{bd}	98.83 ^{bc}	$\chi^2(2) = 120.84, p < .001$
Comorbid mood disorder (%)		10.0 ^d	12.50 ^d	66.28 ^{bc}	$\chi^2(2) = 50.78, p < .001$
Comorbid single major depressive disorder (%)		2.0 ^d	4.12	12.79 ^b	$\chi^2(2) = 5.52, ns$
Comorbid recurrent major depressive disorder (%)		2.0 ^d	8.33 ^d	36.0 ^{bc}	$\chi^2(2) = 24.72, p < .001$

Measure	Control	Principal Specific Phobia: Single Fears	Principal Specific Phobia: Multiple Fears	Non-principal/Additional Specific Phobia	Group Effect
Comorbid anxious misery disorder (%)		6.0 ^d	8.30 ^d	84.89 ^{bc}	$\chi^2(2) = 97.21, p < .001$
Demographics					
Age at assessment (Years)	31.79 (11.61)	34.98 (13.97)	36.83 (11.90)	36.20 (12.75)	$F(3, 232) = 1.99, ns$
Gender (% Female)	65.79	72.0	79.17	72.09	$\chi^2(3) = 1.84, ns$
Race (% Caucasian)	84.21	86.0	75.0	77.91	$\chi^2(3) = 2.43, ns$
College graduate (%)	60.53 ^d	56.0 ^d	54.17 ^d	32.56 ^{abc}	$\chi^2(3) = 14.58, p < .001$

Note. STAI-Trait = Trait scale of State Trait Anxiety Inventory (41); BDI = Beck Depression Inventory (42); MASQ Anhedonia = Anhedonia subscale of the Mood and Anxiety Symptom Questionnaire (43); ASI = Anxiety Sensitivity Index (44); FSS = Fear Survey Schedule (45); STAXI-Trait = Trait scale of State Trait Anger Expression Inventory (46); IIRS = Illness Intrusiveness Rating Scale (47); QMI Total = Questionnaire on Mental Imagery (48); Fear/avoidance of worst phobia = Severity ratings from ADIS-IV (9-point scale ranging from 0, *No fear/Never avoid*, to 8, *Very severe fear/Always avoids*) (34). Interference/distress secondary to worst phobia = Severity ratings from ADIS-IV (9-point scale ranging from 0, *None*, to 8, *Very severe*); Specific phobia chronicity = years from patient-reported onset of earliest phobia to assessment; Prognosis = clinician-rated estimate of treatment outcome (4-point scale ranging from 1, *Excellent*, to 4, *Poor*); Superscripts = Results of Tukey HSD pairwise comparisons:

^a = Post hoc between-group comparison to control is significant at $p < .05$

^b = Post hoc between-group comparison to single phobia is significant at $p < .05$

^c = Post hoc between-group comparison to multiple phobia is significant at $p < .05$.

^d = Post hoc between-group comparison to non-principal phobia is significant at $p < .05$.