Figure 2 Loose body grasped with tool

Figure 3 Loose body reduced with shaver

Figure 4 Removal of loose body once reduced

TECHNIQUE

Three arthroscopy portals are made: one for viewing, one for grasping the loose body and one for introduction of a shaver. While the scrub nurse holds the camera, the surgeon grasps the loose body and manipulates it to the shaver (Figs 1 and 2). The loose body is then reduced to a suitable size (Fig 3) to enable safe removal through a standard arthroscopic portal (Fig 4).

DISCUSSION

The only potential danger with this technique is inadvertent damage to the grasper by the shaver. This can be avoided by good surgical technique. By using this operative technique, a large arthrotomy can be avoided.

References

1. Henderson MS. Loose bodies in the knee-joint. *J Bone Joint Surg* 1916; **S2-14**: 265–280.
2. Gillquist J, Hagberg G, Oretorp N. Therapeutic arthroscopy of the knee. *Injury* 1978; **10**: 128–132.
3. Lee YS, Ahn JH, Min BH, Nha KW. Arthroscopic loose body removal and cyst decompression using a posterior trans-septal portal in the blind spot during knee arthroscopy – technical report. *Knee* 2011; **18**: 55–58.
4. Froelich JM, Hillard-Sembell D. Symptomatic loose bodies of the knee located in a popliteal cyst. *Orthopedics* 2009; **32**: 918.
5. Ahn JH, Yoo JC, Lee SH. Arthroscopic loose-body removal in posterior compartment of the knee joint: a technical note. *Knee Surg Sports Traumatol Arthrosc* 2007; **15**: 100–106.

Ileoanal pouch: short mesentery? Lengthen the pouch

RWL Ma, DM Gold

University of New South Wales, Sydney, Australia

CORRESPONDENCE TO

Darren Gold, E: d.gold@unsw.edu.au

BACKGROUND

Ileoanal pouch surgery is the preferred option following a proctocolectomy for ulcerative colitis and familial adenomatous polyposis, with the 'J' pouch favoured by most surgeons. The pouch is anastomosed to the anus following complete excision of the rectum and its mucosa.^{1,2} Techniques to lengthen the mesentery include pouch folding,³ mesenteric incisions and division.⁴ For a tension free anastomosis, the pouch should extend 6cm below the pubic symphysis.⁵ We describe a technique to lengthen the 'J' pouch that avoids mesenteric surgery.

TECHNIQUE

Most 'J' pouch enterotomies are made at the apex of the pouch, allowing introduction of the stapling device. This usually corresponds with the point on the ileum with the greatest reach. It was observed that once the pouch is created, extra dependency is obtained by closing the original enterotomy and recreating it approximately 4cm proximal to the apex on the antimesenteric border (Figs 1 and 2). This allows for an extra 4cm of reach. Even if the original enterotomy is made at the very tip of the apex, a further 2–3cm are achieved. This is due to the increased ability for 'stretching' of the surrounding bowel wall, no longer impeded by nearby mesentery.

Figure 1 The increase in length obtainable by creating a new opening 4cm proximal to the apex on the antimesenteric border, compared to the original opening

Figure 2 The new longer pouch: Sutures are in place where the original enterotomy was closed

DISCUSSION

With the described technique all subsequent pouches have reached to the dentate line without the need for mesenteric manoeuvres. Mesenteric manipulation risks vascular compromise. We have experienced no complications related to this technique. It is simple to perform and increases reach by approximately 4cm, which is comparable to other techniques.⁵

References

1. Onaitis MW, Mantyh C. Ileal pouch-anal anastomosis for ulcerative colitis and familial adenomatous polyposis: historical development and current status. *Ann Surg* 2003; **238**(6 Suppl): S42–48.
2. Parks AG, Nicholls RJ. Proctocolectomy without ileostomy for ulcerative colitis. *Br Med J* 1978; **2**: 85–88.
3. Byrne CM, Uraiqat AA, Phillips RK. A systematic stapling defect – a potential cause of ‘anastomotic’ leak after restorative proctocolectomy. *Colorectal Dis* 2008; **10**: 286–288.
4. Burnstein MJ, Schoetz DJ Jr, Collier JA, Veidenheimer MC. Technique of mesenteric lengthening in ileal reservoir-anal anastomosis. *Dis Colon Rectum* 1987; **30**: 863–866.
5. Smith L, Friend WG, Medwell SJ. The superior mesenteric artery. The critical factor in the pouch pull-through procedure. *Dis Colon Rectum* 1984; **27**: 741–744.

Safe and inexpensive alternative to suture passer

JD Kosy, IT Sharpe

Royal Devon and Exeter Hospital, Exeter, UK

CORRESPONDENCE TO

Jonathan Kosy, E: jonkosy@yahoo.co.uk

BACKGROUND

As the name suggests, the suture passer is used to achieve solid fixation of structures, often removed during the operative approach, by pulling an anchoring suture through an osseous tunnel. In addition, the suture passer can be used to pass tendons and other soft tissue structures through bone during reconstructive procedures. The expense of this single use instrument has been highlighted and a potentially dangerous alternative suggested.

Figure 1 Loop of nylon suture passed through drawing-up needle

Our method is inexpensive, successful and uses materials available in all theatres.

TECHNIQUE

A loop of nylon suture is passed through a blunt ended 18G drawing-up needle (Fig 1). Leaving a 5mm loop protruding from the end, this is passed through the tunnel. The suture or tendon for reattachment is passed through the loop and pulled back through the tunnel (Fig 2).

Figure 2 Needle (with suture loop) passed through metatarsal head