References

1. Nanchahal J, Nayagam S, Khan U *et al*. *Standards for the Management of Open Fractures of the Lower Limb*. London: Royal Society of Medicine Press; 2009.
2. Townley WA, Nguyen DQ, Rooker JC *et al*. Management of open tibial fractures – a regional experience. *Ann R Coll Surg Engl* 2010; **92**: 693–696.
3. Gopal S, Majumder S, Batchelor AG *et al*. Fix and flap: the radical orthopaedic and plastic treatment of severe open fractures of the tibia. *J Bone Joint Surg Br* 2000; **82**: 959–966.
4. Nicholson D. *Establishment of Regional Networks of Trauma Care*. London: Department of Health; 2010.

Straight to colonoscopy: the ideal patient pathway for the 2-week suspected cancer referrals?

SA Fallis, K Busby, J Francombe, MJ Osborne

Warwick Hospital, Warwick, UK
doi 10.1308/147870811X591909

CORRESPONDENCE TO

Simon Fallis, E: simonfallis@hotmail.com

COMMENT ON

Beggs AD, Bhate RD, Irukulla S *et al*. Straight to colonoscopy: the ideal patient pathway for the 2-week suspected cancer referrals? *Ann R Coll Surg Engl* 2011; **93**: 114–119
doi 10.1308/003588411X12851639107917

In order to meet two-week referral targets for patients with colorectal symptoms, trusts must provide an efficient and effective patient pathway. On the face of it, the paper by Beggs *et al* suggests an attractive model for streamlining the process and potentially lowering costs. However, when one considers factors other than efficiency and cost effectiveness, the pathway may appear less than ideal.

In the straight-to-colonoscopy pathway described, patients are sent full bowel preparation with instructions. In response to a number of adverse incidents, the National Patient Safety Agency has published clear guidelines for the use of bowel cleansing solutions.¹ These include the need for clinical assessment to be undertaken by the clinician responsible for the procedure to ensure no contraindication or risks from their use. By the time a patient on a straight-to-colonoscopy pathway first sees the clinician responsible he or she will have already taken his or her bowel preparation. This is too late for an assessment of risk and an explanation on the safe use of the product.

A second but related concern pertains to consent. Guidelines for obtaining informed consent are widely circulated by the General Medical Council² and Department of Health.³ For elective procedures it is generally considered good practice to seek the patient's consent well in advance, with the clinician checking before the procedure starts that the person still consents. If a person is not asked to signify his or her consent until just before the procedure is due to start, at a time when he or she may be feeling particularly vulnerable, there may be real doubts as to its validity.

While it remains important to meet targets and continue to improve the efficiency of our service, we would caution colleagues in the potential for compromise in the quality of that service.

References

1. National Patient Safety Agency. *Reducing Risk of Harm from Oral Bowel Cleansing Solutions*. NPSA/2009/RRR012. London: NPSA; 2009.
2. General Medical Council. *Consent: Patients and Doctors Making Decisions Together*. London: GMC; 2008.
3. Department of Health. *Reference Guide to Consent for Examination or Treatment*. 2nd edn. London: DH; 2009.

Author's response

AD Beggs, RD Bhate, S Irukulla, M Achiek, AM Abulafi

Croydon University Hospital, Croydon, UK
doi 10.1308/147870811X591909

CORRESPONDENCE TO

Andrew Beggs, E: andrew@andrewbeggs.org

We thank Fallis *et al* for their observations on our pathway and paper. To answer their specific points, this study was carried out before the National Patient Safety Agency (NPSA) alert was issued. However, our patients collect their bowel preparation from a nurse at the endoscopy unit prior to the procedure. Patients are asked by the nurse whether they have any potential contraindications to bowel preparation as discussed in the NPSA guidance.¹ After the NPSA alert was issued we also changed the referral form so that general practitioners are prompted to answer questions about co-morbidities and to request a blood test that will be available on the computer system when the patient attends the nurse appointment to collect the bowel preparation. If there is any potential problem, the case is discussed with an endoscopist present within the endoscopy unit and the preparation is given only if it is safe to proceed.

With regards to Fallis *et al*'s second point pertaining to consent, patients are sent a booklet describing the risks and benefits of colonoscopy at the time the appointment is booked. Patients are also specifically informed that they can discuss the procedure or raise any concerns either when collecting the bowel preparation or immediately before the procedure. We feel that this ensures valid consent as recommended in the General Medical Council and Department of Health guidelines.^{2,3}

In summary, we feel our pathway allows rapid access to the gold standard diagnostic investigations in an appropriate, safe and timely fashion.

References

1. National Patient Safety Agency. *Reducing Risk of Harm from Oral Bowel Cleansing Solutions*. NPSA/2009/RRR012. London: NPSA; 2009.
2. General Medical Council. *Consent: Patients and Doctors Making Decisions Together*. London: GMC; 2008.
3. Department of Health. *Reference Guide to Consent for Examination or Treatment*. 2nd edn. London: DH; 2009.