Published in final edited form as:

Neuroscience. 2012 July 12; 214: 84–105. doi:10.1016/j.neuroscience.2012.03.047.

Long lasting transcriptional refractoriness triggered by a single exposure to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine

Roberto Pattarini¹, Yongqi Rong¹, Kennie R. Shepherd³, Yun Jiao¹, Chunxu Qu, Richard J. Smeyne¹, and James I. Morgan¹

¹Department of Developmental Neurobiology, St. Jude Children's Research Hospital, Memphis, TN, 38105

²Hartwell Center for Bioinformatics and Biotechnology, St. Jude Children's Research Hospital, Memphis, TN, 38105

³Morehouse School of Medicine, Atlanta, GA, 30310

Abstract

Parkinson's disease (PD) is a progressive neurodegenerative disorder whose etiology is thought to have environmental (toxin) and genetic contributions. The neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine (MPTP) induces pathological features of PD including loss of dopaminergic neurons in the substantia nigra pars compacta (SNpc) and striatal dopamine (DA) depletion. We previously described the striatal transcriptional response following acute MPTP administration in MPTP-sensitive C57BL/6J mice. We identified three distinct phases: early (5h), intermediate (24 h) and late (72 h) and reported that the intermediate and late responses were absent in MPTP-resistant Swiss-Webster (SWR) mice. Here we show that C57BL/6J mice pre-treated with a single 40 mg/kg dose of MPTP and treated 9 days later with 4×20 mg/kg MPTP, display a striatal transcriptional response similar to that of MPTP-resistant SWR mice i.e. a robust acute response but no intermediate or late response. Transcriptional refractoriness is dependent upon the dose of the priming challenge with as little as 10 mg/kg MPTP being effective and can persist for more than 28 days. Priming of SWR mice has no effect on their response to subsequent challenge with MPTP. We also report that paraquat, another free radical producer, also elicits striatal transcriptional alterations but these are largely distinct from those triggered by MPTP. Paraquat-induced changes are also refractory to priming with paraquat. However neither paraquat nor MPTP elicit cross-attenuation. Thus exposure to specific toxins triggers distinct transcriptional responses in striatum that are influenced by prior exposure to the same toxin. The prolonged refractory period described here for MPTP could explain at the molecular level the reported discrepancies between different MPTP administration regimens and may have implications for our understanding of the relationship between environmental toxin exposure and PD.

Keywords

Parkinson's disease; MPTP; C57BL/6J; Swiss Webster; dopamine; paraquat

1. Introduction

Parkinson's disease (PD) is a progressive neurodegenerative disorder characterized by the loss of dopaminergic (DA) neurons in the substantia nigra pars compacta (SNpc). The etiology of sporadic PD, the most frequent form of the disorder, remains uncertain, although it likely involves interplay between environmental toxins (Landrigan et al., 2005, Brown et al., 2006) and predisposing genetic risk factors (Huang et al., 2004, Benmoyal-Segal and Soreq, 2006, Farrer, 2006, Wood-Kaczmar et al., 2006). Several reports indicate that exposure to environmental agents and chronic use of certain substances alters the incidence of PD (Tanner et al., 2011). Cigarette smoking (Tanaka et al., 2010), chronic intake of ibuprofen (Gao and Hong, 2011), or caffeine (Costa et al., 2010), have been associated with a lower incidence of PD. On the other hand several studies indicate that viral infections (Jang et al., 2009, Henry et al., 2010), prenatal exposure to lipopolysaccharide (Barlow et al., 2007), drinking well water (Koller et al., 1990), and life-long exposure to heavy metals (Weisskopf et al., 2010), pesticides and herbicides (Gatto et al., 2009), can increase the risk of developing PD.

The introduction of several toxin- and genetic-based models for PD has advanced our knowledge of the disease, although none of the models fully recapitulate the characteristics of sporadic PD (Bove et al., 2005, Meredith et al., 2008). The 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine (MPTP) model represents the most widely used and better-characterized toxin-based animal model that is currently available (Smeyne and Jackson-Lewis, 2005). This neurotoxin was first identified as the cause of PD-like symptoms in humans (Ballard et al., 1985) and was subsequently used to simulate PD in various animals including mice (Dauer and Przedborski, 2003, Smeyne and Jackson-Lewis, 2005). MPTP induces the characteristic loss of DAergic neurons in SNpc and a concomitant loss of striatal dopamine seen in PD (Dauer and Przedborski, 2003, Smeyne and Jackson-Lewis, 2005). Furthermore, sensitivity to MPTP is genetically influenced: some strains of mice, such as Swiss Webster (SWR), display a limited loss of DAergic neurons after MPTP administration, whereas other strains of mice such as C57BL/6J, display a larger cell loss (Sundstrom et al., 1987, German et al., 1996, Hamre et al., 1999, Vila et al., 2001). These strain differences provide an opportunity to identify predisposing genetic determinants of MPTP toxicity.

The striatum, that is innervated by the SNpc DAergic neurons, is likely to be the proximal target of neurotoxicity in the MPTP model (Boyd et al., 2007) and PD itself (Brooks, 1991). MPTP is a protoxin that rapidly crosses the blood brain barrier and it is metabolized to 1-methyl-4-phenylpyridinium (MPP⁺) by monoamine oxidase B (MAO-B) in glial cells (Smeyne and Jackson-Lewis, 2005). Within hours of MPTP exposure, DA is released in the striatum from synaptic stores (Bazzu et al., 2010), and dopaminergic nerve terminals show evidence of dysfunction (Nagai et al., 2007). Several days after this initial MPTP-induced dopamine depletion, microglia are activated and SNpc DAergic neurons degenerate (Boyd et al., 2007).

Using Affymetrix chip technology coupled with quantitative reverse transcribed PCR (qRT-PCR) we previously described a stereotypic change in gene expression patterns (Pattarini et al., 2008). We identified three distinct phases of MPTP-induced gene expression changes characterized by almost unique sets of genes. The earliest response was seen 5 hour after systemic administration of MPTP, and was followed by an intermediate (24 hour), and a late (72 hour) response. The changes in transcriptional response occur long before any sign of dopaminergic neuron degeneration can be identified in the SNpc (Boyd et al., 2007).

Administration of MPTP follows different dosing regimens that simulate different types of exposures. The acute paradigm uses doses of MPTP injected over a short period of time, whereas other regimens simulate prolonged exposure to lower doses of MPTP (Meredith et al., 2008). All the dosing regimens cause loss of DAergic neurons in SNpc and are often used interchangeably (Meredith et al., 2008). Nonetheless, evidence suggests that they may differ in various respects including the ability to induce proteinaceous inclusions (Alvarez-Fischer et al., 2008, Meredith et al., 2008), the progressiveness of the DAergic lesion in the SNpc, and recovery from injury of the surviving DAergic neurons (Meredith et al., 2008).

Here we have used Affymetrix gene expression analysis to compare mRNA expression profiles in various MPTP regimens. We have complemented this by comparing the expression profiles elicited by paraquat with those of MPTP and whether there is an interaction at the level of gene transcription between these two agents.

2. Experimental procedures

2.1 Animals and Experiments

8 week old female C57BL/6J mice were purchased from the Jackson Laboratories (Bar Harbor, ME, USA), and female Swiss Webster (SWR) mice were purchased from Harlan (Strain ND4, Indianapolis, IN, USA). Animals were housed in micro-isolator units with a 12 hour light-dark schedule and constant temperature with free access to food and water. When animals were 3–4 months of age, MPTP-HCl (Sigma, St. Louis, MO, USA) was administered by intraperitoneal (IP) injections at the dosage and schedule specified (see Results and Figure Legends). Further details regarding MPTP administration and biosafety have been previously reported (Przedborski et al., 2001, Smeyne et al., 2001). Animals were sacrificed at the time points after MPTP administration specified for each experiment (see Results and Figure Legends), and the striatum was rapidly dissected and immediately frozen on dry ice to preserve RNA integrity. Samples were stored at -80°C . All studies were approved by the St. Jude Children's Research Hospital Animal Care and Use Committee (ACUC) and were conducted in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals (NIH Publication No. 80-23, revised 1996). Efforts were made to minimize the number of animals involved in each experiment and their suffering.

2.2 RNA Isolation

Total RNA was extracted with TRIzol[®] Reagent (Invitrogen, Carlsbad, CA, USA) according to manufacturer instructions. Briefly, 1 ml TRIzol[®] was added to frozen samples and immediately homogenized. Samples were mixed with 200 μl of chloroform and centrifuged for 15 min at $12,000 \times g$ at 4°C . The aqueous phase was transferred to a new vial, mixed with 50 μg of glycogen from Roche Applied Science (Penzberg, Germany) and 500 μl of isopropanol and centrifuged (10 min at $12,000 \times g$, 4°C) to precipitate total RNA. The pellet was washed with ice cold 70% ethanol and air dried for 10 min. Total RNA was resuspended in RNase-free water and it was checked for integrity by agarose gel electrophoresis. Samples that appeared degraded were discarded.

2.3 Preparation of Samples for Microarray Analysis

Technical procedures for microarray analysis, including quality control of RNA, labeling, hybridization and scanning of the arrays were performed by the Hartwell Center for Bioinformatics & Biotechnology (HC) at St. Jude Children's Research Hospital (SJCRH) according to standard operating procedures for Affymetrix protocols (GeneChip[®] Expression Analysis manual, Affymetrix, Santa Clara CA, USA). Prior to use, RNA sample integrity was assessed with a 2100 Bioanalyzer Lab-on-a-chip system (Agilent

Technologies, Santa Clara, CA, USA). Total RNA samples that were not degraded were labeled using the Gene Chip IVT Labeling Kit (Affymetrix) according to manufacturer instructions. Further technical details are reported in Pattarini et al. (2008).

2.4 Analysis of Microarrays

The microarrays used in this study (Mouse Genome 430 2.0 Arrays, Affymetrix) contain 45,101 probe sets, representing 39,000 transcripts and variants. Scanned images were analyzed with the Gene Chip Operating Software (GCOSv1.2, Affymetrix). Assessment of probe set present/absent calls was made using the Single Array Analysis method in GCOS using statistical algorithm with default analysis parameters (http://www.affymetrix.com/support/technical/whitepapers/sadd_whitepaper.pdf). Probe set signal values were scaled by global methods to a target value of 500.

Array analysis was performed using Spotfire® DecisionSite 8.2 from TIBCO Software Inc. (Palo Alto, CA, USA) as previously reported in Pattarini et al. (2008).

Gene ontology (GO) and Gene set enrichment analysis (GSEA)—Gene ontology analysis for the probe sets differentially expressed between SM/SS (203 probe sets) and SP/SS (70 probe sets) was performed using DAVID Bioinformatics Resources (<http://david.abcc.ncifcrf.gov/>). GO terms with p-value less than 0.003 were selected.

To identify biological processes and molecular functions differentially represented between MPTP and paraquat treatments, we compared all the probe sets in the array of MPTP-treated mice (SM) to those of paraquat-treated mice (SP), using a permutation approach called gene set enrichment analysis (GSEA). GSEA is a computational method that determines whether an *a priori* defined set of genes shows statistically significant, concordant differences between two biological states. This analysis was performed using the R-version of a publicly available program at <http://www.broad.mit.edu/gsea> (Mootha et al., 2003, Subramanian et al., 2005). Gene sets for this analysis were downloaded from the Broad Institute Molecular Signature database (MSigDB) at <http://www.broadinstitute.org/gsea/msigdb/index.jsp>. Only results with a normalized enrichment score (NES) ≥ 1.7 and false discovery rate (q) ≤ 0.07 were considered. Further details can be found in Pattarini et al. (2007).

2.5 Validation of Microarray Data by Quantitative RT-PCR

Total RNA was reverse transcribed using TaqMan® reverse transcription reagents from Applied Biosystems (Foster City, CA, USA). Primers and probes for real-time PCR (qRT-PCR) were designed with Primer Express Software version 1.5 for Windows® (Applied Biosystems) and synthesized by the HC. Real time PCR was performed using TaqMan® PCR Core Reagent Kit (Applied Biosystem), and the ABI Prism 7900HT system (Applied Biosystem). Absolute quantification was performed using standard curves for each gene of interest. Primers and probes used for qRT-PCR, for both absolute and relative ($\Delta\Delta Ct$) quantification are listed in Table 1.

Standards for absolute quantification were prepared by cloning the coding sequence (CDS) of each gene into a pcDNA3 plasmid (Invitrogen) as previously described (Pattarini et al., 2007, Pattarini et al., 2008). The primers used to prepare the standards, including the restriction site used, are listed in Table 2.

2.6 Statistical Analysis

Statistical methods used to analyze microarray results are explained in the microarray analysis section. Statistical analysis for qRT-PCR results was performed with GraphPad Prism® version 5.0 for Macintosh® (GraphPad Software Inc., San Diego, CA). Results are

expressed as the ratio of number of copies of a specific gene over the number of copies of glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*). Each time point is the average of at least three animals. The temporal profile of each gene was analyzed by one-way ANOVA followed by Bonferroni's multiple comparisons test to assess statistical significance versus respective control.

2.7 HPLC quantitation of dopamine and its metabolites

Striata were rapidly removed at specified time points after a single dose of MPTP and placed on an ice-cooled plate for dissection of the striatum. Tissue was weighed and homogenized in 10 volumes of ice-cooled 0.1 M perchloric acid and centrifuged for 12 minutes at $10,000 \times g$ at 4°C. The supernatant was passed through a 0.2 µm filter and stored at -70°C. Dopamine (DA), and its metabolites 3-4-dihydroxyphenylacetic acid (DOPAC) and homovanillic acid (HVA) were analyzed using reverse-phase ion pairing HPLC combined with electro coupling (EC) detection under isocratic conditions (Acworth and Cunningham, 1999). The guard cell was set at +350 mV, with a screening electrode set at -150 mV, and working electrode set at +220 mV. The mobile phase consisted of 75 mM sodium dihydrogen phosphate monohydrate, 1.7 mM 1-octane sulfonic acid sodium salt, 0.73 mM triethylamine, 25 mM EDTA, 10% acetonitrile/90% water. The pH was adjusted to 3.0 with phosphoric acid. The mobile phase was delivered at a flow rate of 0.6 ml/min onto the MD-150 (3×150 mm, 3 µm) reverse phase column (ESA, Chelmsford, MA, USA). Twenty µl aliquots were injected by an autoinjector with cooling module set at 4°C. The levels of DA, DOPAC, and HVA in unknown samples were determined using standard curves created by injection of known concentrations of each substance.

3 Results

3.1 Differences between the acute and subchronic MPTP regimens

Two widely characterized MPTP dosing regimens, acute and subchronic, were used in this study (Petroske et al., 2001). The acute model consists of four injections of MPTP (20 mg/kg) spaced two hours apart, whereas the subchronic regimen consists of single daily injections of MPTP (40 mg/kg) for 4 days (Figure 1A).

We have previously reported that 24 h after systemic injection of MPTP, mRNA levels for the inducible form of heme oxygenase-1 (*Hmox1*) are elevated in striatal astrocytes (Fernandez-Gonzales et al., 2000). Therefore, we used mRNA expression of this gene as a benchmark to determine the minimal dose of MPTP necessary to elicit a transcriptional response (Figure 1B). C57BL/6J mice were injected with a single dose of MPTP at concentrations of 10, 20 and 40 mg/kg, sacrificed 24 h later and striatal *Hmox1* mRNA levels determined by qRT-PCR. A statistically significant increase in relative *Hmox1* mRNA was detected starting with the 20 mg/kg dose (Figure 1B). Thus, single doses of MPTP used in both the acute (20 mg/kg) and subchronic (40 mg/kg) regimens are able to modulate *Hmox1* expression.

We next compared striatal *Hmox1* expression profiles over time in the two MPTP models by qRT-PCR. For the acute model, we intraperitoneally injected C57BL/6J mice with MPTP (20 mg/kg) or saline over an eight hour period (0, 2, 4 and 6 h) and collected striata at 2, 4, 8, 12, 18, 24, 36 and 48 h after the first injection. Levels of *Hmox1* mRNA increased between 12 and 24 h and returned to basal levels by 36 h (Figure 1C), confirming results from our previous study (Fernandez-Gonzales et al., 2000). Using the subchronic regimen (Figure 1A), animals were intraperitoneally injected with MPTP (40 mg/kg) on day 0, 1, 2, and 3 and their striata were harvested 24 h after each injection (Figure 1D). Striatal *Hmox1* mRNA levels were elevated 24 h after the first injection (Figure 1D), however, further

injections of MPTP (40 mg/kg) failed to induce *Hmox1* gene expression (Figure 1D). Even a fifth dose of MPTP (for a total of 200 mg/kg) a week after the first MPTP injection failed to elevate *Hmox1* mRNA levels (Figure 1D). Thus exposure to a single dose of MPTP elicits a refractory period in which *Hmox1* cannot be reinduced.

3.2 A single dose of MPTP elicits a long-term refractory state for *Hmox1* induction

To confirm and extend our finding of an MPTP-induced refractory period, we investigated the dose-dependency and duration of this phenomenon. C57BL/6J mice were injected on day 0 with a priming dose of MPTP ranging from 1 to 40 mg/kg (Figure 2A). Some mice were sacrificed the next day to assess the effect of the priming dose on *Hmox1* mRNA levels (blue bars in figure 2A). The remaining animals received a second dose of 40 mg/kg MPTP on day 7 and were sacrificed 24 h later (brown bars in figure 2A). The dose-response at 24 h after the priming dose was similar to that reported in Figure 1B: only doses at or higher than 20 mg/kg increased *Hmox1* mRNA levels (compare Figure 1B with blue bars in Figure 2A). Pre-exposure to either 20 or 40 mg/kg abolished *Hmox1* gene expression after the second injection of MPTP (40 mg/kg) (Figure 2A). Interestingly, although 10 mg/kg MPTP did not alter *Hmox1* mRNA levels after 24 hours (blue bar in Figure 2A), it was sufficient to lower the mRNA levels of *Hmox1* after the second injection of MPTP when compared to the 1 mg/kg priming dose (brown bar in Figure 2A).

To investigate the duration of refractoriness, we injected C57BL/6J mice on day 0 with a single dose of MPTP (40 mg/kg) or saline. Some animals were sacrificed after 24 h to assess the effect of the first MPTP injection (green bar in Figure 2B). The remaining animals were injected a second time with MPTP (40 mg/kg) (brown bars in Figure 2B) or saline (blue bars in Figure 2B) on either day 7, 14, 21 or 28 and were sacrificed 24 h after the second injection. Striatal *Hmox1* mRNA levels were elevated 24 h after the priming dose of MPTP (40 mg/kg) (green bar in Figure 2B). When compared to mice that received MPTP followed by saline, mice that received two injections of MPTP did not have statistically significantly elevated levels of *Hmox1* at any time point (compare blue and brown bars in Figure 2B). Although there was a trend towards recovery of a transcriptional response to MPTP starting at day 22, it did not reach statistical significance, indicating that the refractory period elicited by MPTP lasts at least 3–4 weeks.

3.3 mRNA levels for *Cdkn1a* are elevated after multiple injections of MPTP

Shortly (5 h) after administration of MPTP, the cyclin-dependent kinase inhibitor 1a (*Cdkn1a* a.k.a. *p21^{Waf1/Cip1}*) mRNA level is elevated in striatum. This is only one of the 88 genes that are elevated starting at early time points after MPTP injections in both MPTP-sensitive (C57BL/6J) and MPTP-resistant (SWR) mice (Pattarini et al., 2008). Therefore we assessed *Cdkn1a* mRNA levels in striatum by qRT-PCR. C57BL/6J mice were injected with MPTP (40 mg/kg) or saline as control at day 0, 1, 2 and 3 (0, 24, 48 and 72 h, respectively; Figure 3) and sacrificed 5 and 24 h after each injection. *Cdkn1a* mRNA levels were elevated 5 hours after each injection, and returned to baseline 24 h later (Figure 3). Thus in contrast to *Hmox1*, MPTP does not induce a refractory period for *Cdkn1a*.

3.4 Genome-wide transcriptional responses after pre-exposure to MPTP

We used a genome-wide approach to determine whether genes other than *Hmox1* were refractory to repeated administration of MPTP. C57BL/6J mice were “primed” with a single dose of either MPTP (40 mg/kg) (M) or saline (S) on day 0. After 9 days, saline or MPTP-primed mice were injected with either saline or MPTP following the acute regimen (4×20 mg/kg every 2 hours). Mice were sacrificed 5, 24 and 72 h after the first injection. Control animals (S-S) were injected with saline on both days. Animals in the M-S group received a single dose of MPTP (40 mg/kg) on day 0 and saline on day 9 to identify genes that were

still elevated 9 days after the priming dose. Animals of the S-M group were treated with saline on day 0 and MPTP on day 9 to assess the transcriptional response after acute administration of MPTP. Mice in the M-M group were injected with MPTP (40 mg/kg) on day 0 and again with the acute MPTP regimen on day 9 to assess the effect of pre-exposure to MPTP.

Following isolation of total RNA from striatum, Affymetrix analysis was performed (Figure 4). Using hierarchical cluster analysis, a direct comparison of animals in categories M-S and S-S indicated that with few exceptions (Table 3), the transcriptional response elicited by the MPTP injection on day 0 was no longer evident by day 9 (Figure 4). The S-M group revealed relatively unique gene sets in the early (5 h), intermediate (24 h) and late (72 h) phases, that were qualitatively similar to those previously reported (Pattarini et al., 2008). Animals primed with MPTP (40 mg/kg) before receiving an acute MPTP treatment (M-M group) did not differ in the early (5 h) response (compare 5 hour S-M and M-M groups in Figure 4). In contrast, mice of the M-M group lacked almost completely the intermediate and late transcriptional phases (compare 24 and 72 h S-M and M-M groups in Figure 4). Indeed only 10 genes in the intermediate phase were not affected by pre-exposure to MPTP (Table 4).

To confirm the gene chip array results we performed qRT-PCR for representative genes of the early (5 h) and intermediate (24 h) phases (Figures 5 and 6, respectively). Animals for this experiment were treated exactly the same as those used in the microarray experiments. Levels of mRNA for all genes in M-S treated animals were indistinguishable from those that received only saline (S-S) (compare white and brown bars in Figures 5 and 6). Some genes, such as *Cdkn1a* and growth arrest and DNA-damage-inducible, beta (*Gadd45b*), did not differ between S-M and M-M groups whereas other genes, such as nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (*Nfkb1a*) and FBJ murine osteosarcoma viral oncogene homolog B (*Fosb*), displayed a blunted although still significant response after pre-exposure to MPTP (Figure 5). Nuclear receptor subfamily 4, group A, member 1 (*Nr4a1*) was the only early response gene that showed significant refractoriness (Figure 5). For the intermediate response, we assessed the mRNA level profiles of *Hmox1*, tumor necrosis factor receptor superfamily, member 12A (*Tnfrsf12a*), endothelial differentiation, sphingolipid G-protein-coupled receptor 3 (*Edg3*), PDZ and LIM domain 4 (*Pdlim4*), heparin-binding EGF-like growth factor (*Hbegf*) tumor necrosis factor alpha (*Tnfa*) and monocyte chemoattractant protein-1/chemokine (C-C motif) ligand 2 (*Mcp1/Ccl2*). *Hmox1*, *Tnfrsf12a*, *Edg3* and *Tnfa* were profoundly refractory (Figure 6). Elevation of *Pdlim4* mRNA was attenuated but not abolished. *Mcp1/Ccl2* is an interesting gene as it is elevated during both the early and intermediate phases (Pattarini et al, 2007 and Figure 6), and whereas the intermediate response is refractory, the early response is only reduced following pre-exposure to MPTP.

3.5 Effect of MPTP pre-exposure in MPTP-sensitive (C57BL/6J) and MPTP-resistant (SWR) strains of mice

We previously described that the striatal transcriptional response in the MPTP-sensitive C57BL/6J strain of mice differs from that of the MPTP-resistant, SWR strain (Pattarini et al., 2007, Pattarini et al., 2008). Here we assessed if pre-exposure to MPTP modified the transcriptional response in MPTP-resistant SWR mice. We injected both SWR and C57BL/6J mice with MPTP (30 mg/kg) or saline, on day 0 and day 9, and sacrificed animals 5, 24 and 72 hours after the second injection. The dosage used in this experiments was chosen to minimize death of SWR mice due to peripheral toxicity and has been shown previously to elicit transcriptional responses in both strains (Pattarini et al., 2007, Pattarini et al., 2008). qRT-PCR was performed for genes of the early phase (*Cdkn1a*, *Fosb*, *Gadd45b* and *Nfkb1a*; Figure 7) and intermediate phase (*Hmox1*, *Edg3*, *Tnfrsf12a*, and *Pdlim4*; Figure 8). mRNA

levels in animals treated with MPTP on day 0 followed by saline (M-S) did not differ from mice injected with saline on both days (S-S, Figures 7 and 8). Animals in the saline-MPTP group, displayed a qualitatively similar response to that previously reported (Pattarini et al., 2008). In particular, the early response occurred in both strains of mice with some genes induced to similar levels (*Cdkn1a* and *Nfkb1a*) and others at even higher levels in SWR compared to C57BL/6J mice (*Fosb* and *Gadd45b*) (Figure 7). In contrast, the intermediate response was much reduced (*Tnfrsf12a* and *Pdlim4*) or absent (*Hmox1* and *Edg3*) in the S-M group of the MPTP-resistant, SWR strain (white bars in Figure 8). Pre-exposure to MPTP (M-M group) did not alter gene expression in the intermediate phase in SWR mice (Figure 8); i.e. priming doses of MPTP do not convert the transcriptional profile of SWR mice to one resembling an MPTP sensitive strain. As anticipated in C57BL/6J mice of the M-M group, the early phase was largely unaffected (Figure 7) whereas the intermediate response was absent or attenuated (Figure 8).

3.6 Paraquat and MPTP trigger distinct and refractory transcriptional responses but do not elicit cross-attenuation

Several environmental agents have been associated with an elevated risk of PD (Moretto and Colosio, 2011). Therefore it is conceivable that exposure to one toxin may alter the response to future encounters with other toxins. In addition, distinct agents may trigger different transcriptional responses. Paraquat (PQ) is structurally similar to MPP⁺ (Brooks et al., 1999) and can elicit loss of DAergic neurons of the SNpc in mice (McCormack et al., 2005). Therefore, we assessed whether exposure to PQ triggers a transcriptional response similar to MPTP and whether pre-exposure to PQ modifies the response to PQ or MPTP and vice versa. C57BL/6J mice were treated on day 0 with one dose of saline, MPTP (40 mg/kg) or paraquat (40 mg/kg). Four days later mice received injections of saline, MPTP or PQ and were sacrificed 24 hour later. Total striatal RNA was isolated and used for Affymetrix® array analysis as described.

In animals pretreated with saline, PQ induced a transcriptional response that was limited to relatively few genes (66 probe sets corresponding to 56 genes listed in Table 5) that by and large did not overlap with those induced by MPTP (compare columns S-M and S-P in Figure 9 and see Table 6 for a list of 5 genes modulated by both toxicants). This was in striking contrast to the extensive changes elicited by MPTP in the S-M group (column S-M in Figure 4 and column S-M in Figure 9). Pre-exposure to priming doses of MPTP again eliminated or attenuated the intermediate response (M-M group, Figure 9). Interestingly, although PQ induced a small group of genes (Table 5), this response was blunted or absent in PQ pre-treated mice (P-P group in Figure 9). In contrast, mice primed with one toxin, and then challenged with the other one, did not show any sign of refractoriness (compare columns P-M and M-P versus S-M and S-P columns in Figure 9). Therefore, PQ and MPTP elicit distinct striatal transcriptional responses and pre-exposure to one toxin can only alter the response to a subsequent challenge with the same toxin but they do not elicit cross-attenuation.

In Figure 9, arrays from animals receiving the same treatment (e.g. S-M) were grouped together, and these groups were compared versus the S-S group using t-tests. This method identified 245 probe sets that differed between the S-S group and the other groups. To identify potential similarities among treatments not revealed with the previous method, we analyzed data using a different approach. First we compared all treatment groups to each other using ANOVA instead of t-tests to identify probe sets differentially expressed among treatments. This method identified 1171 probe sets that differed significantly. The 1171 probe sets on all the arrays were then compared to each other based on their similarity to one another, rather than the experimental condition. This hierarchical clustering approach revealed three major subgroups (Figure 10): cluster 1 includes mice pre-exposed to either

PQ or saline, and then treated with MPTP on day 9; cluster 2 includes animals treated with the same toxins both on day 0 and day 9 (MM and PP), and mice injected only with saline (SS); cluster 3 includes arrays from mice pre-exposed to either MPTP or saline, and then treated with PQ on day 9. This served to reinforce the notion that MPTP and PQ induce distinct gene expression profile changes and that mice treated twice with the same toxin were indistinguishable from saline-treated animals.

3.7 Gene ontology (GO) and gene set enrichment analysis (GSEA)

Although PQ and MPTP largely affected expression of distinct sets of genes, it is possible that they still influence similar biological processes. Therefore, to compare the responses to MPTP and PQ, gene ontology (GO) and gene set enrichment analyses (GSEA) were performed.

GO analysis revealed several biological and molecular functions (12 GO terms) regulated after MPTP treatment (Table 7), including response to wounding (GO:0009611), protein dimerization activity (GO:0046983), and inflammatory-related responses such as inflammatory response (GO:0006954), acute inflammatory response (GO:0002526), and activation of immune response (GO:0002253). These results are consistent with both the available literature on MPTP (Dauer and Przedborski, 2003, Bove et al., 2005, Smeyne and Jackson-Lewis, 2005, Wood-Kaczmar et al., 2006) and our previous results (Pattarini et al., 2007, Pattarini et al., 2008). GO analysis revealed only protein dimerization activity (GO:0046983), to be regulated after PQ administration (Table 7). This molecular process is also affected by MPTP, although genes regulated by PQ (*Fos*, *Fos12*, *Junb*, *Npas4*, *Nr4a1*, *Vegfa*, *Zbtb16*) differ from those modulated by MPTP (*C1qb*, *Cav2*, *Cebpb*, *Cebpd*, *Jun*, *Npas4*, *Pycard*, *S100a16*, *Sim1*, *Stat3*, *Vwf*, *Wwtr1*).

GSEA confirmed the GO results in that MPTP influenced many biological processes (Table 8), whereas we were unable to identify any over-expressed function in PQ treated mice. Table 8 shows the most prominent biological processes and molecular functions that are significantly higher in MPTP-treated mice. These results are comparable to those reported by several authors (Dauer and Przedborski, 2003, Bove et al., 2005, Smeyne and Jackson-Lewis, 2005, Wood-Kaczmar et al., 2006, Pattarini et al., 2007, Pattarini et al., 2008). Similar to GO analysis, we were unable to identify any biological process or molecular function that is more represented in PQ-treated mice.

3.8 Temporal correlation between DA depletion and refractoriness

DA, but not other catecholamines, induces transcription of *Hmox1* in glial cells (Schmidt et al., 1999). Since DA is known to be rapidly released from SNpc nerve endings as an acute response to MPTP intoxication (Chang and Ramirez, 1986), this catecholamine or some unidentified substance emanating from DAergic nerve endings may trigger part of the transcriptional response. If true, the refractory period might be coincident with the period of DA depletion in striatum. Therefore, we assessed the effect of MPTP on striatal DA levels. Animals were administered a single injection of MPTP (40 mg/kg) and the striatal concentrations of DA and its two main metabolites, 3-4-dihydroxyphenylacetic acid (DOPAC) and homovanillic acid (HVA) assessed at 1, 2, 24 and 72 hours and 1, 2, 3, 4, 5 and 6 weeks following treatment. The concentration of DA in the striatum was substantially reduced 2 h after MPTP administration, and transiently recovered to basal levels at 24 h before decreasing again at day 3, and remaining low for up to 6 weeks (Figure 11). The ratio between DA concentration and that of its two metabolites HVA and DOPAC, confirmed that the turnover of DA in the striatum is significantly altered for a prolonged period of time. At the longest time point there was a suggestion that DA levels were beginning to recover.

Nevertheless, the long-term refractory period occurs concomitantly with a reduction of striatal DA.

4 Discussion

Although the etiology of sporadic PD remains to be fully elucidated, it is thought to involve both environmental agents and predisposing genetic factors (Vance et al., 2010). The current study provides an experimental platform that can potentially lead to a better understanding of both aspects of this issue.

Many epidemiological studies have shown a link between exposure to various pesticides and herbicides and the risk of developing PD (Alavanja et al., 2004, Landrigan et al., 2005, Gatto et al., 2009). Our data show an unexpected relationship between the frequency of exposure to a neurotoxin and the molecular response it elicits, suggesting this parameter might play a role in the etiology of PD. The different regimens used here to induce experimental Parkinsonism may model different exposure profiles in man. The mode of administration of MPTP can elicit different pathologies both in terms of their scope and timing. Sonsalla and Heikkila (1986) were the first to report that dose and dose intervals influenced MPTP neurotoxicity. More recently, differences in striatal glutamate synaptic regulation were reported between the acute and subchronic MPTP regimens (Robinson et al., 2003). In addition to the scope of injury, the mechanism of SNpc DAergic neuron death induced by MPTP has also been shown to be dependent on MPTP administration schedule: apoptosis is implicated in the subchronic model (Tatton and Kish, 1997), whereas necrosis was described following the acute regimen (Jackson-Lewis et al., 1995). Finally, although controversial (Alvarez-Fischer et al., 2008), Fornai and colleagues reported the presence of Lewy body-like aggregates in animals continuously perfused with MPTP (30 mg/kg/day) for a prolonged period of time (Fornai et al., 2005). The data presented here suggest that some of these differences could be the result of distinct transcriptional responses triggered in the acute and subchronic MPTP regimens and further that prior exposure to a toxin can profoundly influence the transcriptional response to a second encounter with the same toxin.

A major question raised by this study is the relevance of the various phases of gene expression to pathological outcome. SWR mice lack the delayed transcriptional responses to MPTP and do not exhibit degeneration (Figure 8 and Pattarini et al., 2007, Pattarini et al., 2008); i.e. the presence of the delayed (but not acute) response is predictive of degeneration, but it is unclear whether this is cause or effect. As shown in Figure 4, C57BL/6J mice primed with MPTP show an almost identical gene expression profile following subsequent challenges with MPTP to the MPTP-resistant SWR mice. Therefore, if the expression of one or more genes in the delayed response directly contributes to neuronal demise, then priming with MPTP might render normally MPTP-sensitive strains resistant to subsequent challenges with the same toxin. Alternatively, genes within the intermediate and late response groups may contribute to neuroprotection, in which case priming may further sensitize the animals to subsequent challenges with MPTP. Finally, the delayed phases of gene expression may represent the response to damage rather than being the cause of the damage and therefore be a reflection of the underlying biological processes that accompany the response to an MPTP lesion. Future detailed studies of cell loss in the SNpc in these dosing scenarios will help shed light on the pathological relevance of the various transcriptional alterations triggered by MPTP.

Another key finding in the present study is that whereas MPTP and paraquat both elicit parkinsonian-like responses in mice (Dauer and Przedborski, 2003, Smeyne and Jackson-Lewis, 2005, Tanner et al., 2011) they trigger quite distinct transcriptional responses (Figure 9). Epidemiological studies have linked rural living (Rajput et al., 1987) and exposure to

various pesticides and herbicides including rotenone, paraquat and manganese ethylene bisdithiocarbamate (Maneb) to risk for PD (Alavanja et al., 2004, Landrigan et al., 2005). Additionally, there are reported interactions between PD-associated toxins in animal models. For example, the extent of the lesion achieved by concomitant administration of PQ and Maneb is considerably larger than that obtained with either toxin alone (Thiruchelvam et al., 2003). Priming mice with lipopolysaccharide (LPS), to cause a persistent inflammatory response, also increased the neurodegenerative effects of PQ (Mangano and Hayley, 2009) as did pre-treatment with MPTP (Shepherd et al., 2006). The current data show that MPTP, but not PQ, triggers transcriptional changes with the hallmark of an inflammatory response and this inflammation could underlie the synergy reported between these agents. That the two mechanisms of action are distinct is further underlined by our finding that priming with MPTP or PQ makes the transcriptional response to subsequent exposure to the same toxin refractory but it does not result in cross-attenuation. The lack of overlapping biological responses and molecular functions revealed by GO and GSEA analysis further highlights the differences between the toxicity of MPTP and PQ. It will be important in the future to assess the transcriptional profiles elicited by different toxins that cause PD-like lesions in mice to establish whether they all have different repertoires of gene expression changes or whether they fall into distinct categories, suggesting common mechanisms of action. If we can define toxin mechanisms by gene expression profiling it will then be interesting to establish the gene expression patterns triggered by co-administration of agents from different categories and relate this to pathological outcomes.

Another key question is the molecular and cellular basis of the refractoriness induced by MPTP. Genes within the acute phase have an over-representation of immediate-early transcription factors such as Fosb and c-Jun (Pattarini et al., 2008). Therefore, it was postulated that these gene products contributed to the regulation of genes in the delayed response phases in a temporal transcriptional cascade (Pattarini et al., 2008). However, the present data do not support this hypothesis. MPTP triggers a robust acute transcriptional response in both sensitive and resistant strains of mice as well as in a sensitive strain of mice primed with MPTP, yet the delayed responses are greatly attenuated or absent in the latter two circumstances (Figures 4 and 6). Therefore, it is more likely that the acute response underlies a global, stereotyped cellular stress reaction to MPTP intoxication rather than MPTP priming causing an uncoupling of transcriptional regulation of delayed genes by acute phase genes.

The duration of the refractory period elicited by a single administration of MPTP (Figures 1, 2 and 4) is coincident with the period of depleted striatal dopamine content in this model (Figure 11). Dopamine, but not other catecholamines, triggers expression of Hmox1 in cultured astrocytes (Schmidt et al., 1999), and striatal astrocytes are the source of elevated Hmox1 in the MPTP model (Fernandez-Gonzales et al., 2000). Therefore, dopamine dumping from SNpc DAergic nerve endings in the striatum as a result of MPTP intoxication might play a role as the initial trigger for the delayed transcriptional responses. However, once dopamine pools are depleted, subsequent treatments with MPTP cannot trigger a second delayed response. Although further studies are required to prove this interpretation, it would also fit the observation that the delayed response is selective for the striatum, in that it receives extensive DA-ergic innervation, whereas other brain areas with lower DA content only show an acute response to MPTP (Pattarini et al., 2008). The fact that the acute response is still intact in MPTP-primed mice (Figure 4) and occurs globally in brain (Pattarini et al., 2008) indicates that this rapid transcriptional response is likely triggered by something other than dopamine.

The intermediate response has an over-representation of genes involved in inflammation (Table 7 and 8) such as cytokines, chemokines and their receptors (Pattarini et al., 2007,

Pattarini et al., 2008). This is consistent with these alterations being a reflection of the inflammatory response known to be elicited in the striatum of MPTP treated mice (Pattarini et al., 2007, Pattarini et al., 2008) and also observed in PD patients, as well as humans intoxicated with MPTP (Langston et al., 1999, Lee et al., 2009). Therefore, the refractoriness of the delayed responses following priming may be a result of long-term negative feedback mechanisms limiting the re-initiation of an inflammatory response. This might occur through the production of anti-inflammatory cytokines and/or a shift in the functional status of cells in the brain that mediate and mount such responses. Nevertheless, the finding that exposure to a neurotoxin, even at sub-clinical doses, can have a long lasting effect on the ability of the mouse to mount an inflammatory response to a much higher level of toxin, has significant implications for our understanding of the role of environmental toxin exposure to the genesis of PD.

Genetic predisposing factors have been linked to the etiology of sporadic PD (Gao and Hong, 2011). The MPTP model is potentially useful to identify these predisposing genes as susceptibility to MPTP has a genetic component (Giovanni et al., 1991, 1994, Miller et al., 1998, Hamre et al., 1999). The present (Figure 4) and a previous gene expression study of MPTP-resistant SWR mice (Pattarini et al., 2008) demonstrate an attenuated intermediate and late transcriptional response after administration of MPTP. Strains of mice have also been generated that harbor genetic mutations implicated in familial and sporadic PD (Meredith et al., 2008). It will now be interesting to assess both the basal and MPTP-treated transcriptional profiles of these mice to determine if specific PD-associated mutations influence particular genes, or phases of the response. This may then point to additional cellular and molecular targets for therapeutic intervention in PD.

Acknowledgments

The authors would like to thank the Hartwell Center for Bioinformatics and Biotechnology for the synthesis of qRT-PCR primers and probes, sequencing and for processing samples for Affymetrix GeneChip technology. This work was supported in part by the National Institutes of Health Cancer Center CORE Grant CA 21765, the American Lebanese Syrian Associated Charities (ALSAC) and the National Institutes of Health grants R01-NS042828 and R01-ES010772 to J.I.M and R01-NS39006 to R.J.S.

Abbreviations

<i>Cdkn1a</i>	cyclin-dependent kinase inhibitor 1a
DA	dopamine
DOPAC	3–4-dihydroxyphenylacetic acid
<i>Edg3</i>	endothelial differentiation, sphingolipid G-protein-coupled receptor 3
<i>Fosb</i>	FBJ murine osteosarcoma viral oncogene homolog B
<i>Gadd45b</i>	growth arrest and DNA-damage-inducible, beta
<i>Hbegf</i>	heparin-binding EGF-like growth factor
<i>Hmox1</i>	heme oxygenase 1
HVA	homovanillic acid
<i>Mcp1/Ccl2</i>	monocyte chemoattractant protein-1/chemokine (C-C motif) ligand 2
MPTP	1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine
<i>Nfkb1a</i>	nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor alpha

<i>Nr4a1</i>	Nuclear receptor subfamily 4, group A, member 1
PD	Parkinson's disease
<i>Pdlim4</i>	PDZ and LIM domain 4
qRT-PCR	quantitative RT-PCR
SNpc	substantia nigra pars compacta
SWR	Swiss Webster mice
<i>Tnfa</i>	tumor necrosis factor alpha
<i>Tnfrsf12a</i>	tumor necrosis factor receptor superfamily, member 12A
MAO-B	monoamine oxidase B
PQ	paraquat
MPP+	1-methyl-4-phenylpyridinium
IP	intraperitoneal
CDS	coding sequence
LPS	lipopolysaccharide

References

- Acworth, IN.; Cunningham, ML. The measurement of monoamine neurotransmitters in microdialysis perfusates using HPLC-ECD. In: Harry, J.; Tilson, HA., editors. *Methods in Molecular Medicine*. Vol. vol. 22. New Jersey: Humana Press; 1999. p. 219-236.
- Alavanja MC, Hoppin JA, Kamel F. Health effects of chronic pesticide exposure: cancer and neurotoxicity. *Annu Rev Public Health*. 2004; 25:155–197. [PubMed: 15015917]
- Alvarez-Fischer D, Guerreiro S, Hunot S, Saurini F, Marien M, Sokoloff P, Hirsch EC, Hartmann A, Michel PP. Modelling Parkinson-like neurodegeneration via osmotic minipump delivery of MPTP and probenecid. *J Neurochem*. 2008; 107:701–711. [PubMed: 18761710]
- Ballard PA, Tetrad JW, Langston JW. Permanent human parkinsonism due to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP): seven cases. *Neurology*. 1985; 35:949–956. [PubMed: 3874373]
- Barlow BK, Cory-Slechta DA, Richfield EK, Thiruchelvam M. The gestational environment and Parkinson's disease: evidence for neurodevelopmental origins of a neurodegenerative disorder. *Reprod Toxicol*. 2007; 23:457–470. [PubMed: 17350799]
- Bazzu G, Calia G, Puggioni G, Spissu Y, Rocchitta G, Debetto P, Grigoletto J, Zusso M, Migheli R, Serra PA, Desole MS, Miele E. alpha-Synuclein- and MPTP-generated rodent models of Parkinson's disease and the study of extracellular striatal dopamine dynamics: a microdialysis approach. *CNS Neurol Disord Drug Targets*. 2010; 9:482–490. [PubMed: 20522009]
- Benmoyal-Segal L, Soreq H. Gene-environment interactions in sporadic Parkinson's disease. *J Neurochem*. 2006; 97:1740–1755. [PubMed: 16805780]
- Bove J, Prou D, Perier C, Przedborski S. Toxin-induced models of Parkinson's disease. *NeuroRx*. 2005; 2:484–494. [PubMed: 16389312]
- Boyd JD, Jang H, Shepherd KR, Faherty C, Slack S, Jiao Y, Smeyne RJ. Response to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) differs in mouse strains and reveals a divergence in JNK signaling and COX-2 induction prior to loss of neurons in the substantia nigra pars compacta. *Brain Res*. 2007; 1175:107–116. [PubMed: 17884023]
- Brooks AI, Chadwick CA, Gelbard HA, Cory-Slechta DA, Federoff HJ. Paraquat elicited neurobehavioral syndrome caused by dopaminergic neuron loss. *Brain Res*. 1999; 823:1–10. [PubMed: 10095006]

- Brooks DJ. Detection of preclinical Parkinson's disease with PET. *Geriatrics*. 1991; 46(Suppl 1):25–30. [PubMed: 1909983]
- Brown TP, Rumsby PC, Capleton AC, Rushton L, Levy LS. Pesticides and Parkinson's disease--is there a link? *Environ Health Perspect*. 2006; 114:156–164. [PubMed: 16451848]
- Chang GD, Ramirez VD. The mechanism of action of MPTP and MPP+ on endogenous dopamine release from the rat corpus striatum superfused in vitro. *Brain Res*. 1986; 368:134–140. [PubMed: 3485463]
- Costa J, Lunet N, Santos C, Santos J, Vaz-Carneiro A. Caffeine exposure and the risk of Parkinson's disease: a systematic review and meta-analysis of observational studies. *J Alzheimers Dis*. 2010; 20(Suppl 1):S221–S238. [PubMed: 20182023]
- Dauer W, Przedborski S. Parkinson's disease: mechanisms and models. *Neuron*. 2003; 39:889–909. [PubMed: 12971891]
- Farrer MJ. Genetics of Parkinson disease: paradigm shifts and future prospects. *Nat Reviews*. 2006; 7:306–318.
- Fernandez-Gonzales A, Perez-Otano I, Morgan JI. MPTP selectively induces Heme Oxygenase-1 expression in striatal astrocytes. *Eur J Neurosci*. 2000; 12:1573–1583. [PubMed: 10792435]
- Fornai F, Schluter OM, Lenzi P, Gesi M, Ruffoli R, Ferrucci M, Lazzeri G, Busceti CL, Pontarelli F, Battaglia G, Pellegrini A, Nicoletti F, Ruggieri S, Paparelli A, Sudhof TC. Parkinson-like syndrome induced by continuous MPTP infusion: convergent roles of the ubiquitin-proteasome system and alpha-synuclein. *Proc Natl Acad Sci U S A*. 2005; 102:3413–3418. [PubMed: 15716361]
- Gao HM, Hong JS. Gene-environment interactions: key to unraveling the mystery of Parkinson's disease. *Prog Neurobiol*. 2011; 94:1–19. [PubMed: 21439347]
- Gatto NM, Cockburn M, Bronstein J, Manthripragada AD, Ritz B. Well-water consumption and Parkinson's disease in rural California. *Environ Health Perspect*. 2009; 117:1912–1918. [PubMed: 20049211]
- German DC, Nelson EL, Liang CL, Speciale SG, Sinton CM, Sonsalla PK. The neurotoxin MPTP causes degeneration of specific nucleus A8, A9 and A10 dopaminergic neurons in the mouse. *Neurodegeneration*. 1996; 5:299–312. [PubMed: 9117541]
- Giovanni A, Sieber BA, Heikkila RE, Sonsalla PK. Correlation between the neostriatal content of the 1-methyl-4-phenylpyridinium species and dopaminergic neurotoxicity following 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine administration to several strains of mice. *J Pharmacol Exp Ther*. 1991; 257:691–697. [PubMed: 2033514]
- Giovanni A, Sieber BA, Heikkila RE, Sonsalla PK. Studies on species sensitivity to the dopaminergic neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. Part 1: Systemic administration. *J Pharmacol Exp Ther*. 1994; 270:1000–1007. [PubMed: 7932147]
- Hamre K, Tharp R, Poon K, Xiong X, Smeyne RJ. Differential strain susceptibility following 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) administration acts in an autosomal dominant fashion: quantitative analysis in seven strains of *Mus musculus*. *Brain Res Mol Brain Res*. 1999; 828:91–103.
- Henry J, Smeyne RJ, Jang H, Miller B, Okun MS. Parkinsonism and neurological manifestations of influenza throughout the 20th and 21st centuries. *Parkinsonism Relat Disord*. 2010
- Huang Y, Cheung L, Rowe D, Halliday G. Genetic contributions to Parkinson's disease. *Brain Res Mol Brain Res*. 2004; 46:44–70.
- Jackson-Lewis V, Jakowec M, Burke RE, Przedborski S. Time course and morphology of dopaminergic neuronal death caused by the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Neurodegeneration*. 1995; 4:257–269. [PubMed: 8581558]
- Jang H, Boltz D, Sturm-Ramirez K, Shepherd KR, Jiao Y, Webster R, Smeyne RJ. Highly pathogenic H5N1 influenza virus can enter the central nervous system and induce neuroinflammation and neurodegeneration. *Proc Natl Acad Sci U S A*. 2009; 106:14063–14068. [PubMed: 19667183]
- Koller W, Vetere-Overfield B, Gray C, Alexander C, Chin T, Dolezal J, Hassanein R, Tanner C. Environmental risk factors in Parkinson's disease. *Neurology*. 1990; 40:1218–1221. [PubMed: 2381528]

- Landrigan PJ, Sonawane B, Butler RN, Trasande L, Callan R, Droller D. Early environmental origins of neurodegenerative disease in later life. *Environ Health Perspect.* 2005; 113:1230–1233. [PubMed: 16140633]
- Langston JW, Forno LS, Tetrad J, Reeves AG, Kaplan JA, Karluk D. Evidence of active nerve cell degeneration in the substantia nigra of humans years after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine exposure. *Ann Neurol.* 1999; 46:598–605. [PubMed: 10514096]
- Lee JK, Tran T, Tansey MG. Neuroinflammation in Parkinson's disease. *J Neuroimmune Pharmacol.* 2009; 4:419–429. [PubMed: 19821032]
- Mangano EN, Hayley S. Inflammatory priming of the substantia nigra influences the impact of later paraquat exposure: Neuroimmune sensitization of neurodegeneration. *Neurobiol Aging.* 2009; 30:1361–1378. [PubMed: 18187236]
- McCormack AL, Atienza JG, Johnston LC, Andersen JK, Vu S, Di Monte DA. Role of oxidative stress in paraquat-induced dopaminergic cell degeneration. *J Neurochem.* 2005; 93:1030–1037. [PubMed: 15857406]
- Meredith GE, Sonsalla PK, Chesselet MF. Animal models of Parkinson's disease progression. *Acta Neuropathol.* 2008; 115:385–398. [PubMed: 18273623]
- Miller DB, Ali SF, O'Callaghan JP, Laws SC. The impact of gender and estrogen on striatal dopaminergic neurotoxicity. *Ann N Y Acad Sci.* 1998; 844:153–165. [PubMed: 9668673]
- Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstrale M, Laurila E, Houstis N, Daly MJ, Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES, Hirschhorn JN, Altshuler D, Groop LC. PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet.* 2003; 34:267–273. [PubMed: 12808457]
- Moretto A, Colosio C. Biochemical and toxicological evidence of neurological effects of pesticides: the example of Parkinson's disease. *Neurotoxicology.* 2011; 32:383–391. [PubMed: 21402100]
- Nagai Y, Obayashi S, Ando K, Inaji M, Maeda J, Okauchi T, Ito H, Suhara T. Progressive changes of pre- and post-synaptic dopaminergic biomarkers in conscious MPTP-treated cynomolgus monkeys measured by positron emission tomography. *Synapse.* 2007; 61:809–819. [PubMed: 17598150]
- Pattarini R, Rong Y, Qu C, Morgan JI. Distinct mechanisms of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine resistance revealed by transcriptome mapping in mouse striatum. *Neuroscience.* 2008; 155:1174–1194. [PubMed: 18675323]
- Pattarini R, Smeyne RJ, Morgan JI. Temporal mRNA profiles of inflammatory mediators in the murine 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine model of Parkinson's disease. *Neuroscience.* 2007; 145:654–668. [PubMed: 17258864]
- Petroske E, Meredith GE, Callen S, Totterdell S, Lau YS. Mouse model of Parkinsonism: A comparison between subacute MPTP and chronic MPTP/probenecid treatment. *Neuroscience.* 2001; 106:589–601. [PubMed: 11591459]
- Przedborski S, Jackson-Lewis V, Naini AB, Jakowec M, Petzinger G, Miller R, Akram M. The parkinsonian toxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP): a technical review of its utility and safety. *J Neurochem.* 2001; 76:1265–1274. [PubMed: 11238711]
- Rajput AH, Uitti RJ, Stern W, Laverty W, O'Donnell K, O'Donnell D, Yuen WK, Dua A. Geography, drinking water chemistry, pesticides and herbicides and the etiology of Parkinson's disease. *Can J Neurol Sci.* 1987; 14:414–418. [PubMed: 3676917]
- Robinson S, Freeman P, Moore C, Touchon JC, Krentz L, Meshul CK. Acute and subchronic MPTP administration differentially affects striatal glutamate synaptic function. *Exp Neurol.* 2003; 180:74–87. [PubMed: 12668150]
- Schmidt J, Mertz K, Morgan JI. Regulation of heme oxygenase-1 expression by dopamine in cultured C6 glioma and primary astrocytes. *Brain Res Mol Brain Res.* 1999; 73:50–59. [PubMed: 10581397]
- Shepherd KR, Lee ES, Schmued L, Jiao Y, Ali SF, Oriaku ET, Lamango NS, Soliman KF, Charlton CG. The potentiating effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) on paraquat-induced neurochemical and behavioral changes in mice. *Pharmacol Biochem Behav.* 2006; 83:349–359. [PubMed: 16580056]

- Smeyne M, Goloubeva O, Smeyne RJ. Strain-dependent susceptibility to MPTP and MPP+-induced Parkinsonism is determined by glia. *Glia*. 2001; 74:73–80. [PubMed: 11307156]
- Smeyne RJ, Jackson-Lewis V. The MPTP model of Parkinson's disease. *Brain Res Mol Brain Res*. 2005; 134:57–66. [PubMed: 15790530]
- Sonsalla PK, Heikkila RE. The influence of dose and dosing interval on MPTP-induced dopaminergic neurotoxicity in mice. *Eur J Pharmacol*. 1986; 129:339–345. [PubMed: 3490988]
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005; 102:15545–15550. [PubMed: 16199517]
- Sundstrom E, Stromberg I, Tsutsumi T, Olson L, Jonsson G. Studies on the effect of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) on central catecholamine neurons in C57BL/6 mice. Comparison with three other strains of mice. *Brain Res*. 1987; 405:26–38. [PubMed: 2882814]
- Tanaka K, Miyake Y, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M. Active and passive smoking and risk of Parkinson's disease. *Acta Neurol Scand*. 2010; 122:377–382. [PubMed: 20175761]
- Tanner CM, Kamel F, Ross GW, Hoppin JA, Goldman SM, Korell M, Marras C, Bhudhikanok GS, Kasten M, Chade AR, Comyns K, Richards MB, Meng C, Priestley B, Fernandez HH, Cambi F, Umbach DM, Blair A, Sandler DP, Langston JW. Rotenone, paraquat, and Parkinson's disease. *Environ Health Perspect*. 2011; 119:866–872. [PubMed: 21269927]
- Tatton NA, Kish SJ. In situ detection of apoptotic nuclei in the substantia nigra compacta of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated mice using terminal deoxynucleotidyl transferase labelling and acridine orange staining. *Neuroscience*. 1997; 77:1037–1048. [PubMed: 9130785]
- Thiruchelvam M, McCormack A, Richfield EK, Baggs RB, Tank AW, Di Monte DA, Cory-Slechta DA. Age-related irreversible progressive nigrostriatal dopaminergic neurotoxicity in the paraquat and maneb model of the Parkinson's disease phenotype. *Eur J Neurosci*. 2003; 18:589–600. [PubMed: 12911755]
- Vance JM, Ali S, Bradley WG, Singer C, Di Monte DA. Gene-environment interactions in Parkinson's disease and other forms of parkinsonism. *Neurotoxicology*. 2010; 31:598–602. [PubMed: 20430055]
- Vila M, Jackson-Lewis V, Vukosavic S, Djaldetti R, Liberatore G, Offen D, Korsmeyer SJ, Przedborski S. Bax ablation prevents dopaminergic neurodegeneration in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease. *Proc Natl Acad Sci U S A*. 2001; 98:2837–2842. [PubMed: 11226327]
- Weisskopf MG, Weuve J, Nie H, Saint-Hilaire MH, Sudarsky L, Simon DK, Hersh B, Schwartz J, Wright RO, Hu H. Association of cumulative lead exposure with Parkinson's disease. *Environ Health Perspect*. 2010; 118:1609–1613. [PubMed: 20807691]
- Wood-Kaczmar A, Gandhi S, Wood NW. Understanding the molecular causes of Parkinson's disease. *Trends Mol Med*. 2006; 12:521–528. [PubMed: 17027339]

Figure 1. Comparison of mRNA changes of *Hmox1* in the striatum C57BL/6J mice treated with the acute or subchronic MPTP regimens

Panel A: Schematic representation of the different MPTP injection schedules used in this study. Animals treated according to the acute regimen received 4 injections of 20 mg/kg MPTP-HCl (MPTP) spaced 2 hours apart during a single day. The subchronic model consisted of daily injections of 40 mg/kg MPTP for 4 consecutive days. **Panel B:** quantitative assessment of *Hmox1* mRNA levels after a single variable dose of MPTP in the striatum of MPTP-sensitive C57BL/6J mice. Animals were injected at time zero with a single dose of MPTP (10, 20 or 40 mg/kg) or saline as control, and sacrificed at 24 h. Levels of mRNA for *Hmox1* was determined by qRT-PCR using the $\Delta\Delta C_t$ relative expression

method as described in the Materials and Methods section. Data are presented as mean $\pm$ S.E.M. of 4 animals for each condition. Differences versus control (saline, white bar) were analyzed with one-way ANOVA and Bonferroni post-hoc tests (** $P < 0.01$, *** $P < 0.001$). MPTP causes a dose-dependent increase of *Hmox1* mRNA levels starting at the dose of 20 mg/kg. **Panel C:** quantitative assessment of *Hmox1* mRNA levels after MPTP intoxication following the acute regimen in the striatum of MPTP-sensitive C57BL/6J mice. Animals were injected at time zero with 20 mg/kg MPTP according to the schematic reported on top of the bar graph. Control animals were injected with saline (white bar). Animals were sacrificed at 2, 4, 8, 12, 18, 24, 36 and 48 h after the first injection. Animals sacrificed at 2 and 4h received only 1 and 2 injections, respectively. Levels of mRNA for *Hmox1* were determined by qRT-PCR using the absolute quantization methods. Data are presented as mean $\pm$ S.E.M. of 3 animals for each condition. Differences versus control (saline, white bar) were analyzed with one-way ANOVA and Bonferroni post-hoc tests (** $P < 0.01$, *** $P < 0.001$). Following acute administration of MPTP, *Hmox1* mRNA levels increase starting at 18 h and return to baseline at 36 h. **Panel D:** quantitative assessment of *Hmox1* mRNA levels after MPTP intoxication following the subchronic regimen in the striatum of MPTP-sensitive C57BL/6J mice. Animals were injected with 40 mg/kg MPTP according to the schematic reported on top of the bar graph. Control animals were injected with saline (white bar). Animals were sacrificed 24 h after each injection. Mice were sacrificed on day 1, 2, 3, 4 and 8 and received 1, 2, 3, 4 and 5 injections, respectively. Levels of mRNA for *Hmox1* were determined by qRT-PCR using the $\Delta\Delta C_t$ relative expression method. Data are presented as mean $\pm$ S.E.M. of 15 (time point 0), 12 (time points 1, 2 and 3), 7 (time point 4) and 5 (time point 8) animals. Differences versus control (white bar) were analyzed with one-way ANOVA and Bonferroni post-hoc tests (** $P < 0.01$, *** $P < 0.001$).

Figure 2. Dose dependency and time course of the latency period caused by a single injection of MPTP

Panel A: MPTP-sensitive C57BL/6J mice were injected according to the schematic on the top of the bar graph. Animals received a single injection of MPTP (1, 5, 10, 20 or 40 mg/kg) on day 0. Some animals were sacrificed 24 h later (blue bars) while remaining animals were injected a second time with 40 mg/kg MPTP after 7 days and sacrificed after 24 hours (brown bars). Control animals were injected with saline using the same schedules. Levels of mRNA for *Hmox1* were determined by qRT-PCR using the $\Delta\Delta C_t$ relative expression method. Data are presented as mean $\pm$ S.E.M. of 6 to 9 animals for each condition. Differences versus control (Saline) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated with asterisks (***) $P < 0.001$). Differences among treatments were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^A $P < 0.001$). Exposure to 10 mg/kg MPTP does not significantly elevate *Hmox1* mRNA levels (blue bar) but can reduce the response to acute administration of MPTP (brown bar). **Panel B:** MPTP-sensitive C57BL/6J mice were injected according to the schematic on top of the bar graph. Animals received a single injection of 40 mg/kg MPTP or saline as control on day 0. Some animals were sacrificed 24 h later (green and white bars) whereas remaining animals received a second injection of 40 mg/kg MPTP or saline as control either 7, 14, 21 or 28 days later and were sacrificed 24 h after the second injection. Levels of mRNA for *Hmox1* were determined by qRT-PCR using the $\Delta\Delta C_t$ relative expression method. Data are presented as mean $\pm$ S.E.M. of 7 to 27 animals for each condition. Differences versus control (Saline) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated with asterisks (***) $P < 0.001$. Differences versus MPTP (green bar) and between MPTP-Saline and MPTP-MPTP (blue and brown bars, respectively) were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^A $P < 0.001$).

Figure 3. Quantitative assessment of Cdkn1a mRNA changes in the striatum of C57BL/6J mice treated with the subchronic MPTP regimen

MPTP-sensitive C57BL/6J mice were injected according to the schematic shown on the top of the bar graph. Animals were injected daily with a single dose of 40 mg/kg MPTP and sacrificed 5 or 24 hr after each injection. Control animals were injected with saline (white bar). Levels of mRNA for *Cdkn1a* were determined by qRT-PCR using the $\Delta\Delta C_t$ relative expression method. Data are presented as mean $\pm$ S.E.M. of 10 animals for each condition. Differences versus control (white bar) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated with asterisks (*** $P < 0.001$, * $P < 0.05$). Differences among time points were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^A $P < 0.001$, ^B $P < 0.01$).

Figure 4. Hierarchical cluster analysis reveals differences between the early (5 hr) and intermediate (24 hr) MPTP-induced responses in striatum of C57BL/6J mice following a priming dose of MPTP

Animals were injected with either 40 mg/kg MPTP or saline as control on day 0, and injected again after 9 days using the acute MPTP regimen. Mice were sacrificed and striatum harvested at 5 (3 injections), 24 or 72 hours. In this hierarchical cluster analysis each vertical column represents a single mouse treated as indicated on top of the figure and each horizontal row is an individual probe set. Probe sets that are upregulated in treated compared to control mice appear in red, those that are downregulated appear in green. The relative log₂ (ratio) is reflected by the intensity of the color. Whereas the early phase occurs

independent of previous exposure to MPTP, the intermediate and late phases are absent in mice pretreated with MPTP.

Figure 5. Quantitative assessment of striatal mRNA changes of genes of the early phase in the striatum of C57BL/6J mice previously exposed to 40mg/kg MPTP

MPTP sensitive C57BL/6J mice were injected at time zero with a single priming dose of 40 mg/kg MPTP or saline. Mice received a second exposure to either saline or the acute MPTP regimen starting 9 days after the priming injection, and animals were sacrificed and striatum harvested 5 (3 injections), 24 or 72 hours later. Levels of mRNA for candidate genes of the early (*Cdkn1a*, *Nfkb1a*, *Fosb*, *Gadd45b* and *Nr4a1*) response phase were evaluated by qRT-PCR using the absolute quantitative method. Results are expressed as the number of copies of a specific mRNA normalized versus the number of copies of *Gapdh* mRNA. Data are presented as mean $\pm$ S.E.M. of 4 (saline-treated) and 3 (MPTP-treated) animals. Differences

versus control (Saline-Saline) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated with asterisks (** $P < 0.001$, * $P < 0.01$). Differences among treatments were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^B $P < 0.01$, ^C $P < 0.05$).

Figure 6. Quantitative assessment of striatal mRNA changes of genes of the intermediate phase in the striatum of C57BL/6J mice previously exposed to 40 mg/kg MPTP

MPTP-sensitive C57BL/6J mice were injected at time zero with a single priming dose of 40 mg/kg MPTP or saline. Mice received a second exposure to either saline or the acute MPTP regimen starting 9 days after the priming injection, and animals were sacrificed and striatum harvested 5 (3 injections), 24 or 72 hours later. Levels of mRNA for candidate genes of the intermediate (*Hmox1*, *Tnfrsf12a*, *Edg3*, *Pdlim4*, *Hbegf*, *Tnf- α* and *Mcp1/Ccl2*) response phase were evaluated by qRT-PCR using the absolute quantitative method. Results are expressed as the number of copies of a specific mRNA normalized versus the number of copies of *Gapdh* mRNA. Data are presented as mean $\pm$ S.E.M. of 4 (control) and 3 (MPTP-

treated) animals. Differences versus control (Saline-Saline) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated with asterisks (** $P<0.001$, * $P<0.05$). Differences among treatments were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^A $P<0.001$, ^C $P<0.05$).

Figure 7. Comparison of the MPTP-induced refractory state between MPTP-sensitive C57BL/6J and MPTP-resistant SWR strains of mice

C57BL/6J and SWR mice were injected with either 30 mg/kg MPTP or saline on day zero. After 9 days they were injected again with either a single injection of 30 mg/kg MPTP or saline and sacrificed 5, 24 and 72 hour later and striata collected. Levels of mRNA for candidate genes of the early (*Cdkn1a*, *Fosb*, *Gadd45b* and *Nfkb1a*) response phase were evaluated by qRT-PCR. Results are expressed as the number of copies of a specific mRNA normalized versus the number of copies of *Gapdh* mRNA. Data are presented as mean $\pm$ S.E.M. of 5 to 6 animals per condition. Differences versus control (S-S) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated with asterisks

(*** $P < 0.001$). Differences among treatments were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^A $P < 0.001$, ^B $P < 0.01$). Results indicate that mRNA levels for genes of the early response phase were elevated after MPTP injection independent of previous exposures in both strains of mice.

Figure 8. Comparison of the MPTP-induced refractory state between MPTP-sensitive C57BL/6J and MPTP-resistant SWR strains of mice

C57BL/6J and SWR mice were injected with either 30 mg/kg MPTP or saline (control) on day zero. After 9 days they were injected again with either a single injection of 30 mg/kg MPTP or saline and sacrificed 5, 24 and 72 hour later and striata collected. Levels of mRNA for candidate genes of the intermediate (*Hmox1*, *Edg3*, *Tnfrsf12a* and *Pdlim4*) response phase were evaluated by qRT-PCR. Results are expressed as the number of copies of a specific mRNA normalized versus the number of copies of *Gapdh* mRNA. Data are presented as mean $\pm$ S.E.M. of 5 to 6 animals per condition. Differences versus control (S-S) were analyzed by one-way ANOVA followed by Bonferroni post-hoc tests and are indicated

with asterisks (***) $P < 0.001$). Differences among treatments were analyzed with one-way ANOVA and Bonferroni post-hoc tests and indicated with letters (^A $P < 0.001$). Results indicate that mRNA levels for the genes of the intermediate phase were not elevated following MPTP pre-exposure, and SWR mice did not respond differently to MPTP compared to C57BL/6J after priming.

Figure 9. Hierarchical cluster analysis of basal gene expression differences in striata of MPTP-sensitive C57BL/6J mice following pretreatment with either MPTP or paraquat

C57BL/6J mice were injected on day 0 with a single injection of saline, 40 mg/kg MPTP or 40 mg/kg paraquat. Mice were treated again after 4 days with saline, 40 mg/kg MPTP or 40 mg/kg paraquat and sacrificed 24 h later. All possible treatment combinations were tested. Striatal RNA was extracted and hybridized onto the Affymetrix Mouse Genome 430 2.0 array to identify genes with differential expression between treatments. Each vertical column represents a single mouse treated as indicated on top of the figure (a total of 52 animals were used: 9 each were treated with saline only and PQ-MPTP; 7 each with saline-MPTP, MPTP-MPTP, saline-PQ and PQ-PQ; and 6 with MPTP-PQ), and each horizontal

row is an individual probe set. Probe sets that are upregulated in treated compared to control mice appear in red, those that are downregulated appear in green. The relative log₂ (ratio) is reflected by the intensity of the color. Pre-exposure to MPTP and PQ altered subsequent striatal transcriptional response to the same substance but not to the other one.

Figure 10. Unsupervised hierarchical cluster analysis of gene expression profiles in striata of C57BL/6J mice following treatment with MPTP and/or paraquat

C57BL/6J mice were treated on day 0 with a single injection of saline, 40 mg/kg MPTP or 40 mg/kg paraquat. Mice were treated again after 4 days with saline, 40 mg/kg MPTP or 40 mg/kg paraquat and sacrificed 24 h later. All possible treatment combinations were tested. Striatal RNA was extracted and hybridized onto the Affymetrix Mouse Genome 430 2.0 array and probe sets with differential expression between any two treatments identified by ANOVA. These data were then compared to each other, in an unsupervised cluster analysis based on their similarity to one another regardless of the experimental condition. Probe sets that are upregulated appear in red, those that are downregulated appear in green. The relative

log₂ (ratio) is reflected by the intensity of the color. Each vertical column represents a single mouse treated as indicated at the bottom of the column (a total of 52 animals were used: 9 each were treated with saline only and PQ-MPTP; 7 each with saline-MPTP, MPTP-MPTP, saline-PQ and PQ-PQ; and 6 with MPTP-PQ). Each horizontal row represents a probe set identified by ANOVA. Three major clusters were identified: cluster 1 includes mice pre-exposed to either PQ or saline, and then treated with MPTP on day 9; cluster 2 includes animals treated with the same toxins both on day 0 and day 9 (MM and PP), and mice injected only with saline (SS); cluster 3 includes arrays from mice pre-exposed to either MPTP or saline, and then treated with PQ on day 9. This hierarchical cluster analysis highlights the fact that mice treated twice with the same toxin (MM or PP) are most similar to those treated only with saline (SS).

Figure 11. Temporal profile of dopamine (DA), 3,4-dihydroxyphenylacetic acid (DOPAC) and homovanillic acid (HVA) concentrations after a single injection of 40 mg/kg MPTP
C57BL/6J mice were injected on day 0 with a single dose of either saline or 40 mg/kg MPTP and sacrificed 1, 2, 24, 72 h and 1, 2, 3, 4, 5 and 6 weeks later. **Top Panel:** MPTP elicits a rapid depletion of dopamine in striatum within 2 hours of administration that recovers by 24 hours. However, this recovery is transient as dopamine falls to approximately 10% of control levels by 3 days post-treatment and remains significantly reduced through 6 weeks. **Bottom Panel:** Dopamine turnover can be measured by examining the ratio of its metabolites (DOPAC + HVA)/DA. Three days after a single injection of 40mg/kg MPTP there is a significant increase in (DOPAC + HVA)/DA that remains elevated though 6

weeks. Higher (DOPAC + HVA)/DA ratios suggest dopamine displacement (dumping) since higher cytoplasmic dopamine is converted to DOPAC and rapidly released via reverse transport to promote the formation of HVA. Differences versus control (Saline) were analyzed with one-way ANOVA and Bonferroni post-hoc tests (** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$).

Table 1

List of primers and probes for qRT-PCR

Gene Name	mRNA ID #	Forward Primer	Probe	Reverse Primer
<i>Cdkn1a</i>	NM_007669	5'-TTCCGCACAGGAGCAAAACT-3'	5'-Fam-CCGTTGTCTCTTCCGGTCCCGTGG-BHQ-3'	5'-CGGCGCAAAGTGTCACT-3'
<i>Edg3</i>	NM_010101	5'-AACCAGCCAGATGCGC-3'	5'-Fam-TTGCAGAACGAGAGCCTATTTTCAACACTCTTC-BHQ-3'	5'-GCGTGCAGGCCCGAC-3'
<i>Fosb</i>	NM_008036	5'-CCAGAGCCAGGCCCTAGAAAGAC-3'	5'-Fam-TCGCAGAGAGCGGAACAAGC-BHQ-3'	5'-CCGACGGTTCTCTGCACCTTA-3'
<i>Gadd45b</i>	NM_008655	5'-AGGCGGCCAAACTGATGA-3'	5'-Fam-TGTGGACCCCGACAGCGTGG-BHQ-3'	5'-CATCCTCTCTTCTTCGTCTATGG-3'
<i>Gadph</i>	NM_001001303	5'-TGGATCTGACGTGCCGC-3'	5'-Fam-TGGAGAAACCTGCCAAGTATGATGACATCA-BHQ-3'	5'-TGCCTGCTTCACCACTTC-3'
<i>Hbegf</i>	NM_010415	5'-TGCTGCCGTCGGTGATG-3'	5'-Fam-TGAAAGCTCTTCTGGCCGCAGTGTG-BHQ-3'	5'-ACCGTCAACCAACGCG-3'
<i>Hmox1</i>	NM_010442	5'-GTGATGGAGCGTCCACAGC-3'	5'-Fam-CGACAGCATGCCCCAGGATTTGTC-BHQ-3'	5'-TGGTGGCCTCCTTCAAGG-3'
<i>Mcp-1/Ccl2</i>	NM_011333	5'-GAGCATCCACGTGTTGGCT-3'	5'-Fam-AGCCAGATGCAGTTAACGCCCCACT-BHQ-3'	5'-TGGTGAATGAGTAGCAGCAGGT-3';
<i>NFKBia</i>	NM_010907	5'-CAGCTCACGGAGGACGGGA-3'	5'-Fam-CTCTCCTGCACCTTGGCAATCATCC-BHQ-3'	5'-ATGGTCAGCGGCTTCTCTTC-3'
<i>Nr4a1</i>	NM_010444	5'-GGCCACAGGGAGTGGGA-3'	5'-Fam-CCGGCTGGAGATGCCCTGTATTCAAG-BHQ-3'	5'-CGTTGCTGGTGTTCATATTGA-3'
<i>Pdlim4</i>	NM_019417	5'-AGTGCAACGCGCTGCG-3'	5'-Fam-ACGGGATCGTGGGAACCATTTGTCA-BHQ-3'	5'-ATGGTAGAGCTTGTCTCTCGCC-3'
<i>Tnfrsf12a</i>	NM_013749	5'-GCCGCGGAGAGAAAAAG-3'	5'-Fam-TTACTACCCCCATAGAGGAGACTGGTGGAGAG-BHQ-3'	5'-GCCACCTGGGCAGC-3'
<i>Tnf</i>	NM_013693	5'-TCTATGGCCAGACCCCTCAC-3'	5'-Fam-CTCAGATCATCTTCTCAAAATTCGAGTGACAAGC-BHQ-3'	5'-TTGCTACGACGTGGGCTACA-3'

Table 2

List of primers used to prepare the standards for qRT-PCR

Gene name	Enzyme	Forward primer	Enzyme	Reverse primer
<i>Cdkn1a</i>	HindIII	5'- <u>CCCAAGCTTGGG</u> ATGTCCAATCCTGGTGATGT-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> CTTCAGGGTTTTCTCTTGCA-3'
<i>Edg3</i>	EcoRI	5'- <u>CCGGAATTCCGG</u> ATGGCAACCACGCATGCGCA-3'	XhoI	5'- <u>CCGCTCGAGCGG</u> TCACTTCAGAGGACCCCGT-3'
<i>Fosb</i>	BamHI	5'- <u>CGCGGATCCGCG</u> TGAAACCGACAGAGCCTGG-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> GTTCCTTGCGGGTTTGTGTTG-3'
<i>Gadd45b</i>	BamHI	5'- <u>CGCGGATCCGCG</u> ATGACCCTGGAAGAGCTGGT-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> TGGGTCTCAGCGTTCCTCTA-3'
<i>Gapdh</i>	BamHI	5'- <u>CGCGGATCCGCG</u> ATGGTGAAGGTCGGTGTGAA-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> TTCTTACTCCTTGAGGCCA-3'
<i>Hbegf</i>	HindIII	5'- <u>CCCAAGCTTGGG</u> ATGAAGCTGCTGCCGTCGGT-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> ATAGCTCAGGTCCTCCTCAGTGGG-3'
<i>Hmox1</i>	BamHI	5'- <u>CGCGGATCCGCG</u> TAGCCAGTCCGGTGATGGA-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> TGGGGGCCAGTATTGCATTT-3'
<i>Mcp-1/Ccl2</i>	BamHI	5'- <u>CGCGGATCCGCG</u> TCTCTTCTCCACCACCATG-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> GCATCACAGTCCGAGTCACA-3'
<i>NFkBia</i>	HindIII	5'- <u>CCCAAGCTTGGG</u> AGCCATGTTTCAGCCAGCTG-3'	XhoI	5'- <u>CCGCTCGAGCGG</u> TTATAATGTCAGACGCTGGC-3'
<i>Nr4a1</i>	BamHI	5'- <u>CGCGGATCCGCG</u> TGCTAGAAGGACTGCGGAGC-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> GGCTTAAAGGCACATGGGTG-3'
<i>Pdlim4</i>	HindIII	5'- <u>CCCAAGCTTGGG</u> ATGACCACTCGGTGACCCTG-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> CAGCTCAGACAAGTTCCACCT-3'
<i>Tnf-α</i>	BamHI	5'- <u>CGCGGATCCGCG</u> ATGAGCACAGAAAGCATGAT-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> CTTACAGAGCAATGACTCC-3'
<i>Tnfrsf12a</i>	BamHI	5'- <u>CGCGGATCCGCG</u> GCAATCATGGTCCGGGTTG-3'	EcoRI	5'- <u>CCGGAATTCCGG</u> TGAATCACCACCTCGCCCCA-3'

Table 3

Genes with elevated mRNA levels 9 days after MPTP priming.

Probe Set	Gene Title	Gene Symbol	SS	MS	MS/SS
1418021_at	Complement component 4B (Chido blood group)	C4b	802	1193	1.5
1451978_at	Lysyl oxidase-like 1	Lox1l	161	271	1.7
1415897_a_at	Microsomal glutathione S-transferase 1	Mgst1	531	800	1.5
1418674_at	Oncostatin M receptor	Osmr	187	327	1.8
1417346_at	PYD and CARD domain containing	Pycard	149	276	1.9
1428114_at	Solute carrier family 14 (urea transporter), member 1	Slc14a1	137	290	2.1
1423549_at	Solute carrier family 1 (glutamate/neutral amino acid transporter), member 4	Slc1a4	151	278	1.8
1449682_s_at	Tubulin, beta 2b / tubulin, beta 2a, pseudogene 2	Tubb2b/Tubb2a-ps2	784	1177	1.5

Table 4

Genes that are not refractory 24 h after MPTP priming.

Probe Set Name	Gene Title	Gene Symbol	SS	MS	SM	MM	MM/SM
1448318_at	Adipose differentiation related protein	Adfp	88	95	222	184	0.8
1434449_at	Aquaporin 4	Aqp4	1134	1286	1772	1445	0.8
1417381_at	Complement component 1, q subcomponent, alpha polypeptide	C1qa	1651	1553	3041	2450	0.8
1449401_at	Complement component 1, q subcomponent, C chain	C1qc	1241	1328	2036	1788	0.9
1449164_at	CD68 antigen	Cd68	591	601	1059	883	0.8
1436838_x_at	Coactosin-like 1 (Dictyostelium)	Cotl1	375	516	717	646	0.9
1456567_x_at	Granulin	Grn	1202	1203	1921	1721	0.9
1419573_a_at	Lectin, galactose binding, soluble 1	Lgalsl	355	436	602	511	0.9
1418004_a_at	Transmembrane protein 176B	Tmem176b	1015	967	1530	1391	0.9
1450792_at	TYRO protein tyrosine kinase binding protein	Tyrobp	719	904	1243	1063	0.9

Table 5

List of genes modulated in the striatum 24 h after paraquat treatment.

Probe Set Name	Gene Title	Gene Symbol	SS	SP	SS/SP
1418687_at	Activity regulated cytoskeletal-associated protein	Arc	2281	319	0.1 ↓
1452975_at	Alanine-glyoxylate aminotransferase 2-like 1	Agxt2l1	595	2101	3.5 ↑
1417765_a_at	Amylase 1, salivary	Amy1	309	682	2.2 ↑
1444564_at	Apolipoprotein D	Apod	334	640	1.9 ↑
1428352_at	Arrestin domain containing 2	Arrdc2	230	409	1.8 ↑
1419758_at	ATP-binding cassette, sub-family B (MDR/TAP), member 1A	Abcb1a	220	366	1.7 ↑
1416250_at	B-cell translocation gene 2, anti-proliferative	Btg2	566	332	0.6 ↓
1459372_at	Bhlh-PAS type transcription factor NXF	Nxf	389	119	0.3 ↓
1428485_at	Carbonic anhydrase 12	Car12	1106	679	0.6 ↓
1456314_x_at	CDK2 (cyclin-dependent kinase 2)-associated protein 1	Cdk2ap1	1839	1038	0.6 ↓
1435509_x_at	CDK2 (cyclin-dependent kinase 2)-associated protein 1	Cdk2ap1	2090	1194	0.6 ↓
1417574_at	Chemokine (C-X-C motif) ligand 12	Cxcl12	989	586	0.6 ↓
1416332_at	Cold inducible RNA binding protein	Cirbp	815	2105	2.6 ↑
1434369_a_at	Crystallin, alpha B	Cryab	1963	963	0.5 ↓
1452956_a_at	DNA segment, Chr 12, ERATO Doi 647, expressed	D12Erd647e	992	591	0.6 ↓
1437204_a_at	DNA segment, Chr 8, ERATO Doi 325, expressed	D8Erd325e	255	443	1.7 ↑
1446190_at	Double cortin and calcium/calmodulin-dependent protein kinase-like 1	Deamk11	1402	2399	1.7 ↑
1448830_at	Dual specificity phosphatase 1	Dusp1	1513	812	0.5 ↓
1415834_at	Dual specificity phosphatase 6	Dusp6	1187	682	0.6 ↓
1417065_at	Early growth response 1	Egr1	14097	7195	0.5 ↓
1436329_at	Early growth response 3	Egr3	2839	1642	0.6 ↓
1428142_at	Ets variant gene 5	Etv5	1139	691	0.6 ↓
1433551_at	Expressed sequence A1427515	A1427515	2067	1196	0.6 ↓
1442025_a_at	Expressed sequence A1467657	A1467657	218	392	1.8 ↑
1442026_at	Expressed sequence A1467657	A1467657	474	941	2.0 ↑
1435940_at	Expressed sequence A1836758	A1836758	1606	2790	1.7 ↑
1450779_at	Fatty acid binding protein 7, brain	Fabp7	685	357	0.5 ↓
1423100_at	FBJ osteosarcoma oncogene	Fos	964	200	0.2 ↓
1437247_at	Fos-like antigen 2	Fosl2	346	88	0.3 ↓

Probe Set Name	Gene Title	Gene Symbol	SS	SP	SS/SP
1428642_at	Frc, fringe-like 1 (Drosophila)	Frc1l	845	517	0.6 ↓
1455444_at	Gamma-aminobutyric acid (GABA-A) receptor, subunit alpha 2	Gabra2	856	506	0.6 ↓
1448397_at	Gap junction membrane channel protein beta 6	Gjb6	661	1176	1.8 ↑
1417859_at	Growth arrest specific 7	Gas7	387	237	0.6 ↓
1435194_at	Heat shock protein 4	Hspa4	1262	699	0.6 ↓
1431182_at	Heat shock protein 8	Hspa8	438	253	0.6 ↓
1455789_x_at	Heat shock protein 8	Hspa8	21704	13291	0.6 ↓
1415899_at	Jun-B oncogene	Junb	799	458	0.6 ↓
1449490_at	Methyl-cpg binding domain protein 4	Mbd4	132	217	1.6 ↑
1422860_at	Neurotensin	Nts	389	220	0.6 ↓
1416505_at	Nuclear receptor subfamily 4, group A, member 1	Nr4a1	3523	1074	0.3 ↓
1449374_at	Pipecolic acid oxidase	Pipox	284	165	0.6 ↓
1418595_at	Plasma membrane associated protein, S3-12	S3-12	268	945	3.5 ↑
1448684_at	Protein phosphatase 1, regulatory (inhibitor) subunit 2	Ppp1r2	4593	7927	1.7 ↑
1419247_at	Regulator of G-protein signaling 2	Rgs2	3148	1891	0.6 ↓
1419248_at	Regulator of G-protein signaling 2	Rgs2	3284	1849	0.6 ↓
1447830_s_at	Regulator of G-protein signaling 2	Rgs2	5817	3077	0.5 ↓
1456339_at	RIKEN cDNA 2810410D24 gene	2810410D24Rik	485	293	0.6 ↓
1428375_at	RIKEN cDNA 4932415G12 gene	4932415G12Rik	467	784	1.7 ↑
1436387_at	RIKEN cDNA C330006P03 gene	C330006P03Rik	1776	412	0.2 ↓
1456280_at	RIKEN cDNA E130314M08 gene	E130314M08Rik	814	480	0.6 ↓
1422660_at	RNA binding motif protein 3	Rbm3	1435	3304	2.3 ↑
1429169_at	RNA binding motif protein 3	Rbm3	248	120	0.5 ↓
1434347_s_at	RUN and FYVE domain-containing 2	Rufy2	127	211	1.7 ↑
1416041_at	Serum/glucocorticoid regulated kinase	Sgk	1649	4070	2.5 ↑
1457373_at	Similar to cadherin 19, type 2 preproprotein (LOC227485), mRNA	---	201	334	1.7 ↑
1424481_s_at	Similar to hypothetical protein FLJ12969 (LOC385421), mRNA	---	203	355	1.7 ↑
1427345_a_at	Sulfotransferase family 1A, phenol-preferring, member 1	Sult1a1	292	863	3.0 ↑
1424330_at	SUMO1/sentrin specific protease 1	Senp1	280	166	0.6 ↓
1416029_at	TGFB inducible early growth response 1	Tiegl	373	184	0.5 ↓
1436094_at	Transcribed sequence with weak similarity to protein pir:S05381 (R.norvegicus) S05381 VGF8a protein precursor - rat	---	6183	2368	0.4 ↓

Probe Set Name	Gene Title	Gene Symbol	SS	SP	SS/SP
1439163_at	Transcribed sequence with weak similarity to protein ref:NP_079268.1 (H.sapiens) hypothetical protein FLJ12547 [Homo sapiens]	---	533	882	1.7
1430153_at	Tropomodulin 2	Tmod2	403	664	1.6
1417374_at	Tubulin, alpha 4	Tuba4	6174	3755	0.6
1423778_at	Ubiquitin specific protease 20	Usp20	201	434	2.2
1419063_at	UDP-glucuronosyltransferase 8	Ugt8	2003	1020	0.5
1420909_at	Vascular endothelial growth factor A	Vegfa	663	326	0.5

Table 6

List of genes modulated in the striatum 24 h after injection of MPTP or paraquat.

Probe Set Name	Gene Title	Gene Symbol	SS	SM	SP	SP/SS	SM/SS
1444564_at	Apolipoprotein D	Apod	334	878	640	1.9	↑ 2.6
1418687_at	Activity regulated cytoskeletal-associated protein	Arc	2281	804	319	0.1	↓ 0.4
1416250_at	B-cell translocation gene 2, anti-proliferative	Btg2	566	337	332	0.6	↓ 0.6
1459372_at	Bhlh-PAS type transcription factor NXF	Nxf	389	161	119	0.3	↓ 0.4
1434369_a_at	Crystallin, alpha B	Cryab	1963	1174	963	0.5	↓ 0.6

Table 7

GO Analysis of biological processes and molecular functions enriched after MPTP or PQ treatment. **GO terms enriched 24 hours after MPTP treatment** (*).

Biological Processes	Size	GiG	%	p Value	Genes
Response to wounding (GO:0009611)	327	17	9.88	0.000001	Clqa, Clqb, Clqc, C4b, Cd44, Gjal, Hbegf, Lcp1, Ly86, Nupr1, Pycard, Prosl, Slpr3, Stat3, Tnfrsf1a, Trf, Vwf
Defense response (GO:0006952)	376	17	9.88	0.000004	Clqa, Clqb, Clqc, C4b, Cd44, Clic1, Feerlg, H2-K1, Irgm1, Ly86, Nupr1, Pycard, Ptpre, Slpr3, Stat3, Tnfrsf1a, Trf
Inflammatory response (GO:0006954)	208	12	6.98	0.000021	Clqa, Clqb, Clqc, C4b, Cd44, Ly86, Nupr1, Pycard, Slpr3, Stat3, Tnfrsf1a, Trf,
Acute inflammatory response (GO:0002526)	73	7	4.07	0.000159	Clqa, Clqb, Clqc, C4b, Nupr1, Stat3, Trf
Positive regulation of response to stimulus (GO:0048584)	169	9	5.23	0.000602	Clqa, Clqb, Clqc, C4b, Feerlg, H2-K1, Ptpre, Tgm2, Tnfrsf1a,
Activation of immune response (GO:0002253)	75	6	3.49	0.001482	Clqa, Clqb, Clqc, C4b, Feerlg, Ptpre
Immune effector process (GO:0002252)	114	7	4.07	0.001717	Clqa, Clqb, Clqc, C4b, Feerlg, Ptpre, Ptx3
Response to metal ion (GO:0010038)	50	5	2.91	0.002295	Ccnd1, Mgp, Mtl, Mtl2, S100a16
Complement activation, classical pathway (GO:0006958)	24	4	2.33	0.002318	Clqa, Clqb, Clqc, C4b
Positive regulation of immune response (GO:0050778)	121	7	4.07	0.002323	Clqa, Clqb, Clqc, C4b, Feerlg, H2-K1, Ptpre
Molecular functions	Size	GiG	GiG%	p Value	Genes
Protein dimerization activity (GO:0046983)	326	12	7.0	0.001	Clqb, Cav2, Cebp, Cebp, Jun, Npas4, Pycard, S100a16, Siml, Stat3, Vwf, Wwtr1
Identical protein binding (GO:0042802)	279	10	5.8	0.003	Clqb, Cav2, Cebp, Cebp, Cldn10, Lcp1, Pycard, S100a16, Vwf, Wwtr1
GO terms enriched 24 hours after PQ treatment (*).					
Molecular Function	Size	GiG	%	p Value	Genes
Protein dimerization activity (GO:0046983)	340	7	11.86	0.001	Fos, Fosl2, Junb, Npas4, Nr4a1, Vegfa, Zbtb16

(*) Only GO terms with p-value less than 0.003 are reported.

Table 8

GSEA analysis of biological processes and molecular functions enriched in MPTP-versus PQ-treated mice.

Biological processes	Size	NES	FDR q-val	Gene List
Negative regulation of apoptosis (GO:0043066)	136	2.08	0.029	Anxa5, Arhgdia, Aven, Bax, Bcl10, Bcl2, Bcl3, Becn1, Bnip1, Ccl2, Cd74, Clcf1, Cryab, Dad1, Ddah2, Gpx1, Hsp90b1, Hspa1b, Hspb1, Ier3, Il2, Krt18, Mc11, Myo18a, Nme2, Notch2, Npm1, Rtn4, Scg2, Sema4d, Sfrp1, Socs3, Sphk1, Tnfrsf1a, Tnfrsf1b, Tnfrsf1c, Tnfrsf1d, Tnfrsf1e, Tnfrsf1f, Tnfrsf1g, Tnfrsf1h, Tnfrsf1i, Tnfrsf1j, Tnfrsf1k, Tnfrsf1l, Tnfrsf1m, Tnfrsf1n, Tnfrsf1o, Tnfrsf1p, Tnfrsf1q, Tnfrsf1r, Tnfrsf1s, Tnfrsf1t, Tnfrsf1u, Tnfrsf1v, Tnfrsf1w, Tnfrsf1x, Tnfrsf1y, Tnfrsf1z, Tnfrsf1aa, Tnfrsf1ab, Tnfrsf1ac, Tnfrsf1ad, Tnfrsf1ae, Tnfrsf1af, Tnfrsf1ag, Tnfrsf1ah, Tnfrsf1ai, Tnfrsf1aj, Tnfrsf1ak, Tnfrsf1al, Tnfrsf1am, Tnfrsf1an, Tnfrsf1ao, Tnfrsf1ap, Tnfrsf1aq, Tnfrsf1ar, Tnfrsf1as, Tnfrsf1at, Tnfrsf1au, Tnfrsf1av, Tnfrsf1aw, Tnfrsf1ax, Tnfrsf1ay, Tnfrsf1az, Tnfrsf1ba, Tnfrsf1bb, Tnfrsf1bc, Tnfrsf1bd, Tnfrsf1be, Tnfrsf1bf, Tnfrsf1bg, Tnfrsf1bh, Tnfrsf1bi, Tnfrsf1bj, Tnfrsf1bk, Tnfrsf1bl, Tnfrsf1bm, Tnfrsf1bn, Tnfrsf1bo, Tnfrsf1bp, Tnfrsf1bq, Tnfrsf1br, Tnfrsf1bs, Tnfrsf1bt, Tnfrsf1bu, Tnfrsf1bv, Tnfrsf1bw, Tnfrsf1bx, Tnfrsf1by, Tnfrsf1bz, Tnfrsf1ca, Tnfrsf1cb, Tnfrsf1cc, Tnfrsf1cd, Tnfrsf1ce, Tnfrsf1cf, Tnfrsf1cg, Tnfrsf1ch, Tnfrsf1ci, Tnfrsf1cj, Tnfrsf1ck, Tnfrsf1cl, Tnfrsf1cm, Tnfrsf1cn, Tnfrsf1co, Tnfrsf1cp, Tnfrsf1cq, Tnfrsf1cr, Tnfrsf1cs, Tnfrsf1ct, Tnfrsf1cu, Tnfrsf1cv, Tnfrsf1cw, Tnfrsf1cx, Tnfrsf1cy, Tnfrsf1cz, Tnfrsf1da, Tnfrsf1db, Tnfrsf1dc, Tnfrsf1dd, Tnfrsf1de, Tnfrsf1df, Tnfrsf1dg, Tnfrsf1dh, Tnfrsf1di, Tnfrsf1dj, Tnfrsf1dk, Tnfrsf1dl, Tnfrsf1dm, Tnfrsf1dn, Tnfrsf1do, Tnfrsf1dp, Tnfrsf1dq, Tnfrsf1dr, Tnfrsf1ds, Tnfrsf1dt, Tnfrsf1du, Tnfrsf1dv, Tnfrsf1dw, Tnfrsf1dx, Tnfrsf1dy, Tnfrsf1dz, Tnfrsf1ea, Tnfrsf1eb, Tnfrsf1ec, Tnfrsf1ed, Tnfrsf1ee, Tnfrsf1ef, Tnfrsf1eg, Tnfrsf1eh, Tnfrsf1ei, Tnfrsf1ej, Tnfrsf1ek, Tnfrsf1el, Tnfrsf1em, Tnfrsf1en, Tnfrsf1eo, Tnfrsf1ep, Tnfrsf1eq, Tnfrsf1er, Tnfrsf1es, Tnfrsf1et, Tnfrsf1eu, Tnfrsf1ev, Tnfrsf1ew, Tnfrsf1ex, Tnfrsf1ey, Tnfrsf1ez, Tnfrsf1fa, Tnfrsf1fb, Tnfrsf1fc, Tnfrsf1fd, Tnfrsf1fe, Tnfrsf1ff, Tnfrsf1fg, Tnfrsf1fh, Tnfrsf1fi, Tnfrsf1fj, Tnfrsf1fk, Tnfrsf1fl, Tnfrsf1fm, Tnfrsf1fn, Tnfrsf1fo, Tnfrsf1fp, Tnfrsf1fq, Tnfrsf1fr, Tnfrsf1fs, Tnfrsf1ft, Tnfrsf1fu, Tnfrsf1fv, Tnfrsf1fw, Tnfrsf1fx, Tnfrsf1fy, Tnfrsf1fz, Tnfrsf1ga, Tnfrsf1gb, Tnfrsf1gc, Tnfrsf1gd, Tnfrsf1ge, Tnfrsf1gf, Tnfrsf1gg, Tnfrsf1gh, Tnfrsf1gi, Tnfrsf1gj, Tnfrsf1gk, Tnfrsf1gl, Tnfrsf1gm, Tnfrsf1gn, Tnfrsf1go, Tnfrsf1gp, Tnfrsf1gq, Tnfrsf1gr, Tnfrsf1gs, Tnfrsf1gt, Tnfrsf1gu, Tnfrsf1gv, Tnfrsf1gw, Tnfrsf1gx, Tnfrsf1gy, Tnfrsf1gz, Tnfrsf1ha, Tnfrsf1hb, Tnfrsf1hc, Tnfrsf1hd, Tnfrsf1he, Tnfrsf1hf, Tnfrsf1hg, Tnfrsf1hi, Tnfrsf1hj, Tnfrsf1hk, Tnfrsf1hl, Tnfrsf1hm, Tnfrsf1hn, Tnfrsf1ho, Tnfrsf1hp, Tnfrsf1hq, Tnfrsf1hr, Tnfrsf1hs, Tnfrsf1ht, Tnfrsf1hu, Tnfrsf1hv, Tnfrsf1hw, Tnfrsf1hx, Tnfrsf1hy, Tnfrsf1hz, Tnfrsf1ia, Tnfrsf1ib, Tnfrsf1ic, Tnfrsf1id, Tnfrsf1ie, Tnfrsf1if, Tnfrsf1ig, Tnfrsf1ih, Tnfrsf1ii, Tnfrsf1ij, Tnfrsf1ik, Tnfrsf1il, Tnfrsf1im, Tnfrsf1in, Tnfrsf1io, Tnfrsf1ip, Tnfrsf1iq, Tnfrsf1ir, Tnfrsf1is, Tnfrsf1it, Tnfrsf1iu, Tnfrsf1iv, Tnfrsf1iw, Tnfrsf1ix, Tnfrsf1iy, Tnfrsf1iz, Tnfrsf1ja, Tnfrsf1jb, Tnfrsf1jc, Tnfrsf1jd, Tnfrsf1je, Tnfrsf1jf, Tnfrsf1jg, Tnfrsf1jh, Tnfrsf1ji, Tnfrsf1jj, Tnfrsf1jk, Tnfrsf1jl, Tnfrsf1jm, Tnfrsf1jn, Tnfrsf1jo, Tnfrsf1jp, Tnfrsf1jq, Tnfrsf1jr, Tnfrsf1js, Tnfrsf1jt, Tnfrsf1ju, Tnfrsf1jv, Tnfrsf1jw, Tnfrsf1jx, Tnfrsf1jy, Tnfrsf1jz, Tnfrsf1ka, Tnfrsf1kb, Tnfrsf1kc, Tnfrsf1kd, Tnfrsf1ke, Tnfrsf1kf, Tnfrsf1kg, Tnfrsf1kh, Tnfrsf1ki, Tnfrsf1kj, Tnfrsf1kk, Tnfrsf1kl, Tnfrsf1km, Tnfrsf1kn, Tnfrsf1ko, Tnfrsf1kp, Tnfrsf1kq, Tnfrsf1kr, Tnfrsf1ks, Tnfrsf1kt, Tnfrsf1ku, Tnfrsf1kv, Tnfrsf1kw, Tnfrsf1kx, Tnfrsf1ky, Tnfrsf1kz, Tnfrsf1la, Tnfrsf1lb, Tnfrsf1lc, Tnfrsf1ld, Tnfrsf1le, Tnfrsf1lf, Tnfrsf1lg, Tnfrsf1lh, Tnfrsf1li, Tnfrsf1lj, Tnfrsf1lk, Tnfrsf1ll, Tnfrsf1lm, Tnfrsf1ln, Tnfrsf1lo, Tnfrsf1lp, Tnfrsf1lq, Tnfrsf1lr, Tnfrsf1ls, Tnfrsf1lt, Tnfrsf1lu, Tnfrsf1lv, Tnfrsf1lw, Tnfrsf1lx, Tnfrsf1ly, Tnfrsf1lz, Tnfrsf1ma, Tnfrsf1mb, Tnfrsf1mc, Tnfrsf1md, Tnfrsf1me, Tnfrsf1mf, Tnfrsf1mg, Tnfrsf1mh, Tnfrsf1mi, Tnfrsf1mj, Tnfrsf1mk, Tnfrsf1ml, Tnfrsf1mm, Tnfrsf1mn, Tnfrsf1mo, Tnfrsf1mp, Tnfrsf1mq, Tnfrsf1mr, Tnfrsf1ms, Tnfrsf1mt, Tnfrsf1mu, Tnfrsf1mv, Tnfrsf1mw, Tnfrsf1mx, Tnfrsf1my, Tnfrsf1mz, Tnfrsf1na, Tnfrsf1nb, Tnfrsf1nc, Tnfrsf1nd, Tnfrsf1ne, Tnfrsf1nf, Tnfrsf1ng, Tnfrsf1nh, Tnfrsf1ni, Tnfrsf1nj, Tnfrsf1nk, Tnfrsf1nl, Tnfrsf1nm, Tnfrsf1nn, Tnfrsf1no, Tnfrsf1np, Tnfrsf1nq, Tnfrsf1nr, Tnfrsf1ns, Tnfrsf1nt, Tnfrsf1nu, Tnfrsf1nv, Tnfrsf1nw, Tnfrsf1nx, Tnfrsf1ny, Tnfrsf1nz, Tnfrsf1oa, Tnfrsf1ob, Tnfrsf1oc, Tnfrsf1od, Tnfrsf1oe, Tnfrsf1of, Tnfrsf1og, Tnfrsf1oh, Tnfrsf1oi, Tnfrsf1oj, Tnfrsf1ok, Tnfrsf1ol, Tnfrsf1om, Tnfrsf1on, Tnfrsf1oo, Tnfrsf1op, Tnfrsf1oq, Tnfrsf1or, Tnfrsf1os, Tnfrsf1ot, Tnfrsf1ou, Tnfrsf1ov, Tnfrsf1ow, Tnfrsf1ox, Tnfrsf1oy, Tnfrsf1oz, Tnfrsf1pa, Tnfrsf1pb, Tnfrsf1pc, Tnfrsf1pd, Tnfrsf1pe, Tnfrsf1pf, Tnfrsf1pg, Tnfrsf1ph, Tnfrsf1pi, Tnfrsf1pj, Tnfrsf1pk, Tnfrsf1pl, Tnfrsf1pm, Tnfrsf1pn, Tnfrsf1po, Tnfrsf1pp, Tnfrsf1pq, Tnfrsf1pr, Tnfrsf1ps, Tnfrsf1pt, Tnfrsf1pu, Tnfrsf1pv, Tnfrsf1pw, Tnfrsf1px, Tnfrsf1py, Tnfrsf1pz, Tnfrsf1qa, Tnfrsf1qb, Tnfrsf1qc, Tnfrsf1qd, Tnfrsf1qe, Tnfrsf1qf, Tnfrsf1qg, Tnfrsf1qh, Tnfrsf1qi, Tnfrsf1qj, Tnfrsf1qk, Tnfrsf1ql, Tnfrsf1qm, Tnfrsf1qn, Tnfrsf1qo, Tnfrsf1qp, Tnfrsf1qq, Tnfrsf1qr, Tnfrsf1qs, Tnfrsf1qt, Tnfrsf1qu, Tnfrsf1qv, Tnfrsf1qw, Tnfrsf1qx, Tnfrsf1qy, Tnfrsf1qz, Tnfrsf1ra, Tnfrsf1rb, Tnfrsf1rc, Tnfrsf1rd, Tnfrsf1re, Tnfrsf1rf, Tnfrsf1rg, Tnfrsf1rh, Tnfrsf1ri, Tnfrsf1rj, Tnfrsf1rk, Tnfrsf1rl, Tnfrsf1rm, Tnfrsf1rn, Tnfrsf1ro, Tnfrsf1rp, Tnfrsf1rq, Tnfrsf1rr, Tnfrsf1rs, Tnfrsf1rt, Tnfrsf1ru, Tnfrsf1rv, Tnfrsf1rw, Tnfrsf1rx, Tnfrsf1ry, Tnfrsf1rz, Tnfrsf1sa, Tnfrsf1sb, Tnfrsf1sc, Tnfrsf1sd, Tnfrsf1se, Tnfrsf1sf, Tnfrsf1sg, Tnfrsf1sh, Tnfrsf1si, Tnfrsf1sj, Tnfrsf1sk, Tnfrsf1sl, Tnfrsf1sm, Tnfrsf1sn, Tnfrsf1so, Tnfrsf1sp, Tnfrsf1sq, Tnfrsf1sr, Tnfrsf1ss, Tnfrsf1st, Tnfrsf1su, Tnfrsf1sv, Tnfrsf1sw, Tnfrsf1sx, Tnfrsf1sy, Tnfrsf1sz, Tnfrsf1ta, Tnfrsf1tb, Tnfrsf1tc, Tnfrsf1td, Tnfrsf1te, Tnfrsf1tf, Tnfrsf1tg, Tnfrsf1th, Tnfrsf1ti, Tnfrsf1tj, Tnfrsf1tk, Tnfrsf1tl, Tnfrsf1tm, Tnfrsf1tn, Tnfrsf1to, Tnfrsf1tp, Tnfrsf1tq, Tnfrsf1tr, Tnfrsf1ts, Tnfrsf1tt, Tnfrsf1tu, Tnfrsf1tv, Tnfrsf1tw, Tnfrsf1tx, Tnfrsf1ty, Tnfrsf1tz, Tnfrsf1ua, Tnfrsf1ub, Tnfrsf1uc, Tnfrsf1ud, Tnfrsf1ue, Tnfrsf1uf, Tnfrsf1ug, Tnfrsf1uh, Tnfrsf1ui, Tnfrsf1uj, Tnfrsf1uk, Tnfrsf1ul, Tnfrsf1um, Tnfrsf1un, Tnfrsf1uo, Tnfrsf1up, Tnfrsf1uq, Tnfrsf1ur, Tnfrsf1us, Tnfrsf1ut, Tnfrsf1uu, Tnfrsf1uv, Tnfrsf1uw, Tnfrsf1ux, Tnfrsf1uy, Tnfrsf1uz, Tnfrsf1va, Tnfrsf1vb, Tnfrsf1vc, Tnfrsf1vd, Tnfrsf1ve, Tnfrsf1vf, Tnfrsf1vg, Tnfrsf1vh, Tnfrsf1vi, Tnfrsf1vj, Tnfrsf1vk, Tnfrsf1vl, Tnfrsf1vm, Tnfrsf1vn, Tnfrsf1vo, Tnfrsf1vp, Tnfrsf1vq, Tnfrsf1vr, Tnfrsf1vs, Tnfrsf1vt, Tnfrsf1vu, Tnfrsf1vv, Tnfrsf1vw, Tnfrsf1vx, Tnfrsf1vy, Tnfrsf1vz, Tnfrsf1wa, Tnfrsf1wb, Tnfrsf1wc, Tnfrsf1wd, Tnfrsf1we, Tnfrsf1wf, Tnfrsf1wg, Tnfrsf1wh, Tnfrsf1wi, Tnfrsf1wj, Tnfrsf1wk, Tnfrsf1wl, Tnfrsf1wm, Tnfrsf1wn, Tnfrsf1wo, Tnfrsf1wp, Tnfrsf1wq, Tnfrsf1wr, Tnfrsf1ws, Tnfrsf1wt, Tnfrsf1wu, Tnfrsf1wv, Tnfrsf1ww, Tnfrsf1wx, Tnfrsf1wy, Tnfrsf1wz, Tnfrsf1xa, Tnfrsf1xb, Tnfrsf1xc, Tnfrsf1xd, Tnfrsf1xe, Tnfrsf1xf, Tnfrsf1xg, Tnfrsf1xh, Tnfrsf1xi, Tnfrsf1xj, Tnfrsf1xk, Tnfrsf1xl, Tnfrsf1xm, Tnfrsf1xn, Tnfrsf1xo, Tnfrsf1xp, Tnfrsf1xq, Tnfrsf1xr, Tnfrsf1xs, Tnfrsf1xt, Tnfrsf1xu, Tnfrsf1xv, Tnfrsf1xw, Tnfrsf1xx, Tnfrsf1xy, Tnfrsf1xz, Tnfrsf1ya, Tnfrsf1yb, Tnfrsf1yc, Tnfrsf1yd, Tnfrsf1ye, Tnfrsf1yf, Tnfrsf1yg, Tnfrsf1yh, Tnfrsf1yi, Tnfrsf1yj, Tnfrsf1yk, Tnfrsf1yl, Tnfrsf1ym, Tnfrsf1yn, Tnfrsf1yo, Tnfrsf1yp, Tnfrsf1yq, Tnfrsf1yr, Tnfrsf1ys, Tnfrsf1yt, Tnfrsf1yu, Tnfrsf1yv, Tnfrsf1yw, Tnfrsf1yx, Tnfrsf1yy, Tnfrsf1yz, Tnfrsf1za, Tnfrsf1zb, Tnfrsf1zc, Tnfrsf1zd, Tnfrsf1ze, Tnfrsf1zf, Tnfrsf1zg, Tnfrsf1zh, Tnfrsf1zi, Tnfrsf1zj, Tnfrsf1zk, Tnfrsf1zl, Tnfrsf1zm, Tnfrsf1zn, Tnfrsf1zo, Tnfrsf1zp, Tnfrsf1zq, Tnfrsf1zr, Tnfrsf1zs, Tnfrsf1zt, Tnfrsf1zu, Tnfrsf1zv, Tnfrsf1zw, Tnfrsf1zx, Tnfrsf1zy, Tnfrsf1zz
Jak stat cascade (GO:0007259)	29	2.08	0.011	Ccl2, Clcf1, Hcls1, Lyn, Nmi, Socs3, Stat3
Positive regulation of signal transduction (GO:0009967)	108	2.07	0.009	Bcl10, Bst2, Casp1, Cdkn1c, Clcf1, Cxhc5, Eef1d, Eef1e1, Fadd, Flna, Golt1b, Hcls1, Hmox1, Il12a, Lgals1, Lgals9, Litaf, Ltbr, Lyn, Myd88, Nek6, Notch2, Ptpcr, Rasgrp4, Rela, Rhoc, Rpl17, Slc44a2, Tfg, Tmem101, Tnfrsf1a, Zdhhc13
Regulation of IKK NFκB cascade (GO:0043122)	76	2.05	0.008	Bcl10, Bst2, Casp1, Cxhc5, Eef1d, Fadd, Flna, Golt1b, Hmox1, Lgals1, Lgals9, Litaf, Ltbr, Myd88, Nek6, Rela, Rhoc, Rpl17, Slc44a2, Tfg, Tmem101, Tnfrsf1a, Zdhhc13
Positive regulation of phosphorylation (GO:0042327)	25	2.02	0.007	Ccnd1, Ccnd2, Clcf1, Hcls1, Il5, Lyn
Regulation of signal transduction (GO:0009966)	197	2.001	0.008	Ambp, Arf6, Bcl10, Bst2, Casp1, Cdkn1c, Clcf1, Cxhc5, Eef1d, Eef1e1, Eid2, Fadd, Flna, Git1, Git2, Golt1b, Grk5, Hcls1, Hmox1, Il12a, Lgals1, Lgals9, Litaf, Ltbr, Lyn, Mdfic, Myd88, Nek6, Notch2, Otx2, Plce1, Ptpcr, Rasgrp4, Rela, Rgs2, Rgs20, Rhoc, Rpl17, Slc44a2, Smad2, Tax1bp3, Tfg, Tgfb1, Tmem101, Tnfrsf1a, Zdhhc13
Molecular functions	Size	NES	FDR q-val	Gene List
Structural constituent of cytoskeleton (GO:0005200)	55	2.09	0.002	Arpc1b, Gfap, Msn, Vim,
Specific RNA POL-II transcription factor activity (GO:0003704)	31	2.01	0.007	Atf4, Fos, Glis3, Notch2, Nr2f1, Sox9, Usf1, Znf202
Protease inhibitor activity (GO:0030414)	32	1.92	0.030	Ambp, Cst7, Pros1, Renbp, Serpinb12, Serping1, Serpinh1, Spink5, Spint2, Timp1
Unfolded protein binding (GO:0051082)	39	1.92	0.023	Cct3, Cct4, Cct6a, Cct7, Cdc37, Chaf1b, Grpel1, Hspa1b, Hspa9, Hspd1, Mkks, Nap114, Npm1, Ppiib, Ppic, Ptges3, Ruvbl2, Spg7, Tapbp, Tcpl
Enzyme inhibitor activity (GO:0004857)	107	1.90	0.026	Abce1, Ambp, Anxa5, Cdc42se1, Cdkn1c, Csn2, Cst7, Dnajc3, Gps1, Gpx1, Pkig, Pros1, Prpsap1, Renbp, Rnh1, Serpinb12, Serping1, Serpinh1, Socs3, Spink5, Spint2, Tesc, Timp1, Tnfrsf1a
Oxidoreductase activity (GO:0016616)	50	1.86	0.041	Akr1b1, Akr1b10, Akr7a2, Gpd1, Hpgd, Hsd3b2, Idh1, Idh3b, Impdh2, Ldha, Me1, Rdh5, Rdh10, Rdh16, Spr, Ugdh
Oxidoreductase activity acting on CHOH groups of donors (GO:0016614)	56	1.82	0.051	Akr1b1, Akr1b10, Akr1c4, Akr7a2, Gpd1, Gpd2, H6pd, Hadha, Hpgd, Hsd17b4, Hsd3b2, Idh1, Idh3b, Impdh2, Ldha, Me1, Me2, Rdh5, Rdh10, Rdh16, Spr, Ugdh
Electron carrier activity (GO:0009055)	71	1.80	0.054	Aifm1, Akr1b1, Akr1c4, Akr7a2, Cox11, Cox5a, Cyb561, Cyb5r3, Cyb5r4, Cyba, Cysc, Dhfr3, Dmgdh, Etfb, Etfh, Etfh, Fdx1, Gpx2, Gsr, Haao, Idh3b, Kmo, Me1, Me2, Mtrr, Ndufa13, Ndufa9, Ndufs1, Ndufs3, Ndufs7, Ndufs8, Nqo1, P4ha2, Por, Rdh16, Sardh, Txndc2, Ugdh
Transferase activity transferring acyl groups (GO:0016746)	52	1.76	0.072	Aanat, Agpat2, Edf1, Hadha, Hat1, Lpcat1, Sat1, Smarce1, Soat1, Taf5, Tgm1, Tgm2, Tgm3, Tgm5